

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: Jan Wadstein *et al.*

Serial No.: 09/410,484

Group No.: 1616

Filed: 09/30/99

Examiner: Arnold, Ernst

Entitled: **Method Of Treating Hypertension And Reducing Serum Lipase Activity**

APPELLANTS' BRIEF

EFS WEB FILED

Appeal Brief - Patents

Commissioner for Patents

P.O. Box 1450

Alexandria, VA 22313-1450

Dear Examiner Ernst:

This Brief is in furtherance of the Notice of Appeal filed May 27, 2008.

Appellants hereby authorize the Commissioner of the Patent and Trademark Office to charge the fees required under § 41.20(b)(2), any required fee for any Petition for Extension of Time, and any other fee for filing this Brief to Attorney Deposit Account No. 50-4302. Please reference Attorney Docket No.: NATNUT-03972 when charging the Attorney Deposit Account.

This Brief contains these items under the following headings and in the order set forth below [37 CFR § 1.192(c)]:

I.	REAL PARTY IN INTEREST	3
II.	RELATED APPEALS AND INTERFERENCES	3
III.	STATUS OF CLAIMS	3
IV.	STATUS OF AMENDMENTS	3
V.	SUMMARY OF CLAIMED SUBJECT MATTER	3
VI.	GROUND OF REJECTION TO BE REVIEWED ON APPEAL	5
VII.	ARGUMENT	6
VIII.	CLAIMS APPENDIX	17
IX.	EVIDENCE APPENDIX	19
X.	RELATED PROCEEDINGS APPENDIX	106
XI.	CONCLUSION.....	107

I. REAL PARTY IN INTEREST

The real party in interest is the Assignee, AkerBiomarine ASA, Oslo, Norway.

II. RELATED APPEALS AND INTERFERENCES

There are no appeals or interferences related to the pending appeal.

III. STATUS OF CLAIMS

Claims 1-18 were filed in the original application. The Claims were subject to a Restriction Requirement mailed December 8, 2000. Applicants elected the Group 1 claims, Claims 1-9, with traverse in a Response dated January 5, 2001. Claims 4, 5, 6, 8 and 10-18 were withdrawn in the Office Action mailed February 20, 2001. Claims 4, 5, and 6 were cancelled in the Response dated November 8, 2004.

Claims 1, 2, 3, 7 and 9 are pending and are being appealed.

Appellants appeal the Final Office Action of January 25, 2008.

The Claims, as they now stand, are set forth in Section VIII. CLAIMS APPENDIX.

IV. STATUS OF AMENDMENTS

All previous amendments have been entered.

V. SUMMARY OF CLAIMED SUBJECT MATTER

Independent claim 1 is drawn to a method of treating hypertension in humans comprising: a) providing a hypertensive human patient in need of hypertension treatment and a composition comprising a safe and effective amount conjugated linoleic acid for treating hypertension; and b) administering said conjugated linoleic acid composition to said human patient so that blood pressure of said human patient is reduced.

This method is described in the specification on page 3, lines 23-27, which provides that that in some embodiments of the invention, a conjugated linoleic acid composition is given to a subject under conditions such that the blood pressure of the subject is reduced. The method is further described in the specification at page 10, lines 5-7, which provides that “the data described below in Example 1 indicates that dietary supplementation with CLA results in a decrease in both systolic and diastolic blood pressure in humans with borderline mild hypertension.” Page 11, lines 8-15, further provides that:

In a typical regimen, an individual will begin the hypertension treatment program or serum lipase activity reduction program by ingesting up to several grams (e.g., 0.1 to 5

grams) of CLA with each meal, and monitoring blood pressure over a period of several months. This CLA may be provided in the form of a pill or as a component of a prepared food product. Once the desired blood pressure or serum lipase activity has been attained, a proper maintenance level can be found by gradually reducing the dose and continuing to monitor blood pressure to assure there is no increase.

Dependent Claim 2 specifies the method of Claim 1 wherein the conjugated linoleic acid composition is a mixture of octadecadienoic acid isomers selected from the group of cis-9, trans-11; cis-9, cis-11; trans-9, cis-11; trans-9, trans-11; cis-10, cis-12; cis-10, trans-12; trans-10, cis-12; trans-10, trans-12 octadecadienoic acid. Octadecadienoic acid is another name for conjugated linoleic acid or CLA. The isomers of conjugated linoleic acid are discussed in detail in the specification at page 8, line 25, to page 9, line 28.

Dependent Claim 3 specifies the method of Claim 1 wherein the conjugated linoleic acid composition consists essentially of octadecadienoic acid isomers selected from 9,11 octadecadienoic acid, 10,12 octadecadienoic acid, and mixtures thereof. These preferred isomers of CLA are discussed in detail in the specification at page 9, lines 13-28.

Dependent Claim 7 specifies the method of Claim 1 wherein the conjugated linoleic acid is administered orally. Administration of CLA is discussed in detail in the specification at page 10, line 28 to page 11, line 15 of the specification.

Dependent Claim 9 of the specification specifies the method of Claim 1 wherein said safe and effective amount of conjugated linoleic acid is about 0.1 grams to 20 grams. Dosages of CLA are discussed in detail in the specification at page 11, lines 8-21.

VI. GROUNDS OF REJECTION TO BE REVIEWED ON APPEAL

There is one grounds of rejection to be reviewed on appeal:

Whether Claims 1-3, 7 and 9 are obvious over Cook et al. (U.S. Pat. No. 5,554,646) in view of Kawamura et al. (Hypertension 1996, 27, 408-413) and Shinitzky (U.S. Pat. No. 4,474,773).

VII. ARGUMENT

A. The Claims are Not Obvious

Claims 1-3, 7 and 9 stand rejected as obvious over Cook et al. (U.S. Pat. No. 5,554,646) in view of Kawamura et al. (Hypertension 1996, 27, 408-413) and Shinitzky (U.S. Pat. No. 4,474,773). The Office has found that:

- Cook et al. disclose a method of reducing body fat comprising the administration of a safe and effective amount of conjugated linoleic acid (Final Office Action mailed Jan. 25, 2008 at 3);
- Kawamura et al. provide a nexus teaching between hypertension, weight loss and decreases in blood pressure and that changes in body weight exhibited significant correlations with blood pressure in hypertensive overweight human patients (Id. at 3);
- Shinitzky et al. teach treating hypertension by administering a composition comprising 5-10% linoleic acid (Id. at 3).

The Office admits that the primary reference, Cook et al., does not teach a method of administering CLA to treat hypertension in humans. Id. at 4. The Office goes on to argue, however, that one of ordinary skill would have been motivated to combine the references to arrive at the claimed invention because:

- Cook et al. teaches a method of reducing body fat;
- Kawamura et al. teach that reduction of weight of hypertensive patients results in a lowering of blood pressure;
- Shinitzky et al. teach that linoleic acid (C18:2, cis-9, cis-12) can be used to treat hypertension; and
- Since conjugated linoleic acid is a mixture of positional and geometric isomers of linoleic acid, then one of skill in the art would immediately envision conjugated linoleic acid in the treatment of hypertension.

In rejecting claims under 35 U.S.C. § 103, the Examiner bears the initial burden of presenting a *prima facie* case of obviousness. A *prima facie* case of obviousness requires the Examiner to cite a combination of references which (a) disclose the elements of the claimed invention, (b) suggests or motivates one of skill in the art to combine those elements to yield the claimed combination, and (c) provides a reasonable expectation of success should the claimed combination be carried out. Failure to establish any one of these three requirements precludes a finding of a *prima facie* case of obviousness, and, without more, entitles Applicant to

allowance of the claims at issue. *See, e.g., Northern Telecom Inc. v. Datapoint Corp.*, 15 USPQ2d 1321, 1323 (Fed. Cir. 1990).

When a *prima facie* case is made (for the record, Applicants do not believe a *prima facie* case has been made here), the burden shifts to the applicant to come forward with evidence and/or argument supporting patentability. *In re Sullivan*, 498 F. 3d 1345, 1351, 81 USPQ2d 1034 (Fed. Cir. 2007) (citing *In re Glaug*, 283 F.3d 1335, 1338 (Fed.Cir.2002)). As held in *Sullivan*:

Rebuttal evidence is “merely a showing of facts supporting the opposite conclusion.” *In re Piasecki*, 745 F.2d 1468, 1472 (Fed.Cir.1984). Evidence rebutting a *prima facie* case of obviousness can include: “evidence of unexpected results,” *Pfizer, Inc. v. Apotex, Inc.*, 480 F.3d 1348, 1369 (Fed.Cir.2007), evidence “that the prior art teaches away from the claimed invention in any material respect,” *In re Peterson*, 315 F.3d 1325, 1331 (Fed.Cir.2003), and evidence of secondary considerations, such as commercial success and long-felt but unresolved needs, *WMS Gaming, Inc. v. Int'l Game Tech.*, 184 F.3d 1339, 1359 (Fed.Cir.1999).

Id. Importantly, when a patent applicant puts forth rebuttal evidence, the Office must consider that evidence. *Id.*, *see also In re Soni*, 54 F.3d 746, 750 (Fed.Cir.1995) (stating that “all evidence of nonobviousness must be considered when assessing patentability”); *In re Sernaker*, 702 F.2d 989, 996 (Fed.Cir.1983) (“If, however, a patent applicant presents evidence relating to these secondary considerations, the board must always consider such evidence in connection with the determination of obviousness.”).

The determination whether an invention would have been obvious under 35 U.S.C. § 103 is a legal conclusion based on underlying findings of fact. *Id.* at 1350 (citing *In re Kotzab*, 217 F.3d 1365, 1369 (Fed.Cir.2000)). With respect to the instant invention, the Office has failed to establish a *prima facie* case evidence because the facts the Office has relied on are scientifically incorrect or inaccurate and, in any event, has failed to considered properly established facts submitted in rebuttal. In particular, the Office has made scientifically incorrect and unsupportable assumptions about the teachings of Cook and Shinitzky and has failed to consider pertinent rebuttal evidence established in the Declarations of Mr. Asgeir Saebo and Dr. Inge Bruheim. Applicants detail these errors below and request that the obviousness rejection be withdrawn because of these errors.

1. Shinitzky does not teach that linoleic acid can be used to treat hypertension

The Office alleges that Shinitzky et al. teaches that linoleic acid (C18:2, cis-9, cis-12) can be used to treat hypertension. This greatly overstates the teachings of Shinitzky et al. to one of skill in the art. In making this finding, the Examiner relies on the claims of Shinitzky et al. Examination of the claims reveals, however, that Shinitzky et al. utilizes a complex lipid mixture that contains linoleic acid and provides no indication that linoleic acid is the active ingredient. This issue is specifically addressed in the Bruheim Declaration, provided in Section IX below at p. 20-76.

The Office specifically refers to Claims 1, 4, and 24 of Shinitzky as providing the teaching that linoleic acid can be used to treat hypertension. Office Action mailed June 26, 2007 at 3; Office Action mailed Jan. 25, 2008 at 6. What those claims teach is that hypertension can be treated by administering a complex mixture comprising “a lipid fraction derived from natural sources (AL), said lipid fraction containing 40-80 weight percent glycerides, 3-5 weight percent cholesterol, 10-30 weight percent lecithin (phosphatidyl choline), 5-15 weight percent phosphatidyl ethanolamine and 2-5 weight percent negatively charged phospholipids, wherein the ratio of unsaturated to saturated fatty acids is at least 1:1” (Claim 1) and that the fatty acid component of this complex mixture comprises “Palmitic acid 35-45%, oleic acid 35-45%, linoleic acid 5-10%, stearic acid 5-7%, palmitoleic acid 2-3%, arachidonic acid 0.2-1%” (Claim 4). *See* Shinitzky et al., Claims 1 and 4 and Bruheim Decl. at ¶2. As can be plainly seen, the invention claimed in claims 1, 4, and 24 utilizes a complex lipid mixture, of which linoleic acid is only a small part. Thus, it is scientifically incorrect and inaccurate to state that Shinitzky et al. teaches that linoleic acid can be used to treat hypertension. This statement by the Office is not factually supported.

The Bruheim Declaration directly addresses this issue. As established by Dr. Bruheim:

- Shinitzky et al. does not teach that linoleic acid can be used to treat hypertension or that among all of the components of the complex mixture, linoleic acid is sufficient to treat hypertension.
- Shinitzky et al. teaches that a complex lipid fraction can be used to treat hypertension.
- This is not the same as teaching that linoleic acid can be used to treat hypertension.
- One of skill in the art would recognize that any of the other components could be responsible for the hypertensive effect or that a combination of the components is necessary.

Bruheim Decl. ¶2 (p. 20 in Section IX below). The Office provides no factual or scientifically based rebuttal of these facts in the Office Action mailed January 25, 2008. The Office argues that “it is the Examiner’s position that each component is treating hypertension because that is what the claim language clearly states. There can be no other interpretation of the claim.” *Id.* at 6. This is simply incorrect. Neither the claims nor the specification of Shinitzky teach that each component of the complex lipid mixture treats hypertension.

The Office goes on to argue that “Shinitzky et al. do not have to recite a method that utilizes solely linoleic acid. The Examiner also notes for the record the open claim language of the instant application.” *Id.* at 7. Apparently, the Office is trying to make an argument based on claim interpretation. Appellants note however, the current claims are limited to the use of conjugated linoleic acid, which is a mixture of octadecadienoic acid isomers such as cis-9, trans-11; cis-9, cis-11; trans-9, cis-11; trans-9, trans-11; cis-10, cis-12; cis-10, trans-12; trans-10, cis-12; trans-10, trans-12 octadecadienoic acid. The claims do not encompass linoleic acid, which, as noted by the Examiner, is c9, c12 octadecadienoic acid. Linoleic acid does not have a conjugated double bond. Since the use of a different chemical entity is being claimed, i.e., the use of conjugated linoleic acid as opposed to non-conjugated linoleic acid, then the scientific issue of whether Shinitzky teaches that each component of the complex lipid preparation can be used to treat hypertension is relevant and is not addressed by interpreting the claims broadly. If a person of skill in the art set out to make a composition to treat hypertension based the information in the Shinitzky claims, that person would have no guidance to use linoleic acid, much less conjugated linoleic acid. This is confirmed by the Bruheim Declaration.

Furthermore, Appellants note another key flaw in the Office’s analysis. Cook et al. teaches that administration of conjugated linoleic acid, and **not** linoleic acid, reduces body fat. Cook et al., column 2, lines 48-67. The Office’s reasoning is that one would be motivated to use CLA to treat hypertension because reductions in body weight cause hypertension to be reduced. If, as taught by Cook et al., linoleic acid does not reduce body weight, then why would a person of skill in the art, having knowledge of Cook, think that the linoleic acid in the complex Shinitzky lipid preparation, could be used to treat hypertension? Thus, Cook et al. *teaches away* from combination with Shinitzky et al. Accordingly, it is improper to combine Cook et al. with Shinitzky in the first place.

2. The Office has not rebutted the fact established by Appellants that linoleic acid and conjugated linoleic acid have different properties

The Office's second error is its failure to consider the evidence presented in the Bruheim Declaration that establishes that the different isomers of linoleic acid have different properties. In the Office Action mailed June 26, 2007 at page 5, the Office argues that "it remains obvious to one of ordinary skill in the art that the method of Cook et al. can lower blood pressure via administration of conjugated linoleic acid because the positional and geometric isomers of linoleic acid would be expected to have similar properties to linoleic acid in the absence of evidence to the contrary." This argument is related to the Office's *In re Dillon* argument on pages 3 and 4 of the Office Action that chemical compounds with similar structures would be expected by those of skill in the art to similar utilities.

However, the Bruheim Declaration directly rebuts the assumption that linoleic acid and conjugated linoleic acid have similar properties. In particular, Dr. Bruheim establishes that:

- Conjugated isomers of conjugated linoleic acid have different biological properties than linoleic acid.
- This was known in the art prior to the filing date of this application and has been substantiated in many publications after the filing date of this application.
- The references attached to the Bruheim Declaration establish that the conjugated isomers of linoleic acid have distinct biological properties as compared to standard linoleic c9,c12 linoleic acid.
- Many of these references utilize corn oil, which comprises c9,c12 linoleic acid, as a control.
- All of the references show that the conjugated linoleic acid isomers have distinct biological properties.
- Dr. Mark Cook, the inventor of the cited 5,554,646 patent, is an author on most of these publications.

Bruheim Declaration ¶3 (p. 21 in Section IX below). Thus, Appellants provided evidence that establishes that conjugated linoleic acid and linoleic acid have different properties.

In the Office Action mailed January 25, 2008 at page 7, the Office argued that these facts are not persuasive because Cook et al. teach that free linoleic acid is converted to conjugated linoleic acid in the animal. (Citing Column 3, lines 52-56 of Cook et al.). That portion of Cook et al. provides that "The linoleic acid is converted to CLA in the animal, probably by microorganisms in the animal's gastrointestinal system (S.F. Chin, J. M. Storkson, W. Liu, K. Albright, and M. W. Pariza, 1994, J. Nutr. 124:694-701). This assertion by the Office is irrelevant and scientifically incorrect as applied to humans. The Chin et al. reference is attached to the Bruheim Declaration. Chin et al. plainly teaches that linoleic acid is converted to

conjugated linoleic acid by rumen microorganisms at p. 2344 Column 2 (p. 65 in Section IX below), which provides that:

Kepler et al. [1966] identified the cis-9, trans-11 CLA isomer as an intermediate in the biohydrogenation of linoleic acid by the rumen bacterium *Butyrivibrio fibrisolvens*. Seemingly, some of the CLA produced in this way escapes conversion to stearic acid, is absorbed from the digestive system, and is subsequently incorporated into tissue lipid (including milk phospholipid).

There is no teaching in Cook et al. or Chin et al. that humans can convert linoleic acid into conjugated linoleic acid which is what the Office is attempting to establish. The conversion occurs in rumen microorganisms which are not present in the human gut. Accordingly, the Office has failed to rebut the evidence of the Appellants that establishes that conjugated linoleic acid has different effects than linoleic acid. The Office argues that “any linoleic acid administered to an animal would be expected to be converted into conjugated linoleic acid and have all of the benefits of action of conjugated linoleic acid as per the teachings of Cook et al.” Office Action mailed January 25, 2008 at 7. This argument is clearly incorrect with respect to humans, which the claims are limited to. Of course, this very point is also established by Cook et al. which teaches that conjugated linoleic acid causes a reduction in body fat while linoleic acid does not. If humans were able to convert linoleic acid into conjugated linoleic acid there would be no need for the claimed invention of Cook which is administration of conjugated linoleic acid to a human for the purpose of reducing body fat.

This scientifically irrelevant argument is the *only* evidence offered by the Office to rebut ¶ 3 of the Bruheim Declaration and the facts contained therein, which are summarized above. Thus, the fact that a person of ordinary skill in the art would recognize that linoleic acid and conjugated linoleic acid have different properties as demonstrated by multiple scientific papers is unrebutted. As a result, the Office’s *In re Dillon* argument is inapplicable and properly rebutted.

3. The Office failed to rebut the facts established in the Saebo Declaration that agents that cause weight loss can cause hypertension and that CLA could be expected to cause hypertension

The third error made by the Office is the failure to consider the rebuttal evidence in the Saebo Declaration, provided in full in Section IX below (p. 77-105), that establishes that 1) it is incorrect to make an assumption that a compound that reduces weight will also cause a reduction in hypertension and 2) the art indicates that CLA would be expected to have a hypertensive effect.

The Saebo Declaration establishes that:

- When a biologically active agent, such as CLA, is administered to a subject there can be a variety of effects.
- Just because CLA causes weight loss does not also mean that it would reduce hypertension.
- A person of skill in the art would not reasonable expect CLA to reduce hypertension for two reasons.
- CLA has been shown to elevate the level of F2-isoprostane. Taylor et al., Conjugated Linoleic Acid Impairs Endothelial Function, Arteriosclerosis, Thrombosis, and Vascular Biology 26(2), 307-312 (2006)(attached to Declaration).
- F2-isoprostanes have a vasoconstrictive effect. Cracowski et al., Cardiovascular pharmacology and physiology of the isoprostanes, Fundamental & Clinical Pharmacology 20(5): 417-427 (2006)(attached to Declaration).
- Taken together, it should be expected that administration of CLA would result in an increase in blood pressure.
- Second, administration of other agents known to be effective for weight loss can result in increased hypertension.
- Ephedrine, a commonly used, biologically active weight loss supplement is one such example.
- As established in Haller and Benowitz, Adverse Cardiovascular and Central Nervous System Events Associated with Dietary Supplements Containing Ephedra Alkaloids, New England J. Med. 343(25):1833-1838 (2000)(attached to Declaration), ephedrine can cause an increase in hypertension.
- The Examiner's argument that it would be obvious to use CLA to decrease hypertension because CLA administration also causes weight loss lacks scientific merit.
- How an agent such as CLA acts in the body is complex.
- Whether CLA causes an increase or decrease in hypertension, or has no effect at all, is determined by a variety of factors that have no relation to weight loss.
- It is not scientifically valid to draw a conclusion that because an agent causes weight loss, it can also be expected to decrease hypertension.
- The references cited by the Examiner contain no data that can be interpreted in this manner.

Saebo Declaration ¶3-6 (p. 78 in Section IX below). The Examiner addressed the Saebo Declaration in the Office Action dated June 26, 2007, at page 5. First, the Office **completely failed** to rebut ¶4 of the Saebo Declaration which establishes that CLA would be expected to

increase, not decrease hypertension because of its effect of elevating levels F2-isoprostane. Second, even though the Office recognized other parts of the Saebo Declaration, the Office's arguments are not relevant and do nothing to rebut the facts established in the Saebo Declaration. In attempting to rebut the Saebo Declaration, the Office argues that:

First, Kawamura et al. provide a nexus teaching between weight loss in hypertensive patients and lowering of blood pressure. Cook et al. provide a method of reducing body fat, which is weight loss. Shinitzky et al. teach linoleic acid in the treatment of hypertension. The art recognizes linoleic acid in a method of treating hypertension. (See also pertinent art below.) Therefore, it remains obvious to one of ordinary skill in the art that the method of Cook et al. can lower blood pressure via administration of conjugated linoleic acid because the positional and geometric isomers of linoleic acid would be expected to have similar properties to linoleic acid in the absence of evidence to the contrary.

Office Action dated June 26, 2007, at page 5 (emphasis by the Office). This "rebuttal" merely restates the arguments already made by the Office in attempting to establish a *prima facie* case of obviousness. There is no rebuttal of the actual facts in the Saebo Declaration. For example, there is no rebuttal of the fact that biologically active agents that cause weight loss can also cause hypertension. This is an important point. When a biologically active agent is used to induce weight loss, the agent can have other effects unrelated to the weight loss. There is a difference between weight loss caused by a biologically active agent and weight loss due to dietary changes or exercise. Likewise, there is no rebuttal of the fact that CLA has been shown to elevate F2-isoprostane which can cause hypertension. Applicants further note that, as discussed above, the Bruheim Declaration established that linoleic acid and conjugated linoleic acid cannot be expected to have similar properties as alleged by the Office.

The Office's primary reasoning supporting its alleged *prima facie* case of obviousness is that Kawamura et al. provides a nexus teaching between hypertension, weight loss and decreases in blood pressure. Office Action mailed Jan. 25, 2008 at 3. Thus, accordingly to the Office's reasoning, any agent that reduces body weight would also decrease hypertension. The Saebo Declaration directly rebuts the Office's incorrect reasoning.

4. Taken together, these facts rebut any prima facie case of obviousness established by the Office

As demonstrated above, to the extent that any *prima facie* case of obviousness was established by the Office, it was rebutted by the factual evidence presented during prosecution. The Office's alleged *prima facie* of obviousness is based on the assertions that:

- Cook et al. teaches a method of reducing body fat;
- Kawamura et al. teach that reduction of weight of hypertensive patients results in a lowering of blood pressure;
- Shinitzky et al. teach that linoleic acid (C18:2, cis-9, cis-12) can be used to treat hypertension; and
- Since conjugated linoleic acid is a mixture of positional and geometric isomers of linoleic acid, then one of skill in the art would immediately envision conjugated linoleic acid in the treatment of hypertension.

First, Applicants have demonstrated above that this is not a proper *prima facie* of obviousness, because Cook et al. actually teaches away from combination with Shinitzky because Cook et al. teaches that CLA is effective to reduce body fat, not linoleic acid. Thus, a person of skill in the art would not believe that the linoleic acid in Shinitzky's complex lipid mixture was effective for reducing body fat or, as argued by the Office, hypertension. Moreover, the facts established by Appellants demonstrate that a person of skill in the art would have expected CLA to have a hypertensive effect, not the effect of reducing hypertension. These facts strongly support the non-obviousness of the invention as claimed under the *Graham* factors.

Second, to the extent that a *prima facie* case of obviousness was presented by the Office, Appellants have presented factual evidence that rebuts the Office's arguments in the form of the Saebo and Bruheim Declarations. Many of the facts in these Declarations were not even addressed by the Office and the Office's arguments regarding the remaining facts are scientifically incorrect and insufficient to rebut the facts contained in the Declarations.

Taken together, these Declarations establish the following:

- Shinitzky et al. does not teach that linoleic acid can be used to treat hypertension or that among all of the components of the complex mixture, linoleic acid is sufficient to treat hypertension. Bruheim Declaration ¶ 2 (p. 20 in Section IX below).
- Shinitzky et al. teaches that a complex lipid fraction can be used to treat hypertension. *Id.*
- This is not the same as teaching that linoleic acid can be used to treat hypertension. *Id.*
- One of skill in the art would recognize that any of the other components could be responsible for the hypertensive effect or that a combination of the components is necessary. *Id.*

- Conjugated isomers of conjugated linoleic acid have different biological properties than linoleic acid. Bruheim Declaration ¶ 3 (p. 21 in Section IX below).
- This was known in the art prior to the filing date of this application and has been substantiated in many publications after the filing date of this application. *Id.*
- The references attached to the Bruheim Declaration establish that the conjugated isomers of linoleic acid have distinct biological properties as compared to standard linoleic c9,c12 linoleic acid. *Id.*
- Many of these references utilize corn oil, which comprises c9,c12 linoleic acid, as a control. *Id.*
- All of the references show that the conjugated linoleic acid isomers have distinct biological properties. *Id.*
- Dr. Mark Cook, the inventor of the cited 5,554,646 patent, is an author on most of these publications. *Id.*
- When a biologically active agent, such as CLA, is administered to a subject there can be a variety of effects. Saebo Declaration ¶ 3 (p. 78 in Section IX below).
- Just because CLA causes weight loss does not also mean that it would reduce hypertension. *Id.*
- A person of skill in the art would not reasonable expect CLA to reduce hypertension for two reasons. *Id.*
- CLA has been shown to elevate the level of F2-isoprostane. Taylor et al., Conjugated Linoleic Acid Impairs Endothelial Function, Arteriosclerosis, Thrombosis, and Vascular Biology 26(2), 307-312 (2006). Saebo Declaration ¶ 4 (p. 78 in Section IX below), *see* p. 94-99 of Evidence Appendix for Taylor et al.
- F2-isoprostanes have a vasoconstrictive effect. Cracowski et al., Cardiovascular pharmacology and physiology of the isoprostanes, Fundamental & Clinical Pharmacology 20(5): 417-427 (2006). *Id.*, *see* p. 81-91 of Evidence Appendix for Cracowski et al.
- Taken together, it should be expected that administration of CLA would result in an increase in blood pressure. Saebo Declaration ¶ 4 (p. 78 in Section IX below).
- Second, administration of other agents known to be effective for weight loss can result in increased hypertension. Saebo Declaration ¶ 5 (p. 78 in Section IX below).
- Ephedrine, a commonly used, biologically active weight loss supplement is one such example. *Id.*

- As established in Haller and Benowitz, Adverse Cardiovascular and Central Nervous System Events Associated with Dietary Supplements Containing Ephedra Alkaloids, New England J. Med. 343(25):1833-1838 (2000), ephedrine can cause an increase in hypertension. *Id.*, see p. 100-105 of Evidence Appendix for Haller and Benowitz.
- The Examiner's argument that it would be obvious to use CLA to decrease hypertension because CLA administration also causes weight loss lacks scientific merit. Saebo Declaration ¶ 5.
- How an agent such as CLA acts in the body is complex. *Id.*
- Whether CLA causes an increase or decrease in hypertension, or has no effect at all, is determined by a variety of factors that have no relation to weight loss. *Id.*
- It is not scientifically valid to draw a conclusion that because an agent causes weight loss, it can also be expected to decrease hypertension. *Id.*
- The references cited by the Examiner contain no data that can be interpreted in this manner. *Id.*

These facts are essentially un rebutted by the Office and rebut any *prima facie* case of obviousness established by the Office. Moreover, Appellants have established that there is no teaching in Cook et al. or Chin et al. that humans can convert linoleic acid into conjugated linoleic acid which is what the Office is attempting to establish which is one of the key rebuttal arguments made by the Office. The conversion occurs in rumen microorganisms which are not present in the human gut. The claimed invention is non-obvious and the rejection should be withdrawn.

VIII. CLAIMS APPENDIX

1. (Previously presented) A method of treating hypertension in humans comprising:
 - a) providing a hypertensive human patient in need of hypertension treatment and a composition comprising a safe and effective amount conjugated linoleic acid for treating hypertension; and
 - b) administering said conjugated linoleic acid composition to said human patient so that blood pressure of said human patient is reduced.
2. (Previously presented) The method of Claim 1 wherein the conjugated linoleic acid composition is a mixture of octadecadienoic acid isomers selected from the group of cis-9, trans-11; cis-9, cis-11; trans-9, cis-11; trans-9, trans-11; cis-10, cis-12; cis-10, trans-12; trans-10, cis-12; trans-10, trans-12 octadecadienoic acid.
3. (Previously presented) The method of Claim 1 wherein the conjugated linoleic acid composition consists essentially of octadecadienoic acid isomers selected from 9,11 octadecadienoic acid, 10,12 octadecadienoic acid, and mixtures thereof.
- 4-6. Cancelled.
7. (Previously presented) The method of Claim 1 wherein the conjugated linoleic acid is administered orally.
8. (withdrawn) The method of Claim 1, wherein the conjugated linoleic acid is provided in a prepared food product.
9. (Previously presented) The method of Claim 1 wherein said safe and effective amount of conjugated linoleic acid is about 0.1 grams to 20 grams.
10. (Withdrawn) A method of reducing serum lipase activity in humans comprising
 - a) providing a subject and a composition comprising a safe and effective amount conjugated linoleic acid; and
 - b) administering said conjugated linoleic acid composition to said subject under conditions such that serum lipase activity of said subjects is reduced.

11. (Withdrawn) The method of Claim 10 wherein the conjugated linoleic acid composition is a mixture of octadecadienoic acid isomers selected from the group of cis-9, trans-11; cis-9, cis-11; trans-9, cis-11; trans-9, trans-11; cis-10, cis-12; cis-10, trans-12; trans-10, cis-12; trans-10, trans-12 octadecadienoic acid.

12. (Withdrawn) The method of Claim 10 wherein the conjugated linoleic acid composition consists essentially of octadecadienoic acid isomers selected from 9,11 octadecadienoic acid, 10,12 octadecadienoic acid, and mixtures thereof.

13. (Withdrawn) The method of Claim 10 wherein said conjugated linoleic acid composition comprises esters of conjugated linoleic acid.

14. (Withdrawn) The method of Claim 13 wherein said esters are selected from methyl esters and ethyl esters.

15. (Withdrawn) The method of Claim 10 wherein said conjugated linoleic acid composition comprises triglycerides including at least one conjugated linoleic acid at the SN-1, SN-2, or SN-3 position of said triglycerides.

16. (Withdrawn) The method of Claim 10 wherein the conjugated linoleic acid is administered orally.

17. (Withdrawn) The method of Claim 10 wherein the conjugated linoleic acid is provided in a prepared food product.

18. (Withdrawn) The method of Claim 10 wherein said safe and effective amount of conjugated linoleic acid is about 0.1 to 20 grams.

IX. EVIDENCE APPENDIX

This evidence appendix contains the Declarations of Dr. Inge Bruheim and Mr. Asgeir Saebo. The Bruheim Declaration was entered by the Examiner in the Office Action mailed January 25, 2008. The Saebo Declaration was entered by the Examiner in the Office Action mailed September 26, 2007.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: Jan Wadstein *et al.*
Serial No.: 09/410,484 Group No.: 1614
Filed: 09/30/99 Examiner: Webman
Entitled: **Method Of Treating Hypertension And Reducing Serum Lipase Activity**

Declaration of Dr. Inge Bruheim

EFS WEB-FILED

Assistant Commissioner for Patents
Washington, D.C. 20231

I, Dr. Inge Bruheim, state as follows:

1. My present position is Senior Research Scientist, Aker Biomarine AS, the current owner of the instant patent application. I have a Ph.D. in Chemistry from the University of Oslo (Norway) and more than 4 years of industry experience in chemical research.

2. The Examiner states on page 3 of the Office Action that Shinitzky et al. teaches that linoleic acid (C18:2, cis-9,cis-12) can be used to treat hypertension. One of skill in the art would not interpret Shinitzky as providing this teaching. The Examiner specifically refers to Claims 1, 4, and 24 of Shinitzky as providing the teaching that linoleic acid can be used to treat hypertension. What those claims teach is that hypertension can be treated by administering a complex mixture comprising "a lipid fraction derived from natural sources (AL), said lipid fraction containing 40-80 weight percent glycerides, 3-5 weight percent cholesterol, 10-30 weight percent lecithin (phosphatidyl choline), 5-15 weight percent phosphatidyl ethanolamine and 2-5 weight percent negatively charged phospholipids, wherein the ratio of unsaturated to saturated fatty acids is at least 1:1" (Claim 1) and that the fatty acid component of this complex mixture comprises "Palmitic acid 35-45%, oleic acid 35-45%, linoleic acid 5-10%, stearic acid 5-7%, palmitoleic acid 2-3%, arachidonic acid 0.2-1%" (Claim 4). Shinitzky et al. does not teach that linoleic acid can be used to treat hypertension or that among all of the components of the

complex mixture, linoleic acid is sufficient to treat hypertension. Shinitzky et al. teaches that complex lipid fraction can be used to treat hypertension. This is not the same as teaching that linoleic acid can be used to treat hypertension. Indeed, one of skill in the art would recognize that any of the other components could be responsible for the hypertensive effect or that a combination of the components is necessary.

3. The Examiner further states the positional and geometric isomers of linoleic acid would be expected to have similar properties to linoleic acid in the absence of evidence to the contrary. Isomers of conjugated linoleic acid have different biological properties than linoleic acid. This was known in the art prior to the filing date of this application and has been substantiated in many publications after the filing date of this application. I have included several references, which are attached, that establish that the conjugated isomers of linoleic acid have distinct biological properties as compared to standard linoleic c9,c12 linoleic acid. Many of these references utilize corn oil, which comprises c9,c12 linoleic acid, as a control. All of the references show that the conjugated linoleic acid isomers have distinct biological properties. Furthermore, Dr. Mark Cook, the inventor of the cited 5,554,646 patent, is an author on most of these publications.

Yang M, Cook ME. Dietary conjugated linoleic acid decreased cachexia, macrophage tumor necrosis factor- α production, and modifies splenocyte cytokine production. *Exp Biol Med* (Maywood). 2003 Jan;228(1):51-8.

Li G, Dong B, Butz DE, Park Y, Pariza MW, Cook ME. NF- κ B independent inhibition of lipopolysaccharide-induced cyclooxygenase by a conjugated linoleic acid cognate, conjugated nonadecadienoic acid. *Biochim Biophys Acta*. 2006 Sep;1761(9):969-72.

Park Y, Storkson JM, Liu W, Albright KJ, Cook ME, Pariza MW. Structure-activity relationship of conjugated linoleic acid and its cognates in inhibiting heparin-releasable lipoprotein lipase and glycerol release from fully differentiated 3T3-L1 adipocytes. *J Nutr Biochem*. 2004 Sep;15(9):561-8.

Park Y, Storkson JM, Ntambi JM, Cook ME, Sih CJ, Pariza MW. Inhibition of hepatic stearoyl-CoA desaturase activity by trans-10, cis-12 conjugated linoleic acid and its derivatives. *Biochim Biophys Acta*. 2000 Jul 19;1486(2-3):285-92.

Park Y, Albright KJ, Liu W, Storkson JM, Cook ME, Pariza MW. Effect of conjugated linoleic

PATENT

U.S. Appln. Ser. No.: 09/410,484
Attorney Docket No. NATNUT-03972

acid on body composition in mice. *Lipids*, 1997 Aug;32(8):853-8.

Chin SF, Storkson JM, Albright KJ, Cook ME, Pariza MW. Conjugated linoleic acid is a growth factor for rats as shown by enhanced weight gain and improved feed efficiency. *J Nutr*. 1994 Dec;124(12):2344-9.

Miller CC, Park Y, Pariza MW, Cook ME. Feeding conjugated linoleic acid to animals partially overcomes catabolic responses due to endotoxin injection. *Biochem Biophys Res Commun*. 1994 Feb 15;198(3):1107-12.

4. I further declare that all statement made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.



Dr. Inge Bruheim

Date: 23/10 - 2007

FEEDING CONJUGATED LINOLEIC ACID TO ANIMALS PARTIALLY
OVERCOMES CATABOLIC RESPONSES DUE TO ENDOTOXIN INJECTION

C.C. Miller, Y. Park¹, M.W. Pariza¹, and M. E. Cook²

Poultry Science Dept., Food Research Institute,
U.W. Madison, Madison, WI 53706

Received December 25, 1993

Summary: The ability of conjugated linoleic acid to prevent endotoxin-induced growth suppression was examined. Mice fed a basal diet or diet with 0.5% fish oil lost twice as much body weight after endotoxin injection than mice fed conjugated linoleic acid. By 72 hours post injection, mice fed conjugated linoleic acid had body weights similar to vehicle injected controls; however, body weights of basal and fish oil fed mice injected with endotoxin were reduced. Conjugated linoleic acid prevented anorexia from endotoxin injection. Splenocyte blastogenesis was increased by conjugated linoleic acid.

© 1994 Academic Press, Inc.

Recently, nonessential fatty acids have been shown to modulate acute inflammatory and immune responses possibly by changing cellular fatty acid profiles (1). The ability of two major fatty acid components of fish oil, eicosapentaenoate (EPA) and docosahexaenoate (DHA), to modulate the synthesis of arachidonate and its metabolites was recognized as one mechanism of immune regulation by fatty acids. EPA has been shown to depress arachidonic acid derived prostaglandin E₂ (PGE₂) synthesis in peritoneal leukocytes from rats (2). Unlike corn oil-fed rats, fish oil-fed rats did not exhibit the characteristic depression in food intake when injected with interleukin-1 (IL-1) (3). This phenomena has been partially attributed to the ability of fish oil to block or alter the cyclooxygenase pathway thereby decreasing PGE₂ synthesis and altering the ability of PGE₂ to negatively feed back on IL-1 synthesis.

CLA (conjugated dienoic derivative of linoleic acid) was shown to be a naturally occurring substance in food which has an anticarcinogenic effect on dimethylbenz(a)anthracene induced

²To whom correspondence should be addressed.

mammary tumors (4). Recent studies (5) have shown that CLA decreases arachidonic acid content of select tissues. It was postulated that like fish oil, CLA may exert an effect on the immune system by altering product formation of either the cyclooxygenase or lipoxygenase pathway. The purpose of the following investigation was to determine the effect of CLA on endotoxin induced growth inhibition and food intake depression. Fatty acid levels in tissue and *in vitro* lymphocyte proliferation was examined.

Materials and methods

Preparation of CLA and purity determination: CLA was prepared from linoleic acid by alkali isomerization as previously described (6, 7). Purity was determined to exceed 97%. CLA was stored in an argon atmosphere at -20°C.

Tissue fatty acid determination: Tissue fatty acid levels were determined by the following procedure. Total fat was extracted by chloroform:methanol (2:1, v/v) as described by Polch et al (8). Fatty acid methyl esters were prepared by reaction with 4% HCL in methanol for 20 minutes at 60 C (9) and extracted with hexane. Tridecanoic acid methyl ester was used as an internal standard and fatty acid methyl esters were identified by comparison with standards using gas chromatography (Hewlett Packard 5890 series II) (7).

Mouse experimental protocol: Three-week old mixed-sex mice (U.W. Madison Dairy Science Dept. outbred colony) (4 per group and 3 groups per dietary treatment) were fed a semi-purified basal diet (containing 2.5% linoleic acid) (Harlan Teklad, Madison, WI) or the basal diet containing either 0.5% added Menhaden fish oil with 25-30% omega-3 fatty acids as triglycerides (Sigma Chemical Co., St. Louis, MO) or 0.5% added CLA. All fatty acids were mixed into the basal diet following the protocol of Frische and Johnston (10) to minimize autooxidation. At the end of 15 consecutive feeding days, mice were weighed and i.p. injected with lipopolysaccharide (E. Coli 055:B5, Sigma Chemical Co., St. Louis, MO) at 1 mg/kg body weight in sterile HEPES buffer (Sigma Chemical Co., St. Louis, MO) or HEPES buffer alone. All mice were weighed 3, 8, 24, 48 and 72 hours post injection, and feed intake was determined at 3, 8 and 24 hours post injection.

At sacrifice, spleens were excised and single cell suspensions were made by teasing the spleen apart in calcium and magnesium free buffer (pH 7.0) and passing through an 18-gauge needle. Cells were then centrifuged through Histopaque 1077 (density 1.077) (Sigma Chemical Co., St. Louis, MO) and the buffy layer containing mononuclear cells was collected. Mononuclear cells were counted on a hemocytometer and tested for viability using trypan blue exclusion. Blastogenesis was measured following the procedure of Hughes et al (11). Results are expressed as a DPM index (disintegrations per minute in stimulated sample cells/ DPM in unstimulated sample cells).

Rat and chick experimental protocol: CLA (0.5% mixed into a semi-purified rat diet (Harlan Teklad, Madison, WI) or a standard chick mash) was fed to 20 rats (Harlan Sprague Dawley, Madison, WI) and 24 chicks (Leghorn X New Hampshire, Poultry Research Laboratory, Madison, WI) for 14 and 7 consecutive days, respectively. Control animals were fed their respective basal diets. After the feeding period, animals were injected i.p. with

endotoxin (see mouse protocol) and body weight losses due to the injection were monitored.

Statistics: Data were subjected to analysis of variance using the general linear models procedure of the Statistical Analysis System (12). Mean differences were determined using the least significant difference method.

Results

Feeding CLA to mice prevented the severe body weight loss caused by endotoxin injection as observed in mice fed the basal diet (figure 1). Fish oil (.5% of diet) was ineffective in preventing endotoxin-induced weight loss. Within the first 12 hours after endotoxin injection, body weight loss was equal among treatment groups. However, by 24 hours after injection, body weight losses in basal and fish oil fed mice were twice that of CLA fed mice. Return to pre-injection body weight also appeared to be faster in the groups fed CLA. In a similar manner, twenty-four hours post endotoxin injection, the reduction in body weight gain in CLA fed chicks (+1 g/24 hrs vs. +3 g/24 hrs for noninjected controls) and rats (-8 g/24 hrs vs +9 g/24 hrs for noninjected controls) was significantly less than endotoxin injected chicks (-6 g/24 hrs) and rats (-15 g/24 hrs) fed their respective basal diets.

Mice injected with buffer consumed significantly more feed at all time periods measured than mice fed either the basal diet or basal diet plus fish oil and injected with endotoxin (figure 2). The latter two groups consumed no feed during the 24 hour period after injection. However, mice fed CLA also consumed significantly more feed than mice fed fish oil or the basal diet respectively. Mice from all treatments which were injected with




Figure 1. Change in body weight due to endotoxin injection in mice consuming diets containing CLA or fish oil.

*Not different from CLA or fish oil fed and control inject.

abc Means with different letters are significantly different ($p < 0.05$).




Figure 2. Effect of endotoxin injection on cumulative feed intake in mice.

ab Means with different letters are significantly different ($p < .05$).

buffer had feed consumptions and body weight gains which did not differ.

Mice fed CLA had a 1.5 fold increase in PHA-P induced spleen lymphocyte blastogenesis over basal fed mice (figure 3). CLA fed mice also displayed greater responses than fish oil fed mice, although this difference was not significant.

The muscle content of CLA was increased in rats (6.25 vs. .94 mg/g fat; $p < .05$) and chicks (4.07 vs. not detectable; $p < .05$) fed CLA relative to basal fed controls. Muscle arachidonic acid levels were decreased in CLA fed animals relative to the basal fed control animals (60.8 vs. 39.9 mg/g fat in rats and 43.6 vs. 38.8 mg/g fat in chicks).




Figure 3. Effect of feeding fish oil or CLA on PHA-P* induced spleen lymphocyte blastogenesis in mice.

*PHA-P level = .78 μ g/500,000 cells.

**Significantly greater than control ($p < .08$).

***DPM sample/DPM control.

Discussion

Klasing et al (13) showed that endotoxin induced weight loss was the result of not only reduced food intake, but also reduced fractional protein synthesis rate in the gastrocnemius and increased skeletal muscle degradation. Likewise, the ability of animals fed CLA to maintain body weight gain during endotoxin challenge may be due to factors other than maintenance of feed intake. Endotoxin injection has been shown to enhance IL-1 production, and semi-purified IL-1 was capable of decreasing rate of gain and feed intake to the same extent as endotoxin injection (13). Recently it has been demonstrated that feed intake depression associated with immune stimulation via IL-1 injection can be prevented by feeding 8% fish oil containing omega-3 fatty acids (3). Fish oils added in human diets increased cell membrane EPA (an n-3 fatty acid) and decreased cell membrane arachidonic acid (an n-6 fatty acid). The change in cell membrane fatty acids was thought to alter prostaglandin production and thus prevented body weight loss due to IL-1 injection (3). It was also thought that a depression of PGE₂ at the skeletal muscle level depressed protein degradation during IL-1 stimulation (14) since IL-1 was shown to enhance muscle catabolism by inducing PGE₂ production (15). It is possible that CLA acted through a similar mechanism and thereby altered cell membrane fatty acids and depressed PGE₂ production. Indeed CLA decreased skeletal muscle arachidonic acid levels. Thus animals maintained growth during immune stimulation.

Michel et al (16) recently reported that CLA stimulated mitogen-induced lymphocyte blastogenesis, cytotoxic activity and macrophage killing ability. In agreement with Michel and co-workers, we also have shown that CLA enhances PHA-P induced lymphocyte blastogenesis relative to basal fed controls.

Acknowledgment

Research was partially supported by the College of Agriculture and Life Sciences at the University of Wisconsin.

References

1. Lefkowitz, J.B., Morrison, A., Lee, V., and Rogers, M. (1990) *J. Immunology*. 145, 1523-1529.
2. Endres, S., Ghorbani, R., Kelley, V., Georgilis, K., Lonnemann, G., VanderMeer, J.W.M., Cannon, J.G., Rogers, T.S., Klempner, M.S., Weber, P.C., Schaefer, E.J., Wolff, S.M., Dinarello, C.A. (1989) *N. Engl. J. Med.* 320, 265-271.

3. Hellerstein, M.K., Meydani, S.N., Meydani, M., Wu, K., and Dinarello, C.A. (1989) *J. Clin. Invest.* 84, 228-235.
4. Ip, C., Chin, S.F., Scimeca, J.A., and Pariza, M. (1991) *Cancer Res.* 51, 6118-6124.
5. Cook, M.E., Miller, C.C., Park, Y., and Pariza, M. (1993) *Poultry Sci.* 72, 1301-1305.
6. Chin, S.F., Storkson, J., Ha, Y.L., and Pariza, M.W. (1992) *Food Comp. Anal.* 5, 185-197.
7. Ha, Y.L., Storkson, J. and Pariza, M. (1990) *Cancer Res.* 50, 1097-1101.
8. Folch, J., Lees, M., and Sloane-Stanley, G.H. (1957) *J. Biol. Chem.* 226, 497-509.
9. Carreau, J.P., and Dubaco, J.P. (1978) *J. Chromatograph.* 151, 384-390.
10. Frische, K.L., Johnston, P.V. (1988) *J. Nutr.* 118, 425-426.
11. Hughes, B.J., Hsich, G.C., Jarvis, B.B., and Sharma, R.P. (1989) *Arch. Environ. Contam. Toxicol.* 18, 388-395.
12. SAS Institute Inc. (1985) *SAS Users Guide: Statistics*, pp. 139-199. SAS Institute Inc., Cary, NC.
13. Klasing, K.C., Laurin, D.E., Peng, R.K., and Fry, D.M. (1987) *J. Nutr.* 117, 1629-1637.
14. Rodemann, H.P., and Goldberg, A.L. (1982) *J. Biol. Chem.* 257, 1632-1638.
15. Dinarello, C.A. (1984) *Infect. Dis.* 6, 51-95.
16. Michel, J.J., Chew, B.P., Schultz, T.D., Wong, T.S., and Magnuson, N.S. (1992) *Fed. Am. Soc. Exp. Biol. J.* 6, A1102.

Dietary Conjugated Linoleic Acid Decreased Cachexia, Macrophage Tumor Necrosis Factor- α Production, and Modifies Splenocyte Cytokines Production¹

MINGDER YANG AND MARK E. COOK²

Department of Animal Sciences, University of Wisconsin, Madison, WI 53706

The effect of conjugated linoleic acid (CLA) on macrophage functions were studied *in vitro*, *in vivo*, and *ex vivo*. In RAW macrophage cell line, CLA (mixed isomers) was shown to inhibit lipopolysaccharide (LPS)-stimulated tumor necrosis factor- α (TNF- α) production. Two CLA isomers, c9,t11 and t10,c12, were tested on RAW cells and it was found that the c9,t11 was the isomer responsible for the inhibition of LPS-induced TNF- α production. BALB/c mice were used to determine the effect of dietary CLA on body weight wasting and feed intake after LPS injection. CLA was protective against LPS-induced body weight wasting and anorexia. Plasma TNF- α levels after LPS injection were lower in the CLA group compared with the corn oil-fed control group 2 hr post-LPS injection. In a separate experiment, 30 mice were fed a CLA-supplemented diet or a corn oil-supplemented diet for 6 weeks and peritoneal resident macrophages were obtained for measuring TNF- α and nitric oxide production after *in vitro* exposure to interferon- γ (IFN- γ) and/or LPS. TNF- α production was not found to be different in peritoneal macrophages from mice fed the dietary treatments, but less nitric oxide was produced in macrophages from CLA-fed mice upon stimulation when compared with macrophages from control-fed mice. Splenocytes were also collected from the mice fed the dietary treatments and stimulated to produce cytokines in culture. Supernatant was used to run cytokine enzyme-linked immunosorbent assays. Interleukin-4 (IL-4) was decreased in CLA-fed mice when splenocytes were stimulated with concanavalin A (Con A) for 44 hr; however, IL-2 and the IL-2-to-IL-4 ratio were elevated. *Exp Biol Med* 228:51–58, 2003

Key words: CLA; cachexia; macrophage; cytokines

A biological important role of conjugated linoleic acid (CLA) as an anticarcinogen was first reported by Pariza and Hargraves (1). It was later found that CLA had physiological properties that differed from LA. Synthetic CLA is a mixture of isomers, and it is difficult and costly to purify pure isomers, hence, until recently, most reports on the physiological effects of CLA have used mixed isomers (predominantly c9,t11 and t10,c12). Among these studies, CLA has been shown to decrease carcinogenesis (2, 3), decrease atherosclerosis (4, 5), and change body composition by increasing body protein and water content and decreasing body fat (6–8). In addition to the above-mentioned properties, CLA also modulates the immune system by increasing lymphocyte blastogenesis, lymphocyte cytotoxic activity, and macrophage killing ability, as well as protecting against end-stage body wasting in autoimmune disease (9–12). Other fatty acids also affect immunity as well (13–15).

It has been shown that several dietary oils modulate macrophage function. For example, dietary fish oil decreased interleukin-1 (IL-1) release by peritoneal macrophages when compared with corn oil-fed controls (16). Macrophages are believed to be the principal sources of tumor necrosis factor (TNF)- α produced *in vivo*, and lipopolysaccharide (LPS) is the most potent stimulus of macrophages for TNF- α production. The activity of TNF- α was originally shown to kill tumor cells, but it also has a profound effect in causing body wasting, or cachexia. Direct infusion of TNF- α into rats has been shown to promote muscle degradation (17). Our laboratory has shown that feeding CLA to chicks reduces LPS-induced body weight wasting and feed intake (12, 18). Because macrophages are immune cells known to produce TNF- α upon LPS stimulation, the effect of CLA at inhibiting LPS-induced wasting could be due to decreased TNF- α production by macrophages.

T lymphocytes play an important role in the immune system. Upon activation, naive T cells (Th0) differentiate into either T helper 1 (Th1) or Th2 effector cells as they secrete different cytokines and mediate very different im-

¹ We would like to thank BASF (Aktiengesellschaft, Ludwigshafen 67056 Germany) for partial support of this research. This work was also partly supported by the Wisconsin Alumni Research Foundation and by the University of Wisconsin-Madison.

² To whom requests for reprints should be addressed at Department of Animal Sciences, University of Wisconsin-Madison, 1675 Observatory Drive, Room 1064, Madison, WI 53706. E-mail: mcook@facstaff.wisc.edu

Received March 19, 2002.

Accepted August 14, 2002.

1535-3702/03/2281-0051\$15.00

Copyright © 2003 by the Society for Experimental Biology and Medicine

mune responses (19). Th1 type responses are primarily cell-mediated immunity and inflammation, and Th2 cells and cytokines mediate humoral immunity. Upon activation, development of Th0 cells into either Th1 or Th2 cells depends on the local cytokine milieu. For example, in the presence of IL-12, Th1 clone development is preferred (20, 21). On the other hand, IL-4 is indispensable for Th0 cells to polarize into Th2 clones (22, 23). Moreover, Th1 and Th2 responses are cross-regulated. For example, IFN- γ , a cytokine produced by Th1 cells, inhibits IL-4 production and suppresses Th2 development (24, 25). Conversely, IL-4 and IL-10 produced by Th2 cells block differentiation of Th0 to Th1 (26, 27). Sugano *et al.* (11) showed dietary CLA increased immunoglobulin (Ig) A, IgG, and IgM in both rat serum and cultured lymph node cells, whereas IgE was reduced. This work implied that CLA may promote Th1 cytokine and inhibit Th2 cytokine production, as Ig class switch from IgG to IgE would not occur without IL-4 or IL-13, both which are potent Th2 cytokines (28, 29). In a guinea pig asthma model, allergen-induced trachea histamine production was reduced in CLA-fed guinea pigs (30). IgE-induced histamine release was enhanced by IL-5, a Th2 cytokine (31). It was our interest to test the effect of CLA on the helper T cell cytokines profile modification as suggested by previous studies.

Because macrophages are very sensitive to LPS stimulation, it seemed that macrophages can be a potential target of CLA to decrease LPS-induced responses. In this study, we investigated the effect of CLA on macrophage TNF- α production, as it was hypothesized that CLA inhibits TNF- α production by macrophages. In addition, splenocyte cytokines IL-2 and IL-4 were also measured.

Material and Methods

CLA. CLA used in the *in vivo* and *ex vivo* feeding trials was obtained from Natural Lipids Inc. (Hovdebygd, Norway) and contained approximately 90% CLA (CLA-90) with the following C18:2 conjugated isomer distribution: 43.5% t10,c12; 41.9% c9,t11 and t9,c11; 1.5% t9,t11 and t10,t12; 0.9% c9,c11; and 0.9% c10,c12. Other fatty acids in CLA-90 were 5.6% oleate, 1.4% palmitate, 0.5% linoleate, 0.4% stearate, and 3.4% unidentified compounds. For *in vitro* studies, LA was purchased from Nu-Check Prep (>99% pure; Elysian, MN). The c9,t11 CLA isomer was purchased from Matreya Inc. (Pleasant Gap, PA). The c9,t11 CLA isomer was 96.3% pure and had 2.6% of t9,t11/t10,t12 and 1.1% of other CLA isomers. The t10,c12 isomer was from Natural Lipids, and it had 92.8% of t10,c12; 1.6% of c9,t11; 1.2% of t9,t11/t10,t12; and 2.8% of other CLA isomers.

Macrophages. RAW 264.7 cells were a gift from M.W. Pariza (Food Research Institute, University of Wisconsin, Madison, WI). Cells were cultured in Dulbecco's modified Eagle's medium (DMEM) media (Life Technologies, Gaithersburg, MD) supplemented with 5% fetal bovine serum (FBS) and 1% antibiotics solution (A9909; Sigma,

St. Louis, MO). Macrophages were plated in the density of 100,000 cells/well in a 24-well plate for 24 hr, followed by culture in fresh media containing fatty acid-albumin complex. The final fatty acid concentration was 100 μ M. Cells were cultured with fatty acid-albumin complex for 24 hr. Fresh media (containing corresponding fatty acid-albumin complex) containing 500 ng/ml LPS (from *E. coli*, serotype 055:B5; L-4005; Sigma) in phosphorus-buffered saline (PBS) was then added for 16 hr to stimulate TNF- α secretion. Control cells were treated with PBS. Media was then collected, frozen, and subsequently analyzed for TNF- α as described below. Cell viability was checked under microscope and by total nonwashable protein measurement (cell protein concentration was measured after removing supernatants for TNF- α analysis and washing cells with PBS three times) to confirm that viability of macrophages cultured under the 100 μ M fatty acid-albumin complex did not differ from macrophages cultured with albumin alone.

Preparation of Fatty Acid-Albumin Complex. To make fatty acid-albumin complexes for treating macrophages *in vitro*, 2.8 mg of free fatty acid, already dissolved in 0.5 ml of KOH (0.1 M), was transferred to a scintillation vial, and 4 ml of 2.5 mM bovine albumin PBS solution (LPS free, A-8806; Sigma) was added and gassed with nitrogen. The vial was then wrapped with foil and refrigerated overnight. The pH was then adjusted to 7.2 using 0.1 M NaOH solution, and the volume was brought to 5 ml with PBS. The solution was filtered using a 0.22- μ m syringe filter for use in cell culture. The fatty acid concentration of the preparation was 2 mM.

Diet. A semipurified powdered diet was purchased from Harlan-Teklad (TD94060, 99% basal mix; Madison, WI). The diet had 5% corn oil and it was supplemented with either 0.5% CLA or corn oil and 0.5% sugar. Hence, the control diet had 5.5% corn oil, and the CLA diet had 5% corn oil and 0.5% CLA (10). Fresh diets were prepared every other week and were kept refrigerated. Fresh diet was provided to mice three times a week. Both diets and water were provided *ad libitum*.

Animals. Weanling BALB/c mice were purchased from Harlan-Sprague Dawley (Indianapolis, IN). In experiment one, 12 mice were immediately divided into two groups of six and were fed either CLA or the control diet. Three mice were housed together in a shoebox cage during dietary treatment and were then individually caged right before LPS injection (see below). In the *ex vivo* study, 30 mice were used in the study with 15 mice who were fed the control diet and 15 mice who were fed the CLA diet. Three mice were housed per cage in a temperature- and humidity-controlled room with a 12:12-hr light:dark cycle. The protocol for animal care and use was approved by the institutional animal care and use committee at the College of Agriculture and Life Sciences, University of Wisconsin-Madison.

In Vivo Mouse Trial Treatment. After 6 weeks on the dietary treatments, three mice from each dietary group

were injected intraperitoneally with LPS (0.1 mg/ml in sterile PBS, 1 ml LPS solution/100 g of body weight) and the remaining three were injected with sterile PBS (injection control). Body weight and feed intake were recorded at 0, 24, 48, and 72 hr postendotoxin injection on the individually housed mice. Blood samples were obtained retro-orbitally at 0, 1, and 2 hr after LPS injection (32). Plasma samples were taken from the blood samples after centrifugation and were subsequently analyzed for TNF- α as described below.

Isolation and Culture of Mouse Resident Peritoneal Macrophages. In the *ex vivo* study, 30 mice were either fed the CLA (15 mice) or control diet (15 mice) for 6 weeks. Two or four mice were then sacrificed daily to obtain resident peritoneal macrophages and splenocytes. Peritoneal resident macrophages were collected post euthanasia by injecting 10 ml of ice-cold DMEM media containing 10% FBS and 1% of antibiotics solution (Sigma) into the peritoneal cavities, and then recovering the fluid (33). The fluid was collected and gently laid on top of 3 ml of Histopaque 1081 (Sigma) and was centrifuged at 400g for 15 min at ambient temperature. Mononuclear cells were transferred to a clean centrifuge tube and were washed twice in Mg^{2+} Ca^{2+} -free PBS. Viable cell numbers, as determined by trypan blue exclusion, were greater than 95%. The cells were suspended and adjusted to 1×10^6 viable cells/ml in RPMI 1640 medium with 10% FBS and 1% antibiotics solution (Sigma). One hundred microliters of cell suspension was added to individual wells in a 96-well plate (100,000 cells/well) for 2 hr. Nonadherent cells were removed by washing the monolayer twice with fresh medium. The adherent peritoneal exudate cells are hereafter referred to as macrophages. Macrophages were immediately primed with 5 unit/ml IFN- γ for 4 hr before being stimulated for TNF- α or nitrite production (33). Macrophages were cultured with LPS (500 ng/ml) for 16 hr and the supernatants were collected for TNF- α assay. For nitric oxide assay, phenol-red free DMEM was used, and 500 ng/ml LPS and 5 unit/ml IFN- γ were added to the macrophages and incubated for 44 hr before the culture supernatants were removed for nitrite assay (33).

Splenocyte Isolation. Spleens were removed from the CLA- or control-fed mice and were placed in a petri dish with 10 ml of RPMI 1640 media. A 10-ml sterile syringe plunger was used to disperse the spleen into a single-celled suspension. The media containing the suspended splenocytes were then collected and layered atop 3 ml of Histopaque 1081 (Sigma) in a 15-ml centrifuge tube and were centrifuged at 400g for 15 min at ambient temperature. Mononuclear cells at the interface were collected using a transfer pipette, placed in another centrifugation tube, and 10 ml of media was added. Cells were centrifuged and the pellet was then washed twice with fresh media by centrifugation. Splenocytes were then resuspended to a cell density of 2×10^6 /ml, and 500 μ l was applied into each well in a 24-well plate. Concanavalin A (Con A) at 4 μ g/ml was added into splenocyte culture, and supernatant was col-

lected after 48 hr and frozen for subsequent cytokine analysis.

Cytokine Assay. TNF- α , IL-2, and IL-4 OptEIA enzyme-linked immunoabsorbant assay (ELISA) kits (PharMingen, San Diego, CA) were used to determine the serum TNF- α level as well as culture supernatant IL-2, IL-4, and TNF- α concentration. A standard procedure of cytokine ELISA was performed according to the manufacturer's instructions. Briefly, a plastic plate was coated overnight with a capture antibody for a specific cytokine, followed by washing and blocking the plate. Diluted samples and standards (provided in the kit) were then added and incubated. An extensive wash was applied before secondary antibody and enzyme conjugates were added. Plates then went through another round of incubation and extensive washing. Developing reagent was then added to the plate for 15 min. Color development was stopped during linear increases in substrate utilization by adding 0.5 M sulfuric acid to disrupt enzymatic activity. OD readings of samples were converted to concentration based on the reference curve. At least duplicate samples were analyzed for each cytokine ELISA. The TNF- α kit used in the present study only detects free-form TNF- α but not receptor-bound form. The detection limit for TNF- α , IL-2, and IL-4 is 20, 10, and 10 pg/ml, respectively.

Quantification of Nitrite in Culture Supernatant. Measurement of nitrite has been commonly used as an indirect measurement of nitric oxide formation because nitrite is a stable end product of the highly reactive nitric oxide (34). In the *ex vivo* experiment, 50 μ l of cultured resident macrophages supernatants was taken from each well and mixed with 50 μ l of Griess reagent [1 part of 1% sulfanilamide and 1 part of 0.1% N-(1-naphthyl) ethylenediaminedihydrochloride, mixed immediately before use] in a different 96-well plate. The plate was gently shaken at ambient temperature for 5 min, and OD was measured at 562 nm. A standard curve was established using known $NaNO_2$ preparations. Sample nitrite concentration was extrapolated from the OD based on the standard curve. Duplicate samples were used in every nitrite microplate assay. The typical standard curve had a correlation coefficient of more than 99% between 0 and 20 μ M. Samples were diluted to this range and were analyzed again if they were found to be higher than the upper limit of standards at first analysis.

Colorimetric Tetrazolium Assay. The linear relationship between number of macrophages or splenocytes and cleavage of tetrazolium was measured by a modified 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT; Sigma) assay (35). MTT was dissolved in PBS (5 mg/ml) and filter sterilized through a 0.22- μ m syringe filter. Twenty-five microliters of MTT solution was added to each culture well and these plates were incubated at 37°C for 3 hr. One hundred microliters stop reagent was added to each well and mixed. Stop reagent was prepared by dissolving sodium dodecyl sulfate 20% (w/v) at ambient temperature in 50% N,N-dimethylformamide in demineralized wa-

ter, and pH was adjusted to 4.7 using 80% acetic acid and 2.5% 1 N HCl. The plates were incubated overnight at 37°C to ensure the complete solubilization of cells and the blue crystals of formazan. OD was read on a microplate reader at wavelength of 562 nm. MTT values were determined after Con A stimulation and were used to correct IL-2 and IL-4 data based on cell numbers.

Statistical Analysis. Both nonrepeated and repeated data were analyzed to determine CLA treatment effect by PROC MIX in SAS computer program, version 8 (SAS Institute, Cary, NC) (36). For Figures 1 and 2, data were analyzed by one-way analysis of variance (ANOVA) with two treatments, with repeated measures on experiment unit (mouse). A type "ARH(1)" error structure was used to account for auto-correlated errors. For Figures 3 and 4, data were analyzed by one-way ANOVA with diet as treatment and mice(diet) as error. In addition, the model included a blocking factor to account for variations due to different days of animal sacrifice. For Figure 5, data were analyzed by randomized complete block design with experiment unit




Figure 1. The influence of dietary CLA on LPS-induced body weight loss and feed intake. After feeding CLA or control diet for 6 weeks, mice were injected with either LPS (1 mg/kg) or PBS (data not shown). Cumulative body weight change and feed intake were monitored for 3 days. Each point represents least square mean with pooled error. There were three mice in each group. (a) Repeated data analysis showed dietary CLA was protecting mice against weight loss compared with mice fed the control (corn oil) diet ($P < 0.01$). (b) CLA-fed mice ate more diet after LPS injection compared with control-fed mice ($P < 0.01$).




Figure 2. Influence of dietary CLA on LPS-induced release of plasma TNF- α . Mice were fed a CLA or control diet for 6 weeks before LPS (1 ng/g body weight) or PBS injection. Blood samples were obtained from mice 0, 1, and 2 hr after LPS injection. ELISA was used to determine plasma TNF- α level. Each bar represents least square mean + pooled error (SEM). Plasma TNF- α was unchanged over time in the PBS-injected mice (data not shown). An asterisk indicates significantly different from control at $P < 0.01$.

was mean of quadruplicate wells. Data from five independent experiments were used in the analysis where experiment was treated as block.

Results

Mice fed CLA had significantly less endotoxin-induced body weight loss (0.9 g) over 3 days compared with control-fed endotoxin-injected mice (4.1 g; Fig. 1A). Feed intake after LPS injection was also significantly affected by CLA. After LPS injection, CLA-fed mice ate more feed (2.6 g/day) compared with control-fed mice (1.1 g/day) over a 3-day period (Fig. 1B). In PBS injection treatment, feed intake was 2.8 g/d for the CLA group and 2.6 g/d for the control group. Weight loss in PBS injection treatment was 1.7 g in the control group and 0.1 g in the CLA group (data not shown). In the *in vivo* trial, CLA had a potent inhibitory effect on LPS-induced anorexia and body weight loss.

The evidence that CLA protects against LPS-induced cachexia and the role of dietary CLA on LPS-induced plasma TNF- α was investigated. After mice were fed CLA for 6 weeks, the plasma TNF- α levels were significantly lower in the CLA-fed mice compared with the control-fed mice 2 hr after LPS injection (Fig. 2). The TNF- α production from PBS injection control was unchanged (data not shown).

In the *ex vivo* trial, we studied peritoneal resident macrophage activity after mice were fed the dietary treatments for 6 weeks. We found that even though CLA decreased plasma TNF- α level *in vivo*, CLA did not statistically decreased LPS-induced release of TNF- α in peritoneal resident macrophage in this *ex vivo* system (Fig. 3A). Nitric oxide production, however, was reduced by CLA. Peritoneal resident macrophages from CLA-fed mice had less nitric oxide production when stimulated by IFN- γ and LPS when compared with cells from control-fed mice (Fig. 3B).




Figure 3. Nitric oxide and TNF- α production by resident peritoneal macrophages were affected by CLA feeding. Mice were fed CLA or control diet for 6 weeks before sacrifice. Resident peritoneal macrophages were obtained and primed with IFN- γ . (a) Macrophages were later stimulated with LPS for TNF- α production. (b) Macrophages were later stimulated with LPS and IFN- γ for nitric oxide production. Macrophages from mice fed a CLA diet produced a significantly lower amount of nitrite (53.3 μ M) compared with macrophages from mice fed a control diet (64.2 μ M). TNF- α production, however, was not different between CLA and control groups. Each bar represents least square mean with pooled error. There were 15 mice in each diet treatment. An asterisk indicates significantly different from control at $P < 0.05$.

Spleens were also obtained for lymphocyte cytokine analysis. Splenocytes were treated with Con A for 48 hr before collecting supernatant for cytokine analysis. Splenocytes from CLA-fed mice did show a higher IL-2 production after LPS stimulation and lower IL-4 production than splenocytes from control-fed mice (Fig. 4). The ratio of IL-2 to IL-4 was significantly ($P < 0.05$) higher (ratio = 3.3) for splenocytes from the CLA-fed mice compared with the control-fed mice (ratio = 2.2).

In the *in vitro* experiment, after RAW macrophages were cultured with fatty acid-albumin complexes, they were stimulated with LPS to produce TNF- α . Our data showed that CLA mixture had an inhibitory effect on TNF- α production compared with the LA control. Among isomers, the c9,t11 isomer was responsible for this inhibitory effect (Fig. 5A), but not the t10,c12 isomer (Fig. 5B). In average, 100 μ M c9,t11 CLA isomer decreased macrophage TNF- α production by 60% compared with LA control. This c9,t11 isomer effect was dose responsive as 50 μ M c9,t11 CLA




Figure 4. Cytokine production by splenocytes from mice fed CLA or control diet. After mice were fed CLA or control diet for 6 weeks, splenocytes were harvested and stimulated with concanavalin A for cytokine production. Supernatants were then harvested from the cultured splenocytes and analyzed by ELISA. Cytokine production from each well was corrected by numbers of cells presented at the end of the culture that was determined by colorimetric tetrazolium assay as described in "Materials and Methods." Each bar represents least square mean with pooled error. There are 15 mice in each diet treatment. The IL-2-to-IL-4 ratio is 3.3 for the CLA group and 2.2 for the control group. An asterisk indicates significantly different from control at $P < 0.05$.

alone was 40% inhibitory to macrophage TNF- α production. On the other hand, t10,c12 isomer did not show any effect on macrophage TNF- α production.

Discussion

Dietary fatty acids have been shown to be potent immune regulators. In general, high dietary levels of PUFA or n-3 fatty acids have an inhibitory effect on T cell proliferation and natural killer (NK) cell activity (37). Fish oil, rich in n-3 fatty acids, decreased murine and human LPS-stimulated TNF- α production (38, 39). It has been shown that dietary fish oil prevents endotoxin-induced death in guinea pigs (40), possibly by inhibition of cytokines produced by macrophages (41).

Feeding 0.5% CLA to animals resulted in up to 10 mg of CLA per gram of body fat in several different tissues within 30 days (42), and it took between 2 and 4 weeks to reduce CLA to baseline levels after its withdrawal from the diet (43). Several physiological changes, including body composition change (6), anticarcinogenesis (44), and anti-atherosclerosis (4), were also shown when using up to 0.5% CLA in animal studies.

CLA has been shown to be a potent immune regulator. CLA decreased the LPS-induced cachectic response in chicken, rats, and mice, but at the same time, increased lymphocyte blastogenesis (12, 18). Other groups also reported CLA-increased lymphocyte blastogenesis as well as increased IL-2 production (9, 45, 46). To the best of our knowledge, this is the first report of an elevated Th1/Th2 cytokine ratio in CLA-fed mice. The increased IL-2 production (shown here, and by others) and the decreased IL-4 production associated with dietary CLA may explain the




Figure 5. CLA inhibited TNF- α production from cultured RAW macrophage cell line (a and b). RAW cells were plated (100,000 cells/well) in 24-well plate for 24 hr. Then they were cultured with fatty acid-albumin complex for 24 hr before LPS was added into culture to stimulate TNF- α secretion for 16 hr. Each bar represents least square mean with pooled error (SEM). There were 4 wells for each treatment in a 24-well plate. Bar with a different letter indicates significantly difference at $P < 0.05$. (a) Compared among LA, CLA, and c9, t11 CLA isomer. (b) Compared among LA, CLA, and t10, c12 CLA isomer. Mixed isomers, approximately 42% c9,t11 and 44% t10,c12. Figure shown here represented means of five independent experiments.

reduced IgE production in CLA fed rats (11) as compared with rats fed diets largely based on LA.

Because CLA seems to enhance immunological responses, which may pose a potential risk on immune hypersensitivity, our laboratory has been studying how dietary CLA may affect type I and type III immune hypersensitivity. In a type I hypersensitivity model, CLA reduced antigen-induced histamine and PGE₂ release from tracheae of CLA-fed guinea pigs (30). In an autoimmune (type III hypersensitivity) model, CLA feeding not only protected against end-stage body wasting, but also increased survival days 1.5-fold after the onset of proteinuria in mice (10). Shifted cytokine profiles toward Th1 may interfere with Ig class switch that is needed in producing IgE in immune type I hypersensitivity responses (47). The reduced IgE production by CLA-fed rats (11) would certainly result in a reduced type I hypersensitivity reaction upon challenge, as demonstrated in a guinea pig hypersensitivity model (30). The development of experimentally inducible lupus in mice seems to involve two stages: increased Th1 cytokines followed by elevated Th2 cytokines later in life (48). Increased Th2 cytokine production correlates well to disease progres-

sion. Hence, a CLA-induced shift toward Th1 cytokines could explain, at least in part, increased days of survival post-proteinuria in lupus-prone mice. Inhibited production of TNF- α may also help to explain the reduction in body weight loss in CLA-fed lupus mice at the end stage of systemic lupus disease (10).

Macrophages are sensitive to dietary fatty acid supplementation, as its fatty acid profile reflects such dietary intervention (49, 50). These works also showed that dietary fatty acids also dictate macrophage physiology. Dietary fish oil has been shown to decrease macrophage antigen presentation (51, 52) and to decrease cytokine production as well as mRNA expression (38, 53). In the present study, macrophages were tested as one of the target cells on which CLA modulates immune function. Our data showed that CLA, more specifically c9,t11 CLA, decreased TNF- α production in RAW macrophage cell line. LPS-induced anorexia and cachexia responses were reduced in CLA-fed animals. The plasma TNF- α level after LPS injection was also suppressed in CLA-fed mice. It was unanticipated that mice fed CLA did not suppress *ex vivo* TNF- α production in resident peritoneal macrophages. However, both *in vitro* and plasma samples showed that CLA has an inhibitory effect on TNF- α production. Fatty acid turnover rate in cultured peritoneal resident macrophages may have played a role in the lack of TNF- α response observed in this model. In the cell culture study, CLA was kept at a constant concentration during the *in vitro* experiments, and when blood was drawn for TNF- α analysis, mice were still fed the treatment diet. However, when culture supernatants were collected in the *ex vivo* study, peritoneal macrophages had already been cultured *in vitro* for 24 hr without exogenous exposure to CLA. Moreover, the fetal bovine serum used in the system might not have been as effective as autologous serum (15) in demonstrating CLA's effects on TNF- α production by resident peritoneal macrophages.

The two main isomers of the CLA mixture (c9,t11 and t10,c12) used in this study were considered to have different properties on cytokine regulation. In fact, c9,t11 CLA increased feed efficiency (54, 55), whereas the t10,c12 was the isomer shown to affect body composition (56). In the present study, c9,t11 CLA isomer inhibited TNF- α production from *in vitro* macrophage cultures (Fig. 5a). The slight decrease of TNF- α in 100 μ M t10,c12 CLA isomer treatment probably was thought to be due to c9,t11 CLA isomer contamination of the t10,t12 CLA isomer used (Fig. 5b). Moreover, a recent study reported decreased hepatic TNF- α mRNA expression by c9,t11 CLA in mice (57). It remains to be shown which isomers downregulate IL-4 production and increase IL-2 production. Even though the *in vitro* data suggested that c9,t11 CLA is the active isomer in reducing LPS-induced TNF- α production, further *in vivo* studies with pure CLA isomers are required to validate the *in vitro* data on TNF- α production and determine the effects of different CLA isomers on immune function.

TNF- α was shown to be involved in endotoxin-induced

weight loss and cancer cachexia (58). Muscle degradation was enhanced by TNF- α . Blocking TNF- α function by either injection of anti-TNF- α antibody or TNF- α -binding proteins reduced the cachexia reaction (59, 60). The decreased LPS-induced TNF- α production by CLA may provide at least one mechanism by which CLA enhanced the rate of growth and improved feed efficiency in animals (12, 54, 55).

Thromboxane A_2 (TXA $_2$) is a mediator of renal damage (61). In murine lupus, TXA $_2$ inhibitor prolonged survival of NZB/W F1 mice, and renal TXA $_2$ production was elevated in F1 mice (62). In a study comparing TXB $_2$ (stable end product of TXA $_2$) production and COX gene expression of peripheral blood mononuclear cells (PBMC) among patients with active or inactive lupus nephritis and healthy individuals as a control, COX-2 expression and TXB $_2$ production in PBMC was only elevated in patients with active lupus nephritis (63). Immunostaining of kidney biopsies showed no difference in COX-1, but COX-2 staining in patients with active lupus nephritis was increased compared with patients with inactive lupus nephritis or healthy controls. Double staining of kidney biopsies with anti-COX-2 and anti-CD68 antibodies (macrophage marker) demonstrated that upregulated COX-2 enzyme expression was within the macrophages of glomeruli in patients with active lupus nephritis (63).

Infiltrating macrophages in glomeruli, with upregulated COX-2 enzyme expression, may contribute to the elevated TXA $_2$ level in active lupus nephritis. Regulation of macrophage activities and eicosanoid profiles by CLA may play a role in lupus nephritis. Because COX-2 is the inducible form and the COX-1 is constitutively expressed, decreased antigen-induced TXB $_2$ production but not the basal level of TXB $_2$ production in CLA-fed guinea pigs provides an explanation that CLA may preferentially inhibit COX-2 activity (64). Decreased macrophage activities including TNF- α and nitric oxide productions seen in this study may help to explain the effect of CLA in prolonging survival of NZB/W F1 mice.

In conclusion, these data suggest that cytokine regulation by CLA could be responsible for previous reports demonstrating CLA's anticachectic effects, increased lymphocyte blastogenesis, decreased immune type-I hypersensitivity, decreased IgE production, and prolonged life post proteinuria in the NZB/W F1 autoimmune mouse.

The authors thank Dr. Charles Czuprynski for insightful discussion of this manuscript.

1. Pariza MW, Hargraves WA. A beef-derived mutagenesis modulator inhibits initiation of mouse epidermal tumors by 7,12-dimethylbenz[a]anthracene. *Carcinogenesis* 6:591-593, 1985.
2. Ip MM, Masso-Welch PA, Shoemaker SF, Shea-Eaton WK, Ip C. Conjugated linoleic acid inhibits proliferation and induces apoptosis of normal rat mammary epithelial cells in primary culture. *Exp Cell Res* 250:22-34, 1999.
3. Ip C, Singh M, Thompson HJ, Scimeca JA. Conjugated linoleic acid suppresses mammary carcinogenesis and proliferative activity of the mammary gland in the rat. *Cancer Res* 54:1212-1215, 1994.
4. Nicolosi RJ, Rogers EJ, Kritchevsky D, Scimeca JA, Huth PJ. Dietary conjugated linoleic acid reduces plasma lipoproteins and early aortic atherosclerosis in hypercholesterolemic hamsters. *Artery* 22:266-277, 1997.
5. Kritchevsky D. Antimutagenic and some other effects of conjugated linoleic acid. *Br J Nutr* 83:459-465, 2000.
6. Park Y, Albright KJ, Liu W, Storkson JM, Cook ME, Pariza MW. Effect of conjugated linoleic acid on body composition in mice. *Lipids* 32:853-858, 1997.
7. West DB, Delany JP, Camet PM, Blohm F, Truett AA, Scimeca J. Effects of conjugated linoleic acid on body fat and energy metabolism in the mouse. *Am J Physiol* 44:R667-R672, 1998.
8. Stangl GI. Conjugated linoleic acids exhibit a strong fat-to-lean partitioning effect, reduce serum VLDL lipids and redistribute tissue lipids in food-restricted rats. *J Nutr* 130:1140-1146, 2000.
9. Hayek MG, Han SN, Wu D, Watkins BA, Meydani M, Dorsey JL, Smith DE, Meydani SN. Dietary conjugated linoleic acid influences the immune response of young and old C57BL/6NCrHER mice. *J Nutr* 129:32-38, 1999.
10. Yang M, Pariza MW, Cook ME. Dietary conjugated linoleic acid protects against end stage disease of systemic lupus erythematosus in the NZB/W F1 mouse. *Immunopharmacol Immunotoxicol* 22:433-449, 2000.
11. Sugano M, Tsujita A, Yamasaki M, Noguchi M, Yamada K. Conjugated linoleic acid modulates tissue levels of chemical mediators and immunoglobulins in rats. *Lipids* 33:521-527, 1998.
12. Cook ME, Miller CC, Park Y, Pariza M. Immune modulation by altered nutrient metabolism: nutritional control of immune-induced growth depression. *Poult Sci* 72:1301-1305, 1993.
13. Kelley VE, Ferretti A, Izui S, Strom TB. A fish oil diet rich in eicosapentaenoic acid reduces cyclooxygenase metabolites, and suppresses lupus in MRL-lpr mice. *J Immunol* 134:1914-1919, 1985.
14. Calder PC. Dietary fatty acids and the immune system. *Nutr Rev* 56:S70-S83, 1998.
15. Yaqoob P, Newsholme EA, Calder PC. The effect of dietary lipid manipulation on rat lymphocyte subsets and proliferation. *Immunology* 82:603-610, 1994.
16. Korver DR, Klasing KC. Dietary fish oil alters specific and inflammatory immune responses in chicks. *J Nutr* 127:2039-2046, 1997.
17. Flores EA, Bistrian BR, Pomposelli JJ, Dinarello CA, Blackburn GL, Istfan NW. Infusion of tumor necrosis factor/cachectin promotes muscle catabolism in the rat. A synergistic effect with interleukin 1. *J Clin Invest* 83:1614-1622, 1989.
18. Miller CC, Park Y, Pariza MW, Cook ME. Feeding conjugated linoleic acid to animals partially overcomes catabolic responses due to endotoxin injection. *Biochem Biophys Res Commun* 198:1107-1112, 1994.
19. Spellberg B, Edwards JE Jr. Type 1/Type 2 immunity in infectious diseases. *Clin Infect Dis* 32:76-102, 2001.
20. Hsieh CS, Macatonia SE, Tripp CS, Wolf SF, O'Garra A, Murphy KM. Development of TH1 CD4 $^{+}$ T cells through IL-12 produced by Listeria-induced macrophages. *Science* 260:547-549, 1993.
21. Seder RA, Gazzinelli R, Sher A, Paul WE. Interleukin 12 acts directly on CD4 $^{+}$ T cells to enhance priming for interferon- γ production and diminishes interleukin 4 inhibition of such priming. *Proc Natl Acad Sci U S A* 90:10188-10192, 1993.
22. Kopf M, Le Gros G, Bachmann M, Lamers MC, Bluethmann H, Kohler G. Disruption of the murine IL-4 gene blocks Th2 cytokine responses. *Nature* 362:245-248, 1993.
23. Betz M, Fox BS. Regulation and development of cytochrome c-specific IL-4-producing T cells. *J Immunol* 145:1046-1052, 1990.
24. Demeure CE, Wu CY, Shu U, Schneider PV, Reusser C, Yssel H, Delespasse G. In vitro maturation of human neonatal CD4 T lympho-

- cytes. IL Cytokines present at priming modulate the development of lymphokine production. *J Immunol* 152:4775-4782, 1994.
25. Gajewski TF, Fitch FW. Anti-proliferative effect of IFN- γ in immune regulation. I. IFN- γ inhibits the proliferation of Th2 but not Th1 murine helper T lymphocyte clones. *J Immunol* 140:4245-4252, 1988.
26. D'Andrea A, Aste-Amezaga M, Valiante NM, Ma X, Kubin M, Trinchieri G. Interleukin 10 (IL-10) inhibits human lymphocyte interferon- γ production by suppressing natural killer cell stimulatory factor/IL-12 synthesis in accessory cells. *J Exp Med* 178:1041-1048, 1993.
27. Ohmori Y, Hamilton TA. IL-4-induced STAT6 suppresses IFN- γ -stimulated STAT1-dependent transcription in mouse macrophages. *J Immunol* 159:5474-5482, 1997.
28. Kuhn R, Rajewsky K, Muller W. Generation and analysis of interleukin-4-deficient mice. *Science* 254:707-710, 1991.
29. Punnonen J, Aversa G, Cocks BG, McKenzie AN, Menon S, Zurawski G, de Waal Malefyt R, de Vries JE. Interleukin 13 induces interleukin 4-independent IgG4 and IgE synthesis and CD23 expression by human B cells. *Proc Natl Acad Sci U S A* 90:3730-3734, 1993.
30. Whigham LD, Cook EB, Stahl JL, Saban R, Bjorling DE, Pariza MW, Cook ME. CLA reduces antigen-induced histamine and PGE(2) release from sensitized guinea pig tracheae. *Am J Physiol* 280:R908-R912, 2001.
31. Hirai K, Yamaguchi M, Misaki Y, Takaishi T, Ohta K, Morita Y, Ito K, Miyamoto T. Enhancement of human basophil histamine release by interleukin 5. *J Exp Med* 172:1525-1528, 1990.
32. Roselaar SE, Daugherty A. Lipopolysaccharide decreases scavenger receptor mRNA in vivo. *J Interferon Cytokine Res* 17:573-579, 1997.
33. Coligan JE. *Current Protocols in Immunology*. New York: Greene Publishing Associates and Wiley-Interscience, 1991.
34. Moncada S, Palmer RM, Higgs EA. Nitric oxide: physiology, pathophysiology, and pharmacology. *Pharmacol Rev* 43:109-142, 1991.
35. Hansen MB, Nielsen SE, Berg K. Re-examination and further development of a precise and rapid dye method for measuring cell growth/cell kill. *J Immunol Methods* 119:203-210, 1989.
36. Littell RC, Milliken GA, Stroup WW, Wolfinger RD. *SAS System for Mixed Models*. Cary, NC: SAS Institute, 1996.
37. Kinsella JE, Lokesh B, Broughton S, Whelan J. Dietary polyunsaturated fatty acids and eicosanoids: potential effects on the modulation of inflammatory and immune cells: an overview. *Nutrition* 6:24-44; discussion 59-62, 1990.
38. Renier G, Skamene E, DeSanctis J, Radzioch D. Dietary n-3 polyunsaturated fatty acids prevent the development of atherosclerotic lesions in mice. Modulation of macrophage secretory activities. *Arterioscler Thromb* 13:1515-1524, 1993.
39. Meydani SN, Endres S, Woods MM, Goldin BR, Soo C, Morrill-Labrode A, Dinarello CA, Gorbash SH. Oral (n-3) fatty acid supplementation suppresses cytokine production and lymphocyte proliferation: comparison between young and older women. *J Nutr* 121:547-555, 1991.
40. Mascioli EA, Iwasa Y, Trimbo S, Leader L, Bistran BR, Blackburn GL. Endotoxin challenge after menhaden oil diet: effects on survival of guinea pigs. *Am J Clin Nutr* 49:277-282, 1989.
41. Boutard V, Fouqueray B, Philippe C, Perez J, Baud L. Fish oil supplementation and essential fatty acid deficiency reduce nitric oxide synthesis by rat macrophages. *Kidney Int* 46:1280-1286, 1994.
42. Park Y. Regulation of Energy Metabolism and the Catabolic Effects of Immune Stimulation by Conjugated Linoleic Acid. PhD dissertation. Madison (WI): University of Wisconsin, 1996.
43. Park Y, Albright KJ, Storkson JM, Liu W, Cook ME, Pariza MW. Changes in body composition in mice during feeding and withdrawal of conjugated linoleic acid. *Lipids* 34:243-248, 1999.
44. Ip C, Chin SF, Scimeca JA, Pariza MW. Mammary cancer prevention by conjugated dienoic derivative of linoleic acid. *Cancer Res* 51:6118-6124, 1991.
45. Chew BP, Wong TS, Shultz TD, Magnuson NS. Effects of conjugated dienoic derivatives of linoleic acid and β -carotene in modulating lymphocyte and macrophage function. *Anticancer Res* 17:1099-1106, 1997.
46. Wong MW, Chew BP, Wong TS, Hosick HL, Boylston TD, Shultz TD. Effects of dietary conjugated linoleic acid on lymphocyte function and growth of mammary tumors in mice. *Anticancer Res* 17:987-993, 1997.
47. Snapper CM, Pecanha LM, Levine AD, Mond JJ. IgE class switching is critically dependent upon the nature of the B cell activator, in addition to the presence of IL-4. *J Immunol* 147:1163-1170, 1991.
48. Segal R, Bermas BL, Dayan M, Kalush F, Shearer GM, Mozes E. Kinetics of cytokine production in experimental systemic lupus erythematosus: involvement of T helper cell 1/T helper cell 2-type cytokines in disease. *J Immunol* 158:3009-3016, 1997.
49. Leslie CA, Gonnerman WA, Ullman MD, Hayes KC, Franzblau C, Cathcart ES. Dietary fish oil modulates macrophage fatty acids and decreases arthritis susceptibility in mice. *J Exp Med* 162:1336-1349, 1985.
50. Calder PC, Bond JA, Harvey DJ, Gordon S, Newsholme EA. Uptake and incorporation of saturated and unsaturated fatty acids into macrophage lipids and their effect upon macrophage adhesion and phagocytosis. *Biochem J* 269:807-814, 1990.
51. Fujikawa M, Yamashita N, Yamazaki K, Sugiyama E, Suzuki H, Hamazaki T. Eicosapentaenoic acid inhibits antigen-presenting cell function of murine splenocytes. *Immunology* 75:330-335, 1992.
52. Huang SC, Misfeldt ML, Fritsche KL. Dietary fat influences Ia antigen expression and immune cell populations in the murine peritoneum and spleen. *J Nutr* 122:1219-1231, 1992.
53. Calder PC. n-3 polyunsaturated fatty acids and cytokine production in health and disease. *Ann Nutr Metab* 41:203-234, 1997.
54. Pariza MW, Park Y, Cook ME. Mechanisms of action of conjugated linoleic acid: evidence and speculation. *Exp Biol Med* 223:8-13, 2000.
55. Chin SF, Storkson JM, Albright KJ, Cook ME, Pariza MW. Conjugated linoleic acid is a growth factor for rats as shown by enhanced weight gain and improved feed efficiency. *J Nutr* 124:2344-2349, 1994.
56. Park Y, Storkson JM, Albright KJ, Liu W, Pariza MW. Evidence that the trans-10,cis-12 isomer of conjugated linoleic acid induces body composition changes in mice. *Lipids* 34:235-241, 1999.
57. Roche HM, Noone E, Nugent A, Gibney MJ. Nutrient regulation of SREBP1c gene expression: CLA reduces plasma TAG concentrations by down-regulating hepatic SREBP1c in an isomer-specific effect of CLA on lipoprotein metabolism in man and gene expression in Ob/Ob mice. Ninety-second AOCS Annual Meeting and Expo: 2001 May 13-16; Minneapolis, Minnesota. Champaign (IL): AOCS Press, pp858-859.
58. Tisdale MJ. Biology of cachexia. *J Natl Cancer Inst* 89:1763-1773, 1997.
59. Swiergiel AH, Dunn AJ. The roles of IL-1, IL-6, and TNF α in the feeding responses to endotoxin and influenza virus infection in mice. *Brain Behav Immun* 13:252-265, 1999.
60. Costelli P, Carbo N, Tessitore L, Bagby GJ, Lopez-Soriano FJ, Argiles JM, Baccino FM. Tumor necrosis factor- α mediates changes in tissue protein turnover in a rat cancer cachexia model. *J Clin Invest* 92:2783-2789, 1993.
61. Remuzzi G, FitzGerald GA, Patrono C. Thromboxane synthesis and action within the kidney. *Kidney Int* 41:1483-1493, 1992.
62. Salvati P, Lamberti E, Ferrario R, Ferrario RG, Scampini G, Pugliese F, Barsotti P, Patrono C. Long-term thromboxane-synthase inhibition prolongs survival in murine lupus nephritis. *Kidney Int* 47:1168-1175, 1995.
63. Tomasoni S, Noris M, Zappella S, Gotti E, Casiraghi F, Bonazzola S, Benigni A, Remuzzi G. Upregulation of renal and systemic cyclooxygenase-2 in patients with active lupus nephritis. *J Am Soc Nephrol* 9:1202-1212, 1998.
64. Whigham LD. Inhibition of Lipid Mediators in Type I (Immediate) Hypersensitivity Immune Reaction by Conjugated Linoleic Acid (CLA). PhD dissertation. Madison (WI): University of Wisconsin, 2000.

Rapid report

NF- κ B independent inhibition of lipopolysaccharide-induced cyclooxygenase by a conjugated linoleic acid cognate, conjugated nonadecadienoic acid

Guangming Li^{a,f}, Baiyan Dong^{b,c}, Daniel E. Butz^{d,f}, Yeonhwa Park^e,
Michael W. Pariza^{a,b,c,d}, Mark E. Cook^{a,c,d,f,*}

^a Molecular and Environmental Toxicology, University of Wisconsin-Madison, 1675 Observatory Dr., Madison, WI 53706, USA

^b Department of Food Science, University of Wisconsin-Madison, Dr., Madison, WI 53706, USA

^c Department of Microbiology and Food Toxicology, University of Wisconsin-Madison, Dr., Madison, WI 53706, USA

^d Department of Nutritional Sciences, University of Wisconsin-Madison, Dr., Madison, WI 53706, USA

^e Department of Food Science, University of Massachusetts, 100 Holdsworth Way, Amherst, MA 01003, USA

^f Animal Sciences Department, University of Wisconsin-Madison, 1675 Observatory Dr., Madison, WI 53706, USA

Received 27 June 2006; received in revised form 27 July 2006; accepted 31 July 2006

Available online 5 August 2006

Abstract

10t, 12c-CLA was shown to inhibit COX-2 expression through the NF- κ B pathway. In the current study, conjugated nonadecadienoic acid (CNA) was shown to decrease inducible COX-2 protein and mRNA and PGE₂ release to the similar extent as 10t, 12c-CLA in Raw264.7 macrophage. However, unlike 10t, 12c-CLA, inhibition of COX-2 mRNA/protein by CNA was independent of the NF- κ B pathway. The data indicate the regulation of COX-2 by select conjugated fatty acids and hence their anti-inflammatory actions could operate through different signal transduction pathways.

© 2006 Elsevier B.V. All rights reserved.

Keywords: Conjugated linoleic acid; Conjugated nonadecadienoic acid; Cyclooxygenase; NF- κ B; Prostaglandin

Conjugated linoleic acid (CLA) has been shown to prevent adverse events associated with inflammatory diseases. In animal feeding trials, CLA (mixed isomers of 9c, 11t and 10t, 12c-CLA) was shown to decrease immune-induced wasting [1–3], to decrease inflammatory mediator release from antigen challenged airway [4,5], to increase longevity in a lupus

erythematosus model [6,7], and to decrease mucosal thickening/erosions in a model of colitis [8], and to decrease joint swelling in a monoclonal antibody-induced arthritis (Butz and Cook, unpublished data). Regulation of cyclooxygenase-2 (COX-2) [5,9], tumor necrosis factor alpha (TNF) [3,10], and inducible nitric oxide synthase [10] by CLA suggest that CLA may play a key role in inhibiting the nuclear factor kappa B (NF- κ B) activation. Recent evidence in immune related cells and tissues suggested that CLA-induced down regulation of COX-2 protein and mRNA level may be related to an upstream regulation of signal transduction through the NF- κ B pathway [9]. The isomer shown to inhibit the p50 and p65 subunits binding to DNA was 10t, 12c-CLA.

Structure/function studies on CLA cognates and derivatives [11] showed that a 19-carbon CLA cognate, conjugated nonadecadienoic acid (98.6% CNA, with 10c, 12t-CNA, 42.3%; 11t, 13c-CNA, 48.0%; 10c, 12c-CNA and 11c, 13c-

Abbreviations: CLA, conjugated linoleic acid; COX-2, cyclooxygenase-2; CAN, conjugated nonadecadienoic acid; DMEM, Dulbecco's Modified Eagle's Medium; ELA, enzyme immunoassay; ERK, extracellular signal-related kinase; FBS, fetal bovine serum; G3PDH, glyceraldehyde-3-phosphate dehydrogenase; I κ B α , Inhibitory κ B protein α ; IKK, inhibitor- κ B kinase; JNK, C-jun NH2-terminal kinase; LA, linoleic acid; LPS, lipopolysaccharide; MAPK, mitogen-activated protein kinase; NF- κ B, Nuclear factor κ B; PGE₂, prostaglandin E₂; PVDF, polyvinylidene difluoride; RIPA, radioimmunoprecipitation assay; TNF, tumor necrosis factor α .

* Corresponding author. Animal Sciences Department, University of Wisconsin-Madison, 1675 Observatory Dr., Madison, WI 53706, USA. Tel.: +1 608 262 7747; fax: +1 608 262 5157.

CNA 3.2%; 10t, 12t-CNA and 11t, 13t-CNA 5.1%; and unidentified 1.4%, see reference 12 for the synthesis of CNA), was at least as effective as CLA in regulating fat metabolism in 3T3-L1 adipocytes or growing mice [12]. Both the 10t, 12c-CLA and CNA were shown to similarly reduce leptin secretion [13], and the mRNA and protein level of select adipocyte genes in 3T3-L1 adipocytes [14]. In our continued studies on the mechanisms by which CLA regulates COX-2, we investigated whether the new CLA cognate, CNA, also inhibits COX-2 by a similar mechanism to 10t, 12c-CLA.

We first examined whether CNA could inhibit LPS-induced COX-2 protein expression. Raw264.7 macrophage was incubated with fatty acids for 24 h and then stimulated with LPS (100 ng/ml) for 8 h in the presence of LA, 10t, 12c-CLA or CNA (0, 3.3, 10, 33 and 100 μ M). COX-2 was not induced without LPS stimulation. COX-2 protein level was significantly inhibited by CNA at 33 μ M and 100 μ M (77.8% and 58.7% of LPS-stimulated ethanol vehicle treated cells respectively, Fig. 1A, ethanol <0.2%). Similarly, COX-2 protein level was also reduced by 10t, 12c-CLA at 33 μ M and 100 μ M to 46.7% and 26.4% of LPS-stimulated cells (ethanol vehicle-treated only, Fig. 1A). 10t, 12c-CLA was more potent in reducing COX-2 protein expression compared with CNA (Fig. 1A). Since CNA only contains 48% 11t, 13c isomer (the hypothesized active isomer) in the mixture (the corresponding isomer of 10t, 12c-CLA in CNA), 10t, 12c-CLA and CNA appear equally effective in inhibiting COX-2 protein level.

PGE₂ release from macrophage was also measured to support 10t, 12c-CLA and CNA's inhibitory effects on COX-2 protein level. Basal release of PGE₂ from macrophage was ~1.1 ng/mg protein (without LPS exposure). CNA and 10t, 12c-CLA were both shown to inhibit PGE₂ release in a concentration-dependent manner. At 100 μ M, PGE₂ level in CNA and 10t, 12c-CLA-treated sample was reduced from 20.0 ng/mg protein (LPS stimulated/ethanol vehicle -treated only) to 5.43 ng/mg and 5.15 ng/mg (Fig. 1B). That both CNA and 10t, 12c-CLA were shown to inhibit PGE₂ release at their lowest concentrations suggests these isomers not only reduce protein expression but may also inhibit COX enzymatic activity, as previously shown with 10t, 12c-CLA [9]. LA, as a precursor of AA, was shown to increase PGE₂ release into the culture media in a concentration-dependent manner [9]. PGE₂ release in LA-treated macrophages at 100 μ M was 31.9 ng/mg protein (Fig. 1B).

We further examined whether reduced COX-2 protein level by conjugated fatty acids could be due to the reduced mRNA level. Raw264.7 macrophage was incubated with conjugated fatty acids for 24 h and then was stimulated with LPS for 4 h in the presence of LA, 10t, 12c-CLA or CNA. It was shown that CNA reduced mRNA level significantly at 10 μ M, 33 μ M, and 100 μ M (by 48.9% at 100 μ M) while 10t, 12c-CLA reduced COX-2 mRNA at 100 μ M by 23.6% (Fig. 2). LA was shown to have no effect on COX-2 mRNA level (Fig. 2). We have previously shown differential regulation of 10t, 12c-CLA on COX-2 mRNA and protein level [9].

One pathway leading to the transcription of COX-2 after LPS stimulation involves the activation of the NF- κ B via toll like




Fig. 1. Effect of CNA on LPS-induced COX-2 protein expression (A) and PGE₂ release in Raw264.7 macrophage. 1×10^6 cells were seeded in each well of a 24-well plate and were grown to confluence. LA, 10t, 12c-CLA and CNA (0, 3.3, 10, 33, 100 μ M) were incubated with cells for 24 h before and during LPS (100 ng/ml) stimulation for 8 h (see reference 9). Cell lysate and culture medium were harvested for western blot analysis of COX-2 (A) and PGE₂ determination by enzyme immunoassay (EIA, B). For western blot, the total protein level in the cell lysate was quantitated with Bradford assay (BioRad Laboratories, Hercules, CA) [22] and 25 μ g protein per lane was loaded onto a 10% SDS-polyacrylamide gel and resolved by electrophoresis. Proteins on the gel were transferred to polyvinylidene difluoride (PVDF) membrane. After non-specific blocking in 5% non-fat milk solution, the membrane was incubated with primary COX-2, or G3PDH antibody, then a secondary horseradish peroxidase-coupled antibody before being exposed to Fuji medical X-ray film. In panel A, ratio of COX-2/G3PDH was obtained by normalizing the band area of COX-2 to corresponding band area of G3PDH. CNA was synthesized by Dr. Ken Ku from the Food and Drug Administration, USA. The composition of CNA was 10c, 12c-CNA, 42.3%; 11t, 13c-CNA, 48.0%; 10c, 12c-CNA and 11c, 13c-CNA 3.2%; 10t, 12t-CNA and 11t, 13t-CNA 5.1%; and unidentified 1.4%. 10t, 12c-CLA (>98%) and LA (>99%) were purchased from Matreya, Inc. (Pleasant Gap, PA). COX-2 antibody and PGE₂ EIA kits were from Cayman Chemical (Ann Arbor, MI) and G3PDH antibody was from Trevigen (Gaithersburg, MD). These experimental results were repeatable in at least two independent experiments. Data are presented as means \pm SD from four replicates of one representative experiment. SAS procedure GLM was used to do the multiple regressions. *P* value < 0.05 was considered statistically significant.




Fig. 2. Effect of CNA on LPS-induced COX-2 mRNA level in Raw264.7 macrophage. 1×10^6 cells were seeded in each well of a 24-well plate and were grown to confluence. LA, 10t, 12c-CLA and CNA at different concentrations (0, 3.3, 10, 33, 100 μ M) were incubated with cells for 24 h before and during LPS (100 ng/ml) stimulation for 4 h. mRNA isolation, reverse transcription and polymerase chain reaction were conducted as previously described (see reference 9). Briefly, Raw264.7 macrophage was lysed in 100 μ l lysis buffer solution (4 M guanidium isothiocyanate, 0.5% sarcosyl, 10 mM Tris-HCl, pH 8.0 and 1% dithiothreitol) and DNA was sheared by passing cell lysate through a 25 gauge needle. Binding buffer 200 μ l (10 mM Tris-HCl, pH 8.0, 1 mM EDTA and 400 mM NaCl) was added for mRNA to combine with magnetic particles. Supernatants from the above solution were incubated with magnetized oligo(dT) magnetic particles for at least 5 min. The particles were then washed for 4 times with wash buffer (150 mM NaCl, 10 mM Tris-HCl, pH 8.0 and 1 mM EDTA) and the associated mRNA were rinsed off the beads with 10 μ l elution buffer (2 mM EDTA). Isolated mRNA (2 μ l) was reverse-transcribed in the master mix (reaction buffer 7.0 μ l, random primer 100 μ M 1 μ l, dNTP 10 mM 0.4 μ l, reverse transcriptase 0.2 μ l and H_2O 12.4 μ l). Polymerase chain reaction was used to amplify the individual gene using respective primers. Glyceraldehyde-3-phosphate dehydrogenase (G3PDH) gene was amplified as internal control. The respective primers are: COX-2 upstream: 5'-CAA-GCA-GTG-GCA-AGG-CCT-CCA-3' and downstream 5'-GGC-ACT-TGC-ATT-GAT-GGT-GGC-3'. G3PDH upstream: 5'-GGC-ATT-CTC-GGC-TAC-ACT-GA-3 and downstream: 5'-CAT-ACC-AGG-AAA-ATA-GCT-TGA-C-3'. Ratio of COX-2/G3PDH was obtained by normalizing COX-2 mRNA with corresponding G3PDH mRNA level. These experimental results were repeatable in at least two separate experiments. Data were presented as means \pm SD from 4 replicates of one experiment. SAS procedure GLM was used to do the multiple regressions. P value < 0.05 was considered statistically significant.

receptor-4. NF- κ B is a transcription factor which consists of heterodimer or homodimer of p50, p65, p105, p52, c-Rel or RelB. Upon stimulation by LPS, or TNF, the inhibitory proteins (I κ B α , I κ B β , or I κ B ϵ) associated with the NF- κ B become phosphorylated and degraded in a ubiquitin-proteasome-dependent pathway [15,16]. To examine whether reduced mRNA by CNA was due to the inhibition of the NF- κ B pathway, Raw264.7 macrophages were treated with fatty acids for 24 h and then LPS was introduced for 4 h in the presence of fatty acids. LPS increased I κ B α phosphorylation as shown previously [9]. CNA was shown to have no effect on I κ B α phosphorylation while 10t, 12c-CLA inhibited the I κ B α phosphorylation in a concentration-dependent manner (Fig. 3). Likewise, LA had no effect on I κ B α phosphorylation. Consistent with the effect of CNA, LA and 10t, 12c-CLA on I κ B α phosphorylation, the increased binding of two NF- κ B proteins (p50 and p65) to the DNA consensus sites following LPS exposure was not affected by CNA and LA, but was inhibited by 10t, 12c-CLA in a concentration-dependent manner (Table 1).




Fig. 3. Effects of CNA on phosphorylation of I κ B α in Raw 264.7 macrophage. 1×10^6 cells were seeded in each well of a 24-well plate and were grown to confluence. LA, 10t, 12c-CLA (50 and 100 μ M) and CNA (3.3, 10, 33 and 100 μ M) were incubated with cells for 24 h before and during LPS (1000 ng/ml) stimulation for 4 h. Cell lysate was harvested for western blot analysis of phosphorylated I κ B α and β -actin. The band area of phosphorylated I κ B α was normalized to corresponding band area of β -actin. % Positive Control = (normalized phosphorylated I κ B α area in ethanol vehicle or fatty acids treated samples/normalized phosphorylated I κ B α area in ethanol vehicle and LPS treated samples) \times 100. Data are presented as means \pm SD from four replicates of one experiment. Student's t -test was conducted for the statistical comparison between fatty acids, LPS treated samples and ethanol vehicle, LPS treated samples. P value < 0.05 was considered statistically significant.

These data suggest that CNA regulates COX-2 by an NF- κ B independent process and 10t, 12c-CLA may inhibit other signaling pathway(s) in addition to the NF- κ B pathway. It is well known that stimulation of macrophage by LPS also involves activation of mitogen-activated protein kinase (MAPK) in addition to the NF- κ B pathway [17–19]. MAPK family comprises of the extracellular signal-related kinase (ERK) 1/2, p38 MAP kinase and c-jun NH2-terminal kinase (JNK). Inhibition of individual MAPK was shown to inhibit downstream pro-inflammatory cytokine production. SB202190, a selective inhibitor of p38 MAPK, inhibited LPS-induced IL-1 β and TNF in monocytes [19], LPS-induced COX-2 mRNA level and PGE $_2$ release in Raw264.7 macrophage without inhibiting the NF- κ B activity [20]. Another MAPK inhibitor, PD98059

Table 1
Effect of CNA on nuclear factors p50 and p65 binding to the DNA consensus site

Treatments	p50	p65
Vehicle	0.384 \pm 0.094(c)	0.383 \pm 0.06(d)
Vehicle + LPS	0.728 \pm 0.099(a, b)	0.688 \pm 0.11(a)
LA + LPS	0.689 \pm 0.11(a, b)	0.599 \pm 0.05(a, b)
10t, 12c-CLA + LPS	0.556 \pm 0.07(b, c)	0.440 \pm 0.06(b, c, d)
CNA + LPS	0.664 \pm 0.08(a, b)	0.575 \pm 0.09(a, b)
LA + LPS	0.712 \pm 0.08(a, b)	0.615 \pm 0.20(a)
10t, 12c-CLA + LPS	0.548 \pm 0.07(b, c)	0.401 \pm 0.02(c, d)
CNA + LPS	0.857 \pm 0.34(a)	0.553 \pm 0.07(a, b, c)

1×10^6 cells were seeded in each well of a 24-plate and were grown to confluence. LA, 10t, 12c-CLA and CNA at 50 or 100 μ M were incubated with cells for 24 h before and during LPS (1000 ng/ml) stimulation for 4 h. Cell lysate was harvested for the determination of p50 and p65 binding according to kit instructions (Active Motif, Carlsbad, CA). Data are presented as means \pm SD from four replicates of one experiment. Data were analyzed by one-way ANOVA. Within each NF- κ B subunit (p50 or p65) means without a common letter in parentheses are statistically different (P value < 0.05).

was also shown to inhibit COX-2 mRNA and PGE₂ release independent of the NF- κ B [20]. Potentially, inhibitory effects of CNA and 10*t*, 12*c*-CLA on LPS-induced COX-2 mRNA expression could be attributed to the inhibition of MAPK/ERK/JNK pathways, a hypothesis supported by a recent report of CLA's inhibitory effects on MAPK in osteoclast [21].

References

- [1] M.E. Cook, C.C. Miller, Y. Park, M. Pariza, Immune modulation by altered nutrient metabolism—Nutritional control of immune-induced growth depression, *Poult. Sci.* 72 (1993) 1301–1305.
- [2] E. Graves, A. Hitt, M.W. Pariza, M.E. Cook, D.O. McCarthy, Conjugated linoleic acid preserves gastrocnemius muscle mass in mice bearing the colon-26 adenocarcinoma, *Res. Nurs. Health* 28 (2005) 48–55.
- [3] D. Butz, G. Li, M. Cook, 10*t*, 12*c* conjugated linoleic acid induces compensatory growth after immune challenge, *J. Nutr. Biochem.* (in press).
- [4] L.D. Whigham, E.B. Cook, J.L. Stahl, R. Saban, D.E. Bjorling, M.W. Pariza, M.E. Cook, CLA reduces antigen-induced histamine and PGE₂ release from sensitized guinea pig tracheae, *Am. J. Physiol.: Regul., Integr. Comp. Physiol.* 280 (2001) R908–R912.
- [5] L.D. Whigham, A. Higbee, D.E. Bjorling, Y. Park, M.W. Pariza, M.E. Cook, Decreased antigen-induced sicosanoid release in conjugated linoleic acid-fed guinea pigs, *Am. J. Physiol.: Regul., Integr. Comp. Physiol.* 282 (2002) R1104–R1112.
- [6] M. Yang, M.W. Pariza, M.E. Cook, Dietary conjugated linoleic acid protects against end stage disease of systemic lupus erythematosus in the NZB/W F1 mouse, *Immunopharmacol. Immunotoxicol.* 22 (2000) 433–449.
- [7] M. Yang, M.E. Cook, Dietary CLA decreased weight loss and extended survival following the onset of kidney failure in NZB/W F1 mice, *Lipids* 38 (2003) 21–24.
- [8] R. Hontela, M.J. Wannemuehler, D.R. Zimmerman, D.L. Hutto, J.H. Wilson, D.U. Ahn, R.J. Bassaganya, Nutritional regulation of porcine bacterial-induced colitis by conjugated linoleic acid, *J. Nutr.* 132 (2002) 2019–2027.
- [9] G. Li, D. Barnes, D. Butz, D. Bjorling, M.E. Cook, 10*t*, 12*c*-conjugated linoleic acid inhibits lipopolysaccharide-induced cyclooxygenase expression in vitro and in vivo, *J. Lipid Res.* 46 (2005) 2134–2142.
- [10] M. Yang, M.E. Cook, Dietary conjugated linoleic acid decreased cachexia, macrophage tumor necrosis factor- α production, and modifies splenocyte cytokines production, *Exp. Biol. Medicine* (Maywood) 228 (2003) 51–58.
- [11] Y. Park, J.M. Storkson, J.M. Ntambi, M.E. Cook, C.J. Sih, M.W. Pariza, Inhibition of hepatic stearoyl-CoA desaturase activity by trans-10, cis-12 conjugated linoleic acid and its derivatives, *Biochim. Biophys. Acta* 1486 (2000) 285–292.
- [12] Y. Park, M.W. Pariza, The effects of dietary conjugated nonadecadienoic acid on body composition in mice, *Biochim. Biophys. Acta* 1533 (2001) 171–174.
- [13] K.W. Kang, M.W. Pariza, Trans-10, cis-12-conjugated linoleic acid reduces leptin secretion from 3T3-L1 adipocytes, *Biochem. Biophys. Res. Commun.* 287 (2001) 377–382.
- [14] K. Kang, W. Liu, K.J. Albright, Y. Park, M.W. Pariza, Trans-10, cis-12 CLA inhibits differentiation of 3T3-L1 adipocytes and decreases ppar γ expression, *Biochem. Biophys. Res. Commun.* 303 (2003) 795–799.
- [15] S. Ghosh, M. Karin, Missing pieces in the NF- κ B puzzle, *Cell* 109 (2002) S81–S96.
- [16] A.S. Baldwin Jr., Series introduction: the transcription factor NF- κ B and human disease, *J. Clin. Invest.* 107 (2001) 3–6.
- [17] C.J. Tsai, Y. Chao, C.W. Chen, W.W. Lin, Aurintricarboxylic acid protects against cell death caused by lipopolysaccharide in macrophages by decreasing inducible nitric-oxide synthase induction via I κ B kinase, extracellular signal-regulated kinase, and p38 mitogen-activated protein kinase inhibition, *Mol. Pharmacol.* 62 (2002) 90–101.
- [18] C.C. Chen, J.K. Wang, p38 but not p44/42 mitogen-activated protein kinase is required for nitric oxide synthase induction mediated by lipopolysaccharide in raw 264.7 macrophages, *Mol. Pharmacol.* 55 (1999) 481–488.
- [19] C.L. Mantley, S.W. Wang, S.D. Kinney, Z.B. Yao, SB202190, a selective inhibitor of p38 mitogen-activated protein kinase, is a powerful regulator of LPS-induced mRNA in monocytes, *J. Leukocyte Biol.* 64 (1998) 409–417.
- [20] C.J. Lo, Mapk regulation of prostaglandin E₂ production by lipopolysaccharide-stimulated macrophages is not dependent on nuclear factor κ B, *J. Surg. Res.* 113 (2003) 189–194.
- [21] M.M. Rahman, A. Bhattacharya, G. Fernandes, Conjugated linoleic acid inhibits osteoclast differentiation of raw264.7 cells by modulating rankl signaling, *J. Lipid Res.* 46 (2006) 1739–1748.
- [22] M.M. Bradford, Rapid and sensitive method for quantitation of microgram quantities of protein utilizing principle of protein-dye binding, *Anal. Biochem.* 72 (1976) 248–254.

JNB

STEENBOCK
MEMORIAL LIBRARY

UCI 11 2004

U.W. - MADISON

the Journal of

Nutritional Biochemistry

VOLUME 15 NUMBER 9 SEPTEMBER 2004

In This Issue

- Neuroprotective Mechanism of Green Tea Polyphenols
- Alpha-Lactalbumin Concentration in Human Milk
- Human Responses to L-Arginine Supplementation
- CLA and Lipoprotein Lipase Activity in Adipocytes

Editor-in-Chief
Bernhard Hennig

Structure–activity relationship of conjugated linoleic acid and its cognates in inhibiting heparin-releasable lipoprotein lipase and glycerol release from fully differentiated 3T3-L1 adipocytes

Yeonhwa Park^a, Jayne M. Storkson^a, Wei Liu^a, Karen J. Albright^a, Mark E. Cook^{a,b},
Michael W. Pariza^{a,*}

^aFood Research Institute, University of Wisconsin–Madison, 1925 Willow Drive, Madison, WI 53706, USA

^bDepartment of Animal Sciences, University of Wisconsin–Madison, Madison, WI, USA

Received 5 October 2003; received in revised form 24 March 2004; accepted 19 April 2004

Abstract

Conjugated linoleic acid (CLA) reduces body fat in part by inhibiting the activity of heparin-releasable lipoprotein lipase (HR-LPL) activity in adipocytes, an effect that is induced by the *trans*-10,*cis*-12 CLA isomer. In this study we used a series of compounds that are structurally related to CLA (i.e., CLA cognates) to investigate the structural basis for this phenomenon. None of the 18:1 CLA cognates that were tested, nor *trans*-9,*cis*-12 18:2, *cis*-12-octadecen-10-ynoic acid (10y,*cis*-12) or 11-(2'-(*n*-pentyl)phenyl)-10-undecylenic acid (designated P-10), exhibited any significant effect on HR-LPL activity. Among the CLA derivatives (alcohol, amide, and chloride) that were tested, only the alcohol form inhibited HR-LPL activity, although to a lesser extent than CLA itself. In addition, intracellular TG was reduced only by *trans*-10,*cis*-12 CLA and the alcohol form of CLA. Hence it appears that the *trans*-10,*cis*-12 conjugated double bond in conjunction with a carboxyl group at C-1 is required for inhibition of HR-LPL activity, and that an alcohol group can partially substitute for the carboxyl group. We also studied glycerol release from the cells, observing that this was enhanced by *trans*-10 18:1, *trans*-13 18:1, *cis*-12 18:1, *cis*-13 18:1, P-10 but was reduced by *cis*-9 18:1, the alcohol and amide forms of CLA or 10y,*cis*-12. Accordingly the structural feature or features involved in regulating lipolysis appear to be more complex. Despite enhancing lipolysis in cultured 3T3-L1 adipocytes, *trans*-10 18:1 did not reduce body fat gain when fed to mice. © 2004 Elsevier Inc. All rights reserved.

Keywords: Conjugated linoleic acid; CLA; Lipoprotein lipase; 3T3-L1; *cis*-9; *trans*-11 CLA; *trans*-10; *cis*-12 CLA

1. Introduction

In the last decade, conjugated linoleic acid (CLA) has been studied intensively because of its unusual biological activities [1]. One of the most interesting aspects of CLA is its ability to reduce body fat while increasing lean body mass [2]. As part of the mechanism for these activities, it has been suggested that CLA reduces heparin-releasable lipoprotein lipase (HR-LPL) activity in adipocytes, and in fact this has been demonstrated in 3T3-L1 adipocyte cell culture model [2]. CLA also enhances fatty acid β -oxidation in muscle, which indicates increased use of fat as an energy source [2]. Since CLA is a mixture of isomers, it is of interest to determine which isomer is responsible for these biological activities. Previously we showed that

the *trans*-10,*cis*-12 isomer of CLA (one of two major isomers present in synthetically prepared CLA) is responsible for its reduction in body fat and for inhibition of HR-LPL activity in 3T3-L1 adipocytes [3]. This isomer is also responsible for the inhibition of stearoyl-CoA desaturase (SCD) [4], and reducing apolipoprotein B (apo B) secretion in cultured HepG2 cells [5]. Thus it is of interest whether this CLA isomer may also be solely responsible for inhibition of HR-LPL activity. To determine the key structural feature of *trans*-10,*cis*-12 CLA, we tested fatty acids that are structurally related to *trans*-10,*cis*-12 CLA using 3T3-L1 adipocytes. We also tested these compounds on glycerol release and intracellular triacylglyceride (TG) to estimate lipolysis and total fat deposit, respectively. In addition, since previous findings indicate that a *trans*-10 double bond is a key structure for inhibiting apo B secretion in HepG2 cells [5], we tested *trans*-10 18:1 on body compositional changes compared to *trans*-10,*cis*-12 CLA.

* Corresponding author. Tel.: (608) 263-6955; fax: (608) 263-1114.
E-mail address: mwpariza@facstaff.wisc.edu (M.W. Pariza).

2. Methods and materials

2.1. Materials

Triolein, [9,10-³H (N)] triolein (specific activity 12 Ci/mmol) was obtained from American Radiolabeled Chemicals Incorporation (St. Louis, MO) and [1-¹⁴C] linoleic acid (specific activity 55 mCi/mmol) was obtained from Amersham Life Science (Arlington Heights, IL). 3T3-L1 preadipocytes were purchased from American Type Culture Collection (Rockville, MD). *trans*-9,*cis*-12 octadecadienoic acid (31.7%), *cis*-12 octadecenoic acid (98.1%), *trans*-10 octadecenoic acid (89.9% for cell culture and 85.1% for animal feeding study), 11-(2'-(*n*-pentyl)phenyl)-10-undecylenic acid (P-110, 95%, E:Z about 7:3), *trans*-10,*cis*-12/*cis*-9,*trans*-11-octadecadien-1-ol (CLA-Alc., 95%), *trans*-10,*cis*-12/*cis*-9,*trans*-11-octadecadienamide (CLA-amide, 95%), 1-chloro-*trans*-10,*cis*-12-octadecadiene (CLA-Cl, 95%), and *cis*-12-octadecen-10-ynoic acid (10y,*cis*-12, 94.5%) were prepared by chemical synthesis by Dr. Sih and colleagues, at the Department of Pharmacy, University of Wisconsin–Madison. CLA was prepared as described [6]; the composition of CLA was 45.7% *cis*-9,*trans*-11, 47.6% *trans*-10,*cis*-12, 1.71% *trans*,*trans*, 3.04% other isomer. In addition, *cis*-9,*trans*-11 (96.3%, 2.6% *trans*,*trans* isomer, and 1.08% others) and *trans*-9,*trans*-11 (100%) CLA were purchased from Matreya Inc. (Pleasant Gap, PA). Natural Lipids (Hovdebygd, Norway) kindly provided *trans*-10,*cis*-12 CLA (92.8% *trans*-10,*cis*-12, 1.61% *cis*-9,*trans*-11, 1.16% *trans*,*trans*, and 1.64% others). We purchased *cis*-10 (99%), and *trans*-10 (99%) heptadecenoic acid and *cis*-10 (99%), and *trans*-10 (99%) nonadecenoic acid from Nu-Chek Prep Corporation (Elysian, MN), and purchased *trans*-9 (100%), *trans*-11 (100%), *trans*-12 (100%), *trans*-13 (100%), *cis*-9 (99%), *cis*-11 (100%) and *cis*-13 (100%) octadecenoic acids from Sigma Chemical Co. (St. Louis, MO). Purity was checked by gas chromatography using a Hewlett-Packard 5890 series II chromatograph (Hewlett-Packard, Andover, MA) fitted with a flame ionization detector and 3396A integrator. A Supelcowax-10 fused silica capillary column (60 m × 0.32 mm, inner diameter film thickness 0.25 μm) was used and oven temperature was programmed from 50° to 190°C, increased 20°C/min, held for 40 minutes, increased 10°C/min to 220°C, and held for 20 minutes.

2.2. Cell culture

The 3T3-L1 preadipocytes were cultured as described [7]. Briefly, 3T3-L1 preadipocytes were grown to confluence at 37°C in Dulbecco's modified Eagle's medium (DMEM) containing 10% fetal bovine serum (FBS). At 2 days postconfluence (designated day 0), cell differentiation was induced with a mixture of methylisobutylxanthin (0.5 mmol/L), dexamethasone (0.25 μmol/L), and insulin (1 μg/mL) in DMEM containing 10% FBS. On day 2 this medium was replaced with

medium containing 10% FBS and insulin only. On day 4 and thereafter the medium consisted of DMEM plus 10% FBS only; this medium was subsequently replaced with fresh medium at 2-day intervals. Fatty acid–albumin complexes were prepared as described [2] and added to culture media for 48 hours at day 6, and cells were harvested at day 8. Final concentrations of fatty acids are indicated in each figure legend. All dishes including control had a final concentration of 100 μmol/L albumin. For the experiments with CLA derivatives, test compounds as well as CLA were dissolved in ethanol. In these experiments, control and treatment dishes all contained the same concentration of albumin (with no fatty acid) and ethanol (final concentration of 0.3%). At harvest, medium was collected, centrifuged to remove cell debris, and used for glycerol release analysis. Cells were then washed three times with phosphate buffered saline (PBS) and incubated with heparin containing DMEM (10 U/mL) for 1 hour, and the media was used for determining HR-LPL activity. Subsequently, cells were washed three times with PBS, scraped, and sonicated to give homogenous samples for TG determination. LPL activity (EC 3.1.1.34) was measured as described [8]. Briefly, concentrated substrate was prepared by combining 300 mg triolein, 18 mg lecithin, and 200 μL of [³H] triolein in 5 mL glycerol and homogenizing. Fresh assay substrate was prepared by mixing 1 part concentrated substrate, 4 parts 0.2 mol/L Tris/HCl (pH 8.0) with 3% bovine serum albumin, and 1 part heat inactivated fasted rat serum. The reaction was started by adding 100 μL of samples to the same volume of assay substrate. After incubating at 37°C for 15 minutes, the assay was stopped by adding 3.25 mL methanol:chloroform:heptane solution (1.41:1.25:1). Subsequently, 1.05 mL 0.1 mol/L potassium carbonate/borate buffer (pH 10.5) was added, and the tubes were shaken vigorously and centrifuged at 3000 rpm for 20 minutes. A 200-μL quantity of the upper layer was used for radioactivity count. Nonreacted assay substrate was also used to get total count from each experiment. Recovery of free fatty acid was estimated at 71% using [¹⁴C] linoleic acid.

Free and esterified glycerol were determined in the medium and cell sonicates using a Sigma Diagnostic Kit (GPO Trinder, St. Louis, MO). This kit used a two-step process; free glycerol was measured without lipase, then lipase was added to hydrolyse esterified glycerol. The difference between the amounts of free and total glycerol is esterified glycerol. Protein was determined in LPL and TG samples using Bio-Rad DC Protein assay kit (Hercules, CA).

2.3. Animal studies and body composition analyses

Male retired breeder ICR mice and semipurified diet (TD94060, 99% basal mix) were purchased from Harlan Sprague-Dawley and Harlan Teklad (Madison, WI), respectively. The diet was composed as follows (ingredient, g/kg): sucrose, 476; casein, vitamin-free test, 210; corn starch, 150; DL-methionine, 3; corn oil 55; cellulose, 50; mineral mix, AIN-76, 35; vitamin mix, AIN-76A, 10; calcium car-

Table 1
Effects of *trans*-10 octadecenoic acid on body composition in mice

	ECW (g)	% Fat	% Water	% Protein	% Ash
Control	38.2 ± 1.3	12.71 ^a ± 1.69	59.3 ^{ab} ± 1.2	20.83 ^{ab} ± 0.62	3.70 ± 0.06
t10,c12 CLA	37.8 ± 1.9	7.67 ^b ± 1.11	63.0 ^a ± 0.8	21.77 ^a ± 0.27	3.88 ± 0.16
t10-18:1	40.4 ± 1.2	15.33 ^a ± 1.50	57.9 ^b ± 1.3	19.74 ^b ± 0.41	3.88 ± 0.19

Twenty male retired breeder ICR mice were used. After a 5-day adaptation period mice were randomly separated into three groups and fed control diet, diet supplemented with 0.27% *trans*-10,*cis*-12 CLA (t10,c12 CLA), or diet supplemented with 0.27–0.29% *trans*-10 octadecenoic acid (t10-18:1) for 3 weeks. Numbers are mean ± SE (*n* = 5). Means with different superscript letters in each column are significantly different at *P* < 0.05. CLA = conjugated linoleic acid; ECW = empty carcass weight.

bonate, 4; choline bitartrate, 2; and ethoxyquin, 0.1. The mineral mix (AIN-76) was composed as follows (ingredient, g/kg): calcium phosphate (dibasic), 500; potassium citrate (monohydrate), 220; sodium chloride, 74; potassium sulfate, 52; magnesium oxide, 24; ferric citrate, 6.0; manganese carbonate, 3.5; zinc carbonate, 1.6; cupric carbonate, 0.3; chromium potassium sulfate, 0.55; potassium iodate, 0.01; sodium selenite, 0.01; and sucrose, 118. The vitamin mix (AIN-76A) was composed as follows (ingredient, g/kg): thiamin HCl, 0.6; riboflavin, 0.6; pyridoxine HCl, 0.7; niacin, 3.0; calcium pantothenate, 1.6; folic acid, 0.2; biotin, 0.02; vitamin B₁₂ (0.1% trituration in mannitol), 1.0; dry vitamin A palmitate (500,000 U/g), 0.8; dry vitamin E acetate (500 U/g), 10.0; vitamin D₃ trituration (400,000 U/g), 0.25; menadione sodium bisulfite complex, 0.15, and sucrose, 981.08.

Supplemental CLA (5 g/kg) was added to diets at the expense of corn oil. Diet was stored at −20°C until use. Mice were housed individually in wire-bottomed cages in a windowless room with a 12-h light-dark cycle in strict accordance with guidelines established by the Research Animal Resources Center of University of Wisconsin–Madison. Diet and water, available ad libitum, were freshly provided three times per week. After a 5-day adaptation period, mice were randomly separated into groups and fed control diet, diet supplemented with *trans*-10,*cis*-12 CLA (0.27%), or diet supplemented with *trans*-10 octadecenoic acid (0.29%) for 3 weeks.

For body composition analyses, animals were sacrificed, gut contents were removed (to obtain empty carcass weight [ECW]), and the carcasses frozen at −20°C. Frozen carcasses were chopped and freeze-dried to determine water content. Each dried carcass was ground to give a homogeneous sample before further analysis. Total nitrogen was analyzed by the Kjeldahl method [9] by the Department of Soil Science, University of Wisconsin–Madison. Carcass fat content was measured by extraction with diethyl ether overnight using a Soxhlet apparatus. Total ash content was determined by incineration (500–600°C, overnight).

2.4. Statistical analyses

One-way analysis of variance was performed on data presented in Table 1, and two-way analysis of variance

(treatments and experiments) was performed on data depicted in the figures (individually cited later here). Of major interest here are the comparisons among the treatments; these were computed using the Statistical Analysis System (SAS Institute Inc., Cary, NC) with the general linear mean procedure and least square means options. If the interaction between treatment and experiment was significant, this interaction was then used as the error term in the least square means analysis.

3. Results

The concentrations used in this report are based on the finding that sera CLA levels of rats fed diet supplemented with 0.5% CLA for 28 days were 72 μmol/L (range 23–120 μmol/L) [2]. Thus in this study, we used 50 μmol/L for single isomers and 100 μmol/L for mixed isomers.

Experiments were done in multiple sets of quadruplicates. Figures 1A–1D show inhibition of HR-LPL activity by various 18-carbon fatty acids. As previously reported, the *trans*-10,*cis*-12 isomer of CLA consistently and significantly reduced HR-LPL activity in 3T3-L1 adipocytes [3]. This effect is shown in all four sets of experiments and used as a negative control. In contrast the monounsaturated octadecenoic acids (18:1) with double bonds at *cis*-9, *cis*-13, *trans*-12, *trans*-13 (Fig. 1A), *cis*-11, *trans*-11, *trans*-9 (Fig. 1B), *cis*-12 (Fig. 1C), or *trans*-10 (Fig. 1D) exhibited no effects on HR-LPL activity. In addition the linoleic acid isomer *trans*-9,*cis*-12 18:2 (Fig. 1D), 10*y*,*cis*-12 (an 18-carbon structure similar to *trans*-10,*cis*-12 CLA but with a triple bond instead of a double bond at C:10) (Fig. 1C), and P-110 (11-(2'-(*n*-pentyl)phenyl)-10-undecylenic acid) (a *trans*-10,*cis*-12 CLA related structure in which the *trans*-10 double bond is part of a benzene ring) did not exhibit any effect on HR-LPL activity (Fig. 1B). The effect of *cis*-9 18:1 (oleic acid) was marginal in that it enhanced HR-LPL activity in three experiments but induced no significant effect in three comparable experiments.

To determine the effects of these CLA cognates on lipolysis and fat deposition, respectively, we measured glycerol release into media and the amount of TG within cells. As previously reported [3], *trans*-10,*cis*-12 CLA increased lipolysis in this model. The 18:1 compounds with double




Fig. 1. Effects of 18-carbon fatty acids on heparin-releasable lipoprotein lipase (HR-LPL) activity in 3T3-L1 adipocytes. At day 6 of differentiation, cells were treated with these compounds as an albumin complex for 48 hours. All dishes including control contained the same concentration of albumin (100 μ mol/L). Numbers are mean \pm SE, $n = 23$ –24 for (A) from six independent experiments, $n = 10$ –12 for (B) and (C) from three independent experiments, and $n = 7$ –8 for (D) from two independent experiments. Means with different letters in each figure are significantly different at $P < 0.05$. Abbreviations and final concentrations used are as follows: (A) 10:0,12:0 CLA, *trans*-10,*cis*-12 conjugated linoleic acid, 46 μ mol/L; 9:18:1, *cis*-9 octadecenoic acid, 50 μ mol/L; 13:18:1, *cis*-13 octadecenoic acid, 50 μ mol/L; 12:18:1, *trans*-12 octadecenoic acid, 50 μ mol/L; 13:18:1, *trans*-13 octadecenoic acid, 50 μ mol/L; (B) P-t10, 11-(2'-(*n*-pentyl)phenyl)-10-undecylenic acid, 48 μ mol/L; 11:18:1, *cis*-11 octadecenoic acid, 50 μ mol/L; 11:18:1, *trans*-11 octadecenoic acid, 50 μ mol/L; 9:18:1, *trans*-9 octadecenoic acid, 50 μ mol/L; (C) 10y,12:2, *cis*-12-octadecen-10-ynoic acid, 47 μ mol/L; 12:18:1, *cis*-12 octadecenoic acid, 49 μ mol/L; and (D) 10:18:1, *trans*-10 octadecenoic acid, 45 μ mol/L; 9:12:18:2, *trans*-9,*cis*-12 octadecadienoic acid, 32 μ mol/L.

bonds at the *trans*-12 (Fig. 2A), *cis*-11, *trans*-11, or *trans*-9 (Fig. 2B) had no effect on glycerol release. However, the 18:1 compounds with double bonds at *cis*-9 (Fig. 2A) reduced glycerol release, whereas 18:1 with double bonds at *cis*-13, *trans*-13 (Fig. 2A) (Fig. 2C) or *trans*-10 positions (Fig. 2D) enhanced glycerol release. *Trans*-9, *cis*-12 18:2 had no effect on glycerol release (Fig. 2D). P-t10 increased (Fig. 2B) and 10y, *cis*-12 (Fig. 2C) decreased glycerol release in this system. Hence the structural feature or features involved in regulating lipolysis appear to be complex.

Total intracellular TG was reduced by *trans*-10,*cis*-12 CLA as reported previously [3]. All 18-carbon monounsaturated fatty acids tested had no effect on total TG (Fig. 3A–3D). No effect on TG was observed by treating with 10y, *cis*-12 (Fig. 3C), P-t10 (Fig. 3B), or *trans*-9, *cis*-12 18:2 (Fig. 3D).

We also tested CLA derivatives that have functional groups other than carboxylic acid. Compounds with alcohol, amide, and chloride groups were prepared and were tested for effects on HR-LPL activity, glycerol release, and intra-




Fig. 2. Effects of 18-carbon fatty acids on glycerol release in 3T3-L1 adipocytes. At day 6 of differentiation, cells were treated with these compounds as an albumin complex for 48 hours. At the end of incubation, culture medium was collected and analyzed for free glycerol after centrifugation. All dishes including control contained the same concentration of albumin (100 μ mol/L). Numbers are mean \pm SE, $n = 23$ –24 for (A) from six independent experiments, $n = 10$ –12 for (B) and (C) from three independent experiments, and $n = 7$ –8 for (D) from two independent experiments. Means with different letters in each figure are significantly different at $P < 0.05$. For abbreviations and final concentrations used, see legend to Fig. 1.

cellular TG. Since these compounds do not have the carboxylic group needed to complex with albumin, they were dissolved in ethanol before addition to the culture medium. The HR-LPL activity in cells treated with CLA dissolved in ethanol was not significantly different than when CLA was treated complexed to albumin (Fig. 4B). The alcohol form of CLA slightly reduced HR-LPL activity while the amide form of CLA (Fig. 4A), as well as the chloride form of CLA (Fig. 4B) had no effect on HR-LPL activity.

Glycerol release was increased by CLA and reduced by the alcohol and amide forms of CLA (Fig. 5A). The chloride form of CLA did not have any effect on glycerol release (Fig. 5B). Total TG was reduced by *trans*-10,*cis*-12 CLA as well as the alcohol form of CLA, but not by the amide and chloride forms of CLA (Figs. 6A and 6B).

Of the 17- and 19-carbon monounsaturated fatty acids with a double bond at the 10th position (either *cis* or *trans* configuration) that were studied, only *cis*-10 17:1 inhibited HR-LPL activity, but it was less active than *trans*-10,*cis*-12 CLA (Fig. 7A). In addition, *cis*-10 17:1 increased glycerol release whereas *trans*-10 17:1, *cis*-10 19:1 or *trans*-10 19:1, decreased glycerol release (Fig. 7B). *Cis*-10 17:1 and *cis*-10 19:1 slightly reduced cellular TG compared to control, but *trans*-10 17:1 and *trans*-10 19:1 were without significant effect (Fig. 7C).

Since *trans*-10 18:1 exhibited an effect on glycerol release, we also tested whether *trans*-10 octadecenoic acid




Fig. 3. Effects of 18-carbon fatty acids on esterified glycerol in 3T3-L1 adipocytes. At day 6 of differentiation, cells were treated with these compounds as an albumin complex for 48 hours. All dishes including control contained the same concentration of albumin (100 $\mu\text{mol/L}$). Cells were scraped and sonicated to determine triacylglyceride (as esterified glycerol) and protein. Numbers are mean \pm SE, $n = 23$ –24 for (A) from 6 independent experiments, $n = 10$ –12 for (B) and (C) from three independent experiments, and $n = 7$ –8 for (D) from two independent experiments. Means with different letters in each figure are significantly different at $P < 0.05$. For abbreviations and final concentrations used, see legend to Fig. 1.

might induce body compositional changes similar to those induced by the *trans*-10,*cis*-12 CLA isomer (Table 1). At the end of the experimental period the group fed diet sup-




Fig. 4. Effects of CLA derivatives on heparin releasable lipoprotein lipase (HR-LPL) activity in 3T3-L1 adipocytes. At day 6 of differentiation, cells were treated with these compounds for 48 hours. Except for t10,c12-CLA, all treatment compounds were dissolved in ethanol. All dishes including t10,c12-CLA and control contained the same concentration of albumin (100 $\mu\text{mol/L}$) and ethanol (0.3% final concentration). Numbers are mean \pm SE, $n = 6$ –8 from two independent experiments. Means with different letters in each figure are significantly different at $P < 0.05$. Abbreviations and final concentrations used are as follows: (A) CLA, *trans*-10,*cis*-12 (48%) and *cis*-9,*trans*-11 (46%) isomers, 98 $\mu\text{mol/L}$; CLA-Alc., *trans*-10,*cis*-12/*cis*-9,*trans*-11-octadecadien-1-ol, 95 $\mu\text{mol/L}$; CLA-amide, *trans*-10,*cis*-12/*cis*-9,*trans*-11-octadecadienamide, 95 $\mu\text{mol/L}$; and (B) t10,c12-CLA, *trans*-10,*cis*-12 conjugated linoleic acid, 46 $\mu\text{mol/L}$. Et indicates dissolved in ethanol before treating to cells; CLA-CI, 1-chloro-*trans*-10,*cis*-12-octadecadiene, 48 or 96 $\mu\text{mol/L}$.




Fig. 5. Effects of CLA derivatives on glycerol release in 3T3-L1 adipocytes. At day 6 of differentiation, cells were treated with these compounds as an albumin complex for 48 hours. Except for t10,c12-CLA, all treatment compounds were dissolved in ethanol. All dishes including t10,c12-CLA and control contained the same concentration of albumin (100 $\mu\text{mol/L}$) and ethanol (0.3% final concentration). At the end of incubation culture medium was collected and analyzed for free glycerol after centrifugation. Numbers are mean \pm SE, $n = 6$ –8 from two independent experiments. Means with different letters in each figure are significantly different at $P < 0.05$. For abbreviations and final concentrations used, see legend to Fig. 4.

plemented with *trans*-10,*cis*-12 CLA exhibited significantly reduced body fat (40% reduction compared to control), whereas whole body water was significantly enhanced and whole body protein, and ash were apparently enhanced. By contrast, the group fed *trans*-10 octadecenoic acid exhibited no significant change in any of these measurements (Table 1). There were no significant difference in mean body weights (control group, 40.5 ± 1.2 g; *trans*-10,*cis*-12 CLA-fed group, 40.3 ± 2.2 g; and t10 18:1-fed group, 42.6 ± 1.3 g) or mean total feed intake (control group, 91.2 ± 4.1 g; *trans*-10,*cis*-12 CLA-fed group, 83.7 ± 2.4 g; and t10 18:1-fed group, 89.8 ± 4.2 g).




Fig. 6. Effects of CLA derivatives on esterified glycerol in 3T3-L1 adipocytes. At day 6 of differentiation, cells were treated with these compounds for 48 hours. Except for t10,c12-CLA, all treatment compounds were dissolved in ethanol. All dishes including t10,c12-CLA and control contained the same concentration of albumin (100 $\mu\text{mol/L}$) and ethanol (0.3% final concentration). Cells were scraped and sonicated to determine triacylglyceride (as esterified glycerol) and protein. Numbers are mean \pm SE, $n = 6$ –8 from two independent experiments. Means with different letters in each figure are significantly different at $P < 0.05$. For abbreviations and final concentrations used, see legend to Fig. 4.




Fig. 7. Effects of odd-numbered fatty acids on heparin-releasable lipoprotein lipase (HR-LPL) activity (A), glycerol release (B), and cellular esterified glycerol (C) in 3T3-L1 adipocytes. At day 6 of differentiation, cells were treated with 50 μ M of fatty acid for 48 hours. All treatments had 100 μ M of albumin. Reported values are mean \pm SE ($n = 12$, collected from three independent experiments). Means with different letters in each figure are significantly different at $P < 0.05$. CLA, conjugated linoleic acid; c, *cis*; t, *trans*; 17:1, heptadecenoic acid; 19:1, nonadecenoic acid.

4. Discussion

Table 2 summarizes the results of this study along with those of previously published investigations [2,3,10–12]. Among octadecadienoic acids only *trans*-10,*cis*-12 CLA reduced the activity of HR-LPL, whereas *cis*-9,*trans*-11, *trans*-9,*trans*-11, linoleic acid, or *trans*-9,*cis*-12 octadecadienoic acid did not reduce HR-LPL activity. Various 18-carbon monounsaturated fatty acids with double bonds located between carbons 9 and 13, in either *trans* or *cis* configurations, did not inhibit HR-LPL activity. CLA amide and chloride derivatives were without effect, whereas CLA alcohol exhibited a modest inhibitory effect that was less than that seen with *trans*-10,*cis*-12 CLA. We propose that this is due to the conversion of the alcohol form to free acid form; hence the effect of the alcohol form of CLA is less active than CLA itself. The data support the conclusion that the *trans*-10,*cis*-12 conjugated double bond in conjunction with a carboxyl group at C-1 is required for inhibition of HR-LPL activity.

The TG content in fat cells reflects the balance between fat uptake by HR-LPL, lipogenesis, and lipolysis. Our data consistently show that any compound that reduced HR-LPL activity in adipocytes also reduced body fat in animals. Since the amount of protein secreted into heparin containing medium was not different between treatment groups, it is unlikely that the effect of CLA on HR-LPL inhibition results from less protein secretion. Therefore the effects of CLA on HR-LPL can be due to regulation of HR-LPL expression, modification of location of HR-LPL, or interaction with HR-LPL directly.

Unlike its consistent inhibition of HR-LPL activity, the

effects of CLA on intracellular LPL activity vary, either inhibiting [13] or having no effect [12]. We were able to observe inhibitory effects of CLA (as a mixture of *trans*-10,*cis*-12 and *cis*-9,*trans*-11 isomers) on HR-LPL activity in as short a time as 20 minutes (unpublished observations, [14]). This appears to be too rapid for a transcriptional effect; the effect of CLA on HR-LPL activity may result in part from direct interaction between *trans*-10,*cis*-12 CLA and LPL, similar to the inhibition mechanism of CLA on stearoyl CoA desaturase (SCD, Δ -9 desaturase) [4]. Alternatively, CLA might inhibit LPL enzyme relocation to membrane, making less LPL available for the activity. In addition, we cannot rule out the possibility that CLA may further interact at the transcriptional level, which has been reported by others, for example reduced LPL mRNA in 3T3-L1 adipocytes or adipose tissues by *trans*-10,*cis*-12 CLA [14–16].

Previously, Lin et al. [12] reported that both *trans*-10,*cis*-12 and *cis*-9,*trans*-11 CLA isomers reduced HR-LPL activity from 3T3-L1 adipocytes, although the concentrations were much higher than those we used in this study. When similar concentrations were compared there was no effect by *cis*-9,*trans*-11 CLA, whereas *trans*-10,*cis*-12 CLA effectively reduced HR-LPL activity.

CLA may also affect lipogenesis in adipocytes. Previously our group reported that CLA did not affect activities of acetyl-CoA carboxylase and fatty acid synthetase (key enzymes involved in lipogenesis), both in vivo and during 1 week of feeding [13] and in 3T3-L1 adipocytes [17]. This is consistent with observations by Bee [18]. However, others report that CLA reduced activities and/or mRNA level of these two enzymes in adipocytes [15,16,19–21]. Thus CLA may result in less TG in adipocytes by reducing lipogenesis; however, we did not measure lipogenesis in these experiments.

Glycerol release represents lipolysis in adipocytes. Our results here indicate that glycerol release did not correlate well with TG content in cells or with reduced body fat in vivo. This may be due to the sensitivity of the method used in our experiments. However, nordihydroguaiaretic acid (NDGA) and AA861, both well known inhibitors of lipoxygenase and hormone-sensitive lipase, decreased lipolysis in our experiments, as expected, compared to control (as μ g glycerol/mL: control 181 ± 8 , NDGA (1×10^{-5} mol/L) 131 ± 9 , AA861 (1×10^{-6} mol/L) 150 ± 22 , and CLA 237 ± 11 , $n = 7$ –8 from two independent experiments). This suggests that this method represents the effects of treatment in this model. This was accompanied by reduced TG (as μ g/mg protein: control 153 ± 13 , NDGA (1×10^{-5} mol/L) 93 ± 10 , AA861 (1×10^{-6} mol/L) 95 ± 7 , and CLA 94 ± 6 , $n = 7$ –8 from two independent experiments) and reduced body fat in mice for NDGA (for experimental conditions, see [22]). However, compounds that increased glycerol release only did not result in decreased TG, for example, *trans*-10 18:1 effectively increased glycerol release without affecting HR-LPL activity in 3T3-L1 adipocytes and had no

Table 2
Summary of compounds used

Compound	Inhibition of HR-LPL	Glycerol release	Total TG
Octadecadienoic acid			
<i>trans</i> -10, <i>cis</i> -12 CLA	+++	↑	↓
<i>cis</i> -9, <i>trans</i> -11 CLA [3,12]	– [3]/++ [12]	↑	–
<i>trans</i> -9, <i>trans</i> -11 CLA [3]	–	↑	–
<i>cis</i> -9, <i>cis</i> -12 octadecadienoic acid (linoleic acid) [2,3]	–	↑	↑
<i>trans</i> -9, <i>cis</i> -12 octadecadienoic acid	–	–	–
Octadecenoic acid			
<i>cis</i> -9 octadecenoic acid (oleic acid)	–	↓	–
<i>cis</i> -11 octadecenoic acid	–	–	–
<i>cis</i> -12 octadecenoic acid	–	↑	–
<i>cis</i> -13 octadecenoic acid	–	↑	–
<i>trans</i> -9 octadecenoic acid	–	–	–
<i>trans</i> -10 octadecenoic acid	–	↑ ↑	–
<i>trans</i> -11 octadecenoic acid (vaccenic acid)	–	–	–
<i>trans</i> -12 octadecenoic acid	–	–	–
<i>trans</i> -13 octadecenoic acid	–	↑	–
Eicosadienoic acid			
<i>Cis</i> -11, <i>cis</i> -14 eicosadienoic acid [10]	–	–	–
<i>Cis</i> -11, <i>trans</i> -13 and <i>trans</i> -12, <i>cis</i> -14 CEA [10]	++	↑	↓
CLA derivatives			
<i>trans</i> -10, <i>cis</i> -12/ <i>cis</i> -9, <i>trans</i> -11-octadecadien-1-ol (CLA-Alc.)	+	↓	↓
<i>trans</i> -10, <i>cis</i> -12/ <i>cis</i> -9, <i>trans</i> -11-octadecadienamide (CLA-Armide)	–	↓	–
1-chloro- <i>trans</i> -10, <i>cis</i> -12-octadecadiene (CLA-Cl)	–	–	–
Odd-numbered fatty acids			
<i>cis</i> -10 heptadecenoic acid	+	↑	↓
<i>trans</i> -10 heptadecenoic acid	–	↓	–
<i>cis</i> -10 nonadecenoic acid	–	↓	↓
<i>trans</i> -10 nonadecenoic acid	–	↓	–
<i>Cis</i> -10, <i>cis</i> -13 nonadecadienoic acid [11]	–	↑	–
<i>Cis</i> -10, <i>trans</i> -12 and <i>trans</i> -11, <i>cis</i> -13 CNA [11]	++++	↑	↓
Others			
<i>cis</i> -12-octadecen-10-ynoic acid (10y, <i>cis</i> -12)	–	↓	–
11-(2'-(<i>n</i> -pentyl)phenyl)-10-undecylenic acid (<i>P-110</i>)	–	↑	–

– = effect; + = inhibition; ↑ = increase; ↓ = decrease.

No numbers in square brackets indicate references from which data are collected.

effect on total TG or body fat in mice. This suggests that this lipolysis assay method may not be sensitive enough to enable us to observe the difference in TG; thus further studies are need to clarify the correlation between lipolysis and TG in this cell culture model.

Previously we reported that the *trans*-10,*cis*-12 CLA isomer effectively inhibited hepatic SCD [4]. In addition, *trans*-10,*cis*-12 CLA is also responsible for decreased apo B secretion in HepG2 cells, although a *trans*-10 double bond is a key factor for this activity [5]. It was of interest that odd-numbered fatty acids such as CNA [11] can inhibit HR-LPL activity as well as reduce the TG in the cell. Since CNA and CLA cannot share common metabolites because of the difference of one carbon, we proposed that CLA itself was directly effective. In this study, we observed that *cis*-10 17:1 inhibits HR-LPL activity, although less so than *trans*-10,*cis*-12 CLA, indicating that a double bond at the 10th position may play an important role in these activities.

It appears, then, that *trans*-10,*cis*-12 CLA is directly responsible for reduction of HR-LPL activity, and that the

trans-10 double bond may be the key. In contrast, either a *trans*-10 or *cis*-12 bond may induce glycerol release from adipocytes. This study also suggests that lipolysis may not be a good indicator of predicting TG content in the 3T3-L1 adipocyte model; rather, inhibition of HR-LPL activity may provide a better prediction of TG content in this cell line and fat level in animal models.

Acknowledgments

This research was supported in part by gift funds administered through the University of Wisconsin–Madison Food Research Institute. We thank Dr. Sih at the Department of Pharmacy and Dr. Ken Ku at the Food and Drug Administration for preparing some of the compounds tested. Three of the authors (YP, MEC, MWP) are inventors of CLA use patents that are assigned to the Wisconsin Alumni Research Foundation.

References

- [1] Pariza MW, Park Y, Cook ME. The biologically active isomers of conjugated linoleic acid. *Prog Lipid Res* 2001;40:283–8.
- [2] Park Y, Albright KJ, Storkson JM, Cook ME, Pariza MW. Effect of conjugated linoleic acid on body composition in mice. *Lipids* 1997;32:853–8.
- [3] Park Y, Albright KJ, Storkson JM, Liu W, Pariza MW. Evidence that the *trans*-10,*cis*-12 isomer of conjugated linoleic acid induces body composition changes in mice. *Lipids* 1999;34:235–41.
- [4] Park Y, Storkson JM, Ntambi JM, Cook ME, Sih CJ, Pariza MW. Inhibition of hepatic stearoyl-CoA desaturase activity by *trans*-10,*cis*-12 conjugated linoleic acid and its derivatives. *Biochim Biophys Acta* 2000;1486:285–92.
- [5] Storkson JM, Park Y, Cook ME, Pariza MW. Effects of *trans*-10,*cis*-12 conjugated linoleic acid and cognates on apolipoprotein B secretion in HepG2 cells. *FASEB J* 2001;15:A290.
- [6] Chin SF, Liu W, Storkson JM, Ha YL, Pariza MW. Dietary sources of conjugated dienoic isomers of linoleic acid, a newly recognized class of anticarcinogens. *J Food Comp Anal* 1992;5:185–97.
- [7] Frost SC, Lane MD. Evidence for the involvement of vicinal sulfhydryl groups in insulin-activated hexose transport by 3T3-L1 adipocytes. *J Biol Chem* 1985;260:2646–52.
- [8] Nilsson-Ehle P, Schotz MC. A stable, radioactive substrate emulsion for assay of lipoprotein lipase. *J Lipid Res* 1976;17:536–41.
- [9] Helrich K. Official methods of analysis, 15th ed. Arlington, VA: Association of Official Analytical Chemists, 1990. p. 935–7.
- [10] Pariza MW, Park Y, Albright KJ, Liu W, Storkson JM, Cook ME. Synthesis and biological activities of conjugated eicosadienoic acid. Abstracts for AOCS Annual Meeting & Expo 1998. p. 21.
- [11] Park Y, Pariza MW. The effects of dietary conjugated nonadecadienoic acid (CNA) on body composition in mice. *Biochim Biophys Acta* 2001;1533:171–4.
- [12] Lin Y, Kreeft A, Schuurbiens JAE, Draijer R. Different effects of conjugated linoleic acid isomers on lipoprotein lipase activity in 3T3-L1 adipocytes. *J Nutr Biochem* 2001;12:183–9.
- [13] Xu X, Storkson J, Kim S, Sugimoto K, Park Y, Pariza MW. Short-term intake of conjugated linoleic acid inhibits lipoprotein lipase and glucose metabolism but does not enhance lipolysis in mouse adipose tissue. *J Nutr* 2003;133:663–7.
- [14] Kang K, Liu W, Albright KJ, Park Y, Pariza MW. *trans*-10,*cis*-12 CLA inhibits differentiation of 3T3-L1 adipocytes and decreases PPAR- γ expression. *Biochem Biophys Res Comm* 2003;303:795–9.
- [15] Kang K, Albright KJ, Pariza MW. Conjugated linoleic acid decreases peroxisome proliferators-activated receptor γ and adipogenic mRNA expression in mice. *FASEB J* 2003;17:A1161.
- [16] Baumgard LH, Matitashvili E, Corl BA, Dwyer DA, Bauman DE. *trans*-10,*cis*-12 Conjugated linoleic acid decreases lipogenic rates and expression of genes involved in milk lipid synthesis. *J Dairy Sci* 2002;85:2155–63.
- [17] Park Y. Regulation of energy metabolism and the catabolic effects of immune stimulation by conjugated linoleic acid. Ph.D. Thesis. Madison, WI: University of Wisconsin, 1996.
- [18] Bee G. Dietary conjugated linoleic acids affect tissue lipid composition but not de novo lipogenesis in finishing pigs. *Anim Res* 2001;50:383–99.
- [19] Hsu JM, Ding ST. Effects of polyunsaturated fatty acids on the expression of transcription factor adipocyte determination and differentiation-dependent factor 1 and of lipogenic and fatty acid oxidation enzymes in porcine differentiating adipocytes. *Br J Nutr* 2003;90:507–13.
- [20] Corino C, Mourot J, Magni S, Pastorelli G, Rosi F. Influence of dietary conjugated linoleic acid on growth, meat quality, lipogenesis, plasma leptin and physiological variables of lipid metabolism in rabbits. *J Anim Sci* 2002;80:1020–8.
- [21] Piperova LS, Teter BB, Bruckental I, Sampugna J, Mills SE, Yurawecz MP, Fritsche J, Ku K, Erdman RA. Mammary lipogenic enzyme activity, *trans* fatty acids and conjugated linoleic acids are altered in lactating dairy cow fed a milk fat-depressing diet. *J Nutr* 2000;130:2568–74.
- [22] Park Y, Pariza MW. Lipoxigenase inhibitors enhance body fat reduction in mice by conjugated linoleic acid and inhibit heparin-releasable lipoprotein lipase activity in 3T3-L1 adipocytes. *Biochim Biophys Acta* 2001;1534:27–33.

Inhibition of hepatic stearoyl-CoA desaturase activity by *trans*-10,*cis*-12 conjugated linoleic acid and its derivatives

Yeonhwa Park ^a, Jayne M. Storkson ^a, James M. Ntambi ^{b,c}, Mark E. Cook ^{a,c,d}, Charles J. Sih ^e, Michael W. Pariza ^{a,c,*}

^a Department of Food Microbiology and Toxicology, Food Research Institute, University of Wisconsin-Madison, Madison, WI 53706, USA

^b Department of Biochemistry, University of Wisconsin-Madison, Madison, WI 53706, USA

^c Department of Nutritional Sciences, University of Wisconsin-Madison, Madison, WI 53706, USA

^d Department of Animal Sciences, University of Wisconsin-Madison, Madison, WI 53706, USA

^e Department of Pharmacy, University of Wisconsin-Madison, Madison, WI 53706, USA

Received 29 March 2000; accepted 5 May 2000

Abstract

Conjugated linoleic acid (CLA) has been reported to decrease stearoyl-CoA desaturase (SCD) activity by decreasing mRNA expression. This investigation was designed to determine whether structurally related compounds of CLA have a direct inhibitory effect on SCD activity. *Trans*-10,*cis*-12 CLA had strong inhibitory activity on SCD while *cis*-9,*trans*-11, and *trans*-9,*trans*-11 isomers had no effect. *Trans*-10 octadecenoate was not inhibitory, whereas *cis*-12 octadecenoate was inhibitory, but not as effective as *trans*-10,*cis*-12 CLA. Of the oxygenated derivatives, 9-peroxy-*cis*/*trans*-10, *trans*-12 octadecadienoate was a more effective inhibitor than *trans*-10,*cis*-12 CLA, whereas 9-hydroxy-*trans*-10, *cis*-12 octadecadienoate was less effective. Interestingly, *cis*-11 octadecadienoate and *cis*-12 octadecen-10-ynoate were slightly inhibitory. However, *trans*-9 and *trans*-11 octadecenoates, and *trans*-9,*cis*-12 octadecadienoate were all inactive under test condition, as were linoleate, oleate, and arachidonate. Derivatives of CLA acid modified to alcohol, amide or chloride were all inactive. A *cis*-12 double bond appears to be a key structural feature for inhibiting SCD activity, especially when coupled with a *trans*-10 double, whereas a *cis*-11 double bond is less effective. © 2000 Elsevier Science B.V. All rights reserved.

Keywords: Conjugated linoleic acid; Stearoyl-CoA desaturase; Δ^9 desaturase; *trans*-10,*cis*-12 CLA

1. Introduction

CLA refers to a group of geometric and positional isomers of conjugated linoleic acid. CLA was originally isolated from ground beef as an anticarcinogenic principal which contained mainly the *cis*-9,*trans*-11 isomer [1,2]. This isomer can be formed during the biohydrogenation of linoleic acid to stearic acid by rumen bacteria [3]. Alternatively, the latest research shows that it can form in animals via Δ^9 desaturation of *trans*-11 octadecenoic acid (*trans* vaccenic acid)

Abbreviations: SCD, stearoyl-CoA desaturase; CLA, conjugated linoleic acid; 9-HODE, 9-hydroxy-*trans*-10,*cis*-12 octadecadienoic acid; 9-HOO, 9-hydroperoxy *trans*-10,*cis*-12 octadecadienoic acid; CLA-Alc, *trans*-10,*cis*-12/*cis*-9,*trans*-11-octadecadien-1-ol; CLA-amide, *trans*-10,*cis*-12/*cis*-9,*trans*-11-octadecadienamide; CLA-Cl, 1-chloro-*trans*-10,*cis*-12-octadecadiene; 10y,*cis*-12, *cis*-12-octadecen-10-ynoic acid; AA, arachidonic acid; LA, linoleic acid; ACS, acyl CoA synthetase; LPL, lipoprotein lipase; PUFA, polyunsaturated fatty acid

* Corresponding author. Fax: +1-608-263-1114;
E-mail: mwpariza@facstaff.wisc.edu

[4,5]. CLA typically used in animal studies is prepared by alkali isomerization from pure linoleic acid, and contains more than 95% CLA, mainly *cis*-9,*trans*-11 and *trans*-10,*cis*-12 isomers (85–90%) along with other minor isomers (*trans*,*trans* or *cis*,*cis*). This preparation of CLA has shown a wide range of biologically beneficial activities: reduction of carcinogenesis in several animal models [1,6–8]; decreased severity of atherosclerosis [9,10]; reduction of adverse effects of immune stimulation [11,12]; growth promotion in young rats [13]; and a reduction of body fat and an increase in lean body mass in several animal species [14–18]. However, depending on the source of linoleic acid (pure fatty acid or vegetable oil) and conditions of isomerization, the isomer distribution can vary significantly [19,20]. In some studies, CLA preparations contained a more complex mixture of isomers, such as 8,10 and 11,13 isomers as well as 9,11 and 10,12 isomers [19,20].

Since Ha et al. [6] reported only the *cis*-9,*trans*-11 isomer was incorporated in the phospholipid fraction of mouse forestomach, it has been postulated that *cis*-9,*trans*-11 CLA is the biologically active isomer. More recently, several investigators [17,21,22] reported that all CLA isomers are incorporated in both phospholipids and neutral lipids. However, it was not until 1998 that Lee et al. [23] reported the possible biological activity of the *trans*-10,*cis*-12 isomer on stearoyl-CoA desaturase (SCD). In the first part of that study, they fed CLA (as a mixture of 42% *cis*-9,*trans*-11 and 44% *trans*-10,*cis*-12) to mice and analyzed hepatic SCD mRNA expression, which was significantly decreased by CLA feeding. In the second part of the experiment, H2.35 mouse hepatocytes were tested. CLA generated by rumen bacteria (this preparation contained mainly *cis*-9,*trans*-11 CLA, >60% of total fatty acid, with no *trans*-10,*cis*-12 isomer) did not have any effect on SCD mRNA expression. Based on these observations, Lee et al. [23] suggested that the *trans*-10,*cis*-12 isomer was responsible for the effects of synthetic CLA on SCD. Recently, we clearly demonstrated that the *trans*-10,*cis*-12 isomer was responsible for body composition changes in mice and the decrease in lipoprotein lipase activity in 3T3-L1 adipocytes, while *cis*-9,*trans*-11 and *trans*-9,*trans*-11 isomers did not have any impact [24]. In the current study, we tested the direct effects of CLA on the activity of SCD using

high purity isomers. Other 18-carbon fatty acids, CLA derivatives, as well as arachidonic acid (a known inhibitor of SCD gene) were also compared with isomers of CLA.

2. Materials and methods

[1-¹⁴C]Palmitic acid (50 mCi/mmol) was purchased from American Radiolabeled Chemicals, (St. Louis, MO). 9-hydroxy-, (*E,Z*)-(\pm)-10,12-octadecadienoic acid (9-HODE) was purchased from Cayman (Ann Arbor, MI). 9-Hydroperoxy-10,12 octadecadienoic acid (9-HOO, a mixture of *trans*-10,*cis*-12 and *trans*-10,*trans*-12 in 2:1, 95%), *trans*-9,*cis*-12 octadecadienoic acid (31.7%), *cis*-12 octadecadienoic acid (98.1%), *trans*-10 octadecadienoic acid (89.9%), *trans*-10,*cis*-12/*cis*-9,*trans*-11-octadecadien-1-ol (CLA-Alc, 95%), *trans*-10,*cis*-12/*cis*-9,*trans*-11-octadecadienamide (CLA-amide, 95%), 1-chloro-*trans*-10,*cis*-12-octadecadiene (CLA-Cl, 95%), and *cis*-12-octadecen-10-ynoic acid (10y,*cis*-12, 94.5%) were prepared by chemical synthesis. CLA was prepared as described [2]; the composition of CLA was 45.7% *cis*-9,*trans*-11, 47.6% *trans*-10,*cis*-12, 1.71% *trans*,*trans*, 3.04% other isomer. *Cis*-9,*trans*-11 CLA was purchased from Matreya (Pleasant Gap, PA); 96.3% *cis*-9,*trans*-11/*trans*-9,*cis*-11, 2.6% *trans*,*trans* isomer, and 1.08% others. Natural Lipids (Hovdebygda, Norway) kindly provided *trans*-10,*cis*-12 CLA; 92.8% *trans*-10,*cis*-12, 1.61% *cis*-9,*trans*-11, 1.16% *trans*,*trans*, and 1.64% others. Other fatty acids were purchased from Nu-Chek-Prep, (Elysian, MN). Weanling male ICR mice were purchased from Harlan Sprague Dawley (Madison, WI). Semi-purified diet was purchased from Harlan Teklad (TD96040, Madison, WI). Fat-free carbohydrate diet was from USB (fat-free diet, 15750 PT).

2.1. Preparation of liver microsomal fraction

Mice were housed in a windowless room with a 12-h light–dark cycle in strict accordance with guidelines established by the Research Animal Resources Center of University of Wisconsin-Madison and a protocol approved by the Animal Care Committee. Diet and water were available ad libitum. Animals were fed according to Miller and Ntambi [25]. Mice

were starved for 24 h, refed for 24 h, starved for 24 h, and then fed fat-free carbohydrate diet for 24 h before CO₂ suffocation. Microsomal fractions were prepared from mice livers as described previously [26] with little modification. Liver homogenate was prepared with 4 vols. (w/v) of 0.25 M sucrose, 1 mM EDTA, 10 mM Tris-HCl (pH 7.4) and centrifuged at 12000×g for 15 min. The supernates were re-centrifuged at 100000×g for 1 h, pellets were rinsed once and resuspended before use (microsomal fractions). All preparations were made at 4°C. Three independent preparations of microsomal fractions were used. Protein was determined using Bio-Rad DC Protein assay kit.

The SCD activity was determined as described [27]. The assay system contained 7.2 mM ATP, 0.54 mM CoA, 6 mM MgCl₂, 0.8 mM NADH, 0.1 M potassium phosphate buffer (pH 7.16), 204 nmol [1-¹⁴C]palmitic acid (specific activity 0.98 Ci/mol) in 20 µl ethanol, testing fatty acid complexed with albumin (final concentration of 0–100 µM, as indicated in figure legends) and microsomal fraction (0.5–1.8 mg protein) in 1 ml final volume. The reaction was initiated by the addition of microsomal fraction and incubated at 37°C for 15 min. All fatty acids were dissolved in KOH to complex with albumin before addition to the incubation mixture. 9-HODE, 9-HOO, and CLA derivatives (Fig. 4) were in ethanol solution as indicated in legends. For comparison, CLA was also dissolved in ethanol and tested. For the experiment with acyl CoA synthetase (ACS), ACS (8 mU) was pre-incubated for 5 min with cofactors and fatty acids and then incubated for 5 min with the microsomal fractions. The reaction was stopped by adding 1 ml 12% KOH in ethanol and heating for 45 min at 80°C. Fatty acids were extracted with hexane after acidification with HCl. Fatty acid methyl esters were prepared with 4% HCl/MeOH at 60°C for 20 min, and separated by thin-layer chromatography on silver nitrate-impregnated silica gel G plates [28] using a mixture of hexane:diethyl ether (9:1). To help visualization, cold standards (palmitic and palmitoleic acid methyl esters) were co-spotted with samples. After visualization with 2',7'-dichlorofluorescein, corresponding spots were scraped off the plates, extracted with hexane and subjected to liquid scintillation counting. Enzyme activities were calculated as nmol palmitic

acid converted to palmitoleic acid per min per mg protein.

2.2. Statistics

Because of the experimental structure, the data were divided into six groups so that data were balanced (or close to balanced) with each group. A balanced design means that within a group, the same treatments were used in all experiments in the group. Data were subjected to analysis using the Statistics Analysis System (SAS Users Guide: Statistics, SAS Institute, Cary, NC). Within each group, data were analyzed with PROC MIXED as a two-way analysis of variance with treatment as a fixed factor and experiment and the experiment by treatment interacts as random factors. The error term for comparing treatments was a mix of the residual error and the experiment by treatment interact as determined by PROC MIXED using the Satterthwaith procedure for degrees of freedom [29].

3. Results

Fig. 1 shows the effects of different CLA isomers and linoleic acid derivatives on SCD activity. CLA mixture at 100 µM (either complexed with albumin or dissolved in ethanol) significantly reduced the activity of SCD. Two major isomers of CLA, *cis*-9,*trans*-11 and *trans*-10,*cis*-12, were tested individually at concentrations similar to those in the CLA mixture, 42.6 and 43.6 µM, respectively. Only the *trans*-10,*cis*-12 isomer significantly reduced SCD activity (similar to the CLA mixture), while there was no effect by *cis*-9,*trans*-11 CLA. We also tested linoleic acid derivatives, 9-HODE and 9-HOO, which contained conjugated double bonds at the *trans*-10,*cis*-12 position. Both compounds were inhibitory to SCD, with 9-HOO (51.3 µM) being the more effective while 9-HODE (100 µM) was less active than *trans*-10,*cis*-12 CLA. Ethanol effect was ruled out since CLA treated as albumin complex or dissolved in ethanol showed the same inhibitory effect. To eliminate the possibility that *trans*-10,*cis*-12 CLA may have impacted on acyl CoA formation, additional experiments with acyl CoA synthetase (ACS) were performed. *trans*-10,*cis*-12 isomer was incubated




Fig. 1. Effects of different conjugated linoleic acid (CLA) isomers and linoleic acid derivatives on hepatic stearoyl-CoA desaturase activity. Numbers are mean \pm S.E. ($n=4-6$, collected from three independent experiments). Means with different letters are significantly different ($P<0.05$). The final concentrations were 100 μ M for CLA and 9-HODE, 42.6 μ M for *cis*-9,*trans*-11 CLA, 43.6 μ M for *trans*-10,*cis*-12 CLA, and 51.3 μ M for 9-HOO. CLA, *cis*-9,*trans*-11 CLA, and *trans*-10,*cis*-12 CLA were complexed with albumin. CLA(Et), 9-HODE, and 9-HOO were dissolved in ethanol. All samples contained 100 μ M albumin, including control. CLA, conjugated linoleic acid; 9-HODE, 9-hydroxy-*trans*-10,*cis*-12 octadecadienoic acid; 9-HOO, 9-hydroperoxy-10,12-octadecadienoic acid.

with or without ACS (8 mU) for 5 min before adding microsomal fractions and then incubated for an additional 5 min. Inhibition by this isomer was not affected by addition of ACS (55% inhibition with ACS vs. 52.2% inhibition without ACS), indicating that formation of CoA was not the major process of inhibition by this isomer. The mean of two experiments showed a general linear trend on the inhibition




Fig. 2. Dose response of *trans*-10,*cis*-12 CLA on hepatic stearoyl-CoA desaturase. Reported values are mean \pm S.E. ($n=2-4$, collected from two independent experiments). Means with different letters are significantly different ($P<0.05$). To avoid zero values of log, the value of 1 was added to all data. CLA, conjugated linoleic acid.




Fig. 3. Effects of fatty acids on hepatic stearoyl-CoA desaturase activity. Numbers are mean \pm S.E. ($n=4$, collected from two independent experiments). Means with different letters are significantly different ($P<0.05$). The final concentrations were 50 μ M for *trans*-10,*cis*-12 CLA, *trans*-9,*trans*-11 CLA, *cis*-9,*cis*-12 octadecadienoic acid (linoleic acid), *trans*-9, *trans*-11, *cis*-9, *cis*-11 and *cis*-12 octadecenoic acids, and 10y,*cis*-12, 31.7 μ M for *trans*-9,*cis*-12 octadecadienoic acid, and 49.4 μ M for *trans*-10 octadecenoic acid. CLA, conjugated linoleic acid; LA, linoleic acid; 10y,c12, *cis*-12-octadecen-10-ynoic acid.

of SCD over the dose of *trans*-10,*cis*-12 isomer of CLA tested (Fig. 2).

Fig. 3 shows the effects of other 18-carbon fatty acids including the *trans,trans* isomer of CLA, *trans*-9,*trans*-11 CLA, linoleic acid (*cis*-9,*cis*-12), and *trans*-9,*cis*-12 octadecadienoic acid did not have any effect on SCD. Neither did mono-unsaturated 18-carbon fatty acids, *trans*-9, *trans*-10, *trans*-11 (vaccenic acid), or *cis*-9 (oleic acid) octadecenoic acids. *cis*-11 and *cis*-12 octadecenoic acids inhibited this enzyme, but the effects were less than *trans*-10,*cis*-12 CLA (19% inhibition by *cis*-11 and 51% inhibition by *cis*-12 compared to 70% inhibition by the *trans*-10,*cis*-12 CLA isomer). 10y,*cis*-12 had similar effects as *cis*-12 octadecenoic acid (50% inhibition), but again was less effective than the *trans*-10,*cis*-12 isomer of CLA.

To test if there are any additional requirements for inhibition besides double bond location, we prepared




Fig. 4. Effects of CLA derivatives in hepatic stearoyl-CoA desaturase activity. Numbers are mean \pm S.E. ($n=4$, collected from two independent experiments). Means with different letters are significantly different ($P<0.001$). All final concentrations were 100 μ M. Testing materials including CLA(Et) were dissolved in ethanol, except CLA was complexed to albumin. All samples contained 100 μ M albumin, including control. CLA-Alc, CLA-alcohol form; CLA-amide, CLA-amide form; CLA-Cl, CLA-Cl form. See Section 2 for chemical names of the derivatives.

and tested alcohol, amide or chloride forms of CLA since these compounds have double bonds at the same location, but different functional groups. None of these derivatives showed any effect on SCD (Fig. 4).

To prove the specificity of *trans*-10,*cis*-12 CLA, we also tested arachidonic acid under the same experimental conditions. Polyunsaturated fatty acids, especially arachidonic acid, are known to reduce SCD activity by reducing mRNA expression (similar to previously reported effects of CLA). Fig. 5 shows that, again, the CLA mixture (100 μ M) significantly




Fig. 5. Effects of arachidonic acid on hepatic stearoyl-CoA desaturase activity. Numbers are mean \pm S.E. ($n=4$, collected from two independent experiments). Means with different letters are significantly different ($P<0.05$). The final concentrations were 100 μ M for CLA and 50 and 100 μ M for arachidonic acid. CLA, conjugated linoleic acid; AA, arachidonic acid.

reduced SCD activity. However, unlike CLA, arachidonic acid (at 50 or 100 μ M) did not have any effect in this experiment.

4. Discussion

Previously, Lee et al. [23] reported that feeding CLA mixture significantly reduced hepatic SCD activity in mice by reducing SCD mRNA expression. In that study, it was suggested that since the *cis*-9,*trans*-11 isomer had no effect, *trans*-10,*cis*-12 CLA (the other main isomer) inhibited SCD. Our results clearly support that the *trans*-10,*cis*-12 isomer causes inhibition of SCD. While Lee et al. [23] measured mRNA expression from CLA fed animals, in this study we added CLA directly to the incubation medium with microsomal fractions for a relatively short time period (15 min). Therefore, inhibition by CLA in this study was not due to reduced SCD mRNA levels, indicating that *trans*-10,*cis*-12 CLA directly inhibits SCD enzymatic activity. These results are consistent with the recent publication by Bretillon et al. [30], which showed inhibition of SCD enzyme activity by *trans*-10,*cis*-12, but not by *cis*-9,*trans*-11 isomer.

As a structural requirement of inhibition of SCD, Clarke and Jump [31] proposed that an inhibitory fatty acid must contain two double bonds positioned between carbons 9–10 and 12–13 of an 18-carbon fatty acid. In addition, at least one of these double bonds must be *cis*, but the other can be in the *trans* configuration. Since *trans*-10,*cis*-12 CLA inhibits SCD, but *cis*-9,*trans*-11 does not, it is possible that the *cis*-12 position, or alternatively the *n*-6 position, may be the more important double bond location. This is supported by the intermediate effect of *cis*-12 octadecenoic acid, but not *trans*-10 octadecenoic acid. The minimal effect of *cis*-11 octadecenoic acid may be the result of its similarity to the *cis*-12 isomer.

It is apparent from this study, however, that *trans*-10,*cis*-12 CLA has additional inhibitory mechanisms that are distinct from polyunsaturated fatty acid (PUFA) which are known inhibitors of SCD. For example, linoleic, linolenic, arachidonic and eicosapentaenoic acids have been reported to decrease SCD activity in liver or adipocytes by decreased SCD

mRNA expression either by decreasing transcription or by decreasing mRNA stability [32]. Even though Clarke and Jump [31] reported PUFA effect on transcription response is rapid, 24 h feeding was the shortest time tested in mice [33]. In fact, arachidonic, linoleic, and *trans*-9,*cis*-12 octadecadienoic acids, all of which are either known inhibitors or would be expected to inhibit SCD since they fit the inhibitor requirements described earlier, did not show any effect in our experiments. Because of the relatively short incubation time in our studies (15 min), the interaction between *trans*-10,*cis*-12 CLA and the enzyme itself is more important than any effects on SCD mRNA expression.

Hovik et al. [34] reported direct inhibition of 9 or 10 thia fatty acids on SCD activity. Stearoyl-CoA with sulfur at the 9th or 10th position (9- or 10-thia stearoyl-CoA) apparently does not interfere with binding to the SCD, but the fatty acid cannot be desaturated, thus causing inhibition of Δ^9 desaturation. However, it has been suggested that 3-thia fatty acid reduced the affinity to the enzyme but permits desaturation. *trans*-10,*cis*-12 CLA may have a similar mechanism to thia fatty acid on SCD activity.

Fig. 4 indicates that the acid forms (alternatively CoA form) of the *trans*-10,*cis*-12 CLA isomer, and also 9-HODE and 9-HOO, were active while other forms (alcohol, amide or chloride) were not. This may indicate that there are additional mechanisms or requirements, such as CoA form of fatty acid, that are necessary to interact with the enzyme.

Increased SCD activity in liver and adipose tissue has been reported in relation to obesity [35–37]. In addition to the effects in liver, recently Choi et al. [38] discovered that *trans*-10,*cis*-12 CLA decreased SCD1 mRNA expression and activity in 3T3-L1 adipocytes. Along with previous reports of decreased lipoprotein lipase (LPL) activity in adipocytes and increased fatty acid β -oxidation in muscle, the effects of *trans*-10,*cis*-12 CLA on SCD may help explain CLA's multiple mechanisms on body fat reduction. We previously suggested that CLA acts by mediating prostaglandin mechanisms; however, Sessler et al. [39] indicated that decreased SCD mRNA expression by PUFA is eicosanoid independent. These reports along with the results in this paper suggest that CLA may exert its effects through multiple mechanisms.

When the compounds shown in Figs. 3 and 4 are

tested for LPL activity in 3T3-L1 adipocytes, only CLA-Alc was inhibitory (33%), and it was less effective than the *trans*-10,*cis*-12 CLA isomer (68%, unpublished observation). Interestingly, Cook et al. [40] recently reported that CLA-amide, CLA-Alc and 10y,*cis*-12 reduced fat pad weights in mice, while *cis*-12 octadecenoic acid did not. The discrepancy between these results may be due to different model systems. Animals may convert CLA-amide, CLA-Alc or 10y,*cis*-12 to the *trans*-10,*cis*-12 CLA isomer, but this conversion may not occur in an in vitro assay, particularly since incubation time is limited. These results may indicate the specificity of *trans*-10,*cis*-12 CLA on inhibition of SCD activity. Alternatively, SCD may be unrelated with body fat regulation since *cis*-12 octadecenoic acid had an inhibitory effect on SCD, but no effect on body fat reduction in mice [40].

Based on these results, it is apparent that *trans*-10 and *cis*-12 double bond positions and perhaps conjugation may be key structures for directly inhibiting SCD. These results, along with Lee et al.'s previous study [23] indicate that *trans*-10,*cis*-12 CLA acts on SCD by direct inhibition as well as by decreasing SCD mRNA expression.

In summary, *trans*-10,*cis*-12 CLA directly inhibited hepatic SCD, while *cis*-9,*trans*-11 or *trans*-9,*trans*-11 CLA isomers did not. *Cis*-12 and *cis*-11 octadecenoic acids and 10y,*cis*-12 octadecadienoic acid also inhibited SCD, but to a lesser extent than *trans*-10,*cis*-12 CLA. Linoleic acid derivatives 9-HODE and 9-HOO showed effects similar to *trans*-10,*cis*-12 CLA, while other 18-carbon fatty acids (*trans*-9, *trans*-10, *trans*-11, or *cis*-9 octadecenoic acids, *trans*-9,*cis*-12 octadecadienoic acid, or linoleic acid), CLA derivatives (alcohol, amide and chloride), and arachidonic acid did not have any inhibitory effects using a microsomal enzyme activity assay. A *cis*-12 double bond appears to be a key structure for inhibiting SCD activity, especially when coupled with a *trans*-10 double bond, but not with a 9 double bond.

Acknowledgements

We thank Professor E. Nordheim for assistance with statistical analysis and Dr. B. Fox for helpful discussion. This research was supported in part by

gift funds administered through the University of Wisconsin-Madison Food Research Institute. Three of the authors (Y.P., M.E.C., and M.W.P.) are inventors of CLA use patents that are assigned to the Wisconsin Alumni Research Foundation.

References

- [1] Y.L. Ha, N.K. Grimm, M.W. Pariza, Anticarcinogens from fried ground beef heat-altered derivatives of linoleic acid, *Carcinogenesis* 8 (1987) 1881–1887.
- [2] S.F. Chin, W. Liu, J.M. Storkson, Y.L. Ha, M.W. Pariza, Dietary sources of conjugated dienoic isomers of linoleic acid, a newly recognized class of anticarcinogens, *J. Food Comp. Anal.* 5 (1992) 185–197.
- [3] C.R. Kepler, K.P. Hirons, J.J. McNeill, S.B. Tove, Intermediates and products of the biohydrogenation of linoleic acid by *Butyrivibrio fibrisolvens*, *J. Biol. Chem.* 241 (1966) 1350–1354.
- [4] M.R. Pollard, Desaturation of positional and geometric isomers of monoenoic fatty acids by microsomal preparations from rat liver, *Lipids* 15 (1980) 306–314.
- [5] D.L. Palmquist, J.E. Santora, Endogenous synthesis of rumenic acid in rodents and humans, in: M.P. Yurawecz, M.M. Mossoba, J.K.G. Kramer, M.W. Pariza, G.J. Nelson (Eds.), *Advances in Conjugated Linoleic Acid Research*, Vol. 1, 1999, AOCS Press, Champaign, IL, pp. 201–208.
- [6] Y.L. Ha, J. Storkson, M.W. Pariza, Inhibition of benzo(a)pyrene-induced mouse forestomach neoplasia by conjugated dienoic derivatives of linoleic acid, *Cancer Res.* 50 (1990) 1097–1101.
- [7] C. Ip, S.F. Chin, J.A. Scimeca, M.W. Pariza, Mammary cancer prevention by conjugated dienoic derivative of linoleic acid, *Cancer Res.* 51 (1991) 6118–6124.
- [8] C. Liew, H.A.J. Schut, S.F. Chin, M.W. Pariza, R.H. Dashwood, Protection of conjugated linoleic acids against 2-amino-3-methylimidazo[4,5-f]quinoline-induced colon carcinogenesis in the F344 rat a study of inhibitory mechanisms, *Carcinogenesis* 16 (1995) 3037–3043.
- [9] K.N. Lee, D. Kritchevsky, M.W. Pariza, Conjugated linoleic acid and atherosclerosis in rabbits, *Atherosclerosis* 108 (1994) 19–25.
- [10] R.J. Nicolosi, K.V. Courtemanche, L. Laitinen, J.A. Scimeca, P.J. Huth, Effect of feeding diets enriched in conjugated linoleic acid on lipoproteins and aortic atherosclerosis in hamsters, *Circulation* 88 (Suppl.) (1993) 2458.
- [11] M.E. Cook, C.C. Miller, Y. Park, M.W. Pariza, Immune modulation by altered nutrient metabolism: nutritional control of immune-induced growth depression, *Poultry Sci.* 72 (1993) 1301–1305.
- [12] C.C. Miller, Y. Park, M.W. Pariza, M.E. Cook, Feeding conjugated linoleic acid to animals partially overcomes catabolic responses due to endotoxin injection, *Biochem. Biophys. Res. Commun.* 198 (1994) 1107–1112.
- [13] S.F. Chin, J. Storkson, K.J. Albright, M.E. Cook, M.W. Pariza, Conjugated linoleic acid is a growth factor for rats as shown by enhanced weight gain and improved feed efficiency, *J. Nutr.* 124 (1994) 2344–2349.
- [14] Y. Park, K.J. Albright, W. Liu, J.M. Storkson, M.E. Cook, M.W. Pariza, Effect of conjugated linoleic acid on body composition in mice, *Lipids* 32 (1997) 853–858.
- [15] Y. Park, K.J. Albright, J.M. Storkson, W. Liu, M.E. Cook, M.W. Pariza, Changes in body composition during feeding and withdrawal of dietary conjugated linoleic acid, *Lipids* 34 (1999) 243–248.
- [16] M.E. Cook, D.L. Jerome, T.D. Crenshaw, D.R. Buege, M.W. Pariza, K.J. Albright, S.P. Schmidt, J.A. Scimeca, P.A. Lofgen, E. Hentges, Feeding conjugated linoleic acid improves feed efficiency and reduces whole body fat in pigs, *FASEB J.* 12 (1998) A836.
- [17] Y. Park, Regulation of energy metabolism and the catabolic effects of immune stimulation by conjugated linoleic acid, Ph.D. Thesis, University of Wisconsin-Madison, 1996.
- [18] K.L. Houseknecht, J.P. Vanden Heuvel, S.Y. Moya-Camarena, C.P. Portocarrero, L.W. Peck, K.P. Nickel, M.A. Belury, Dietary conjugated linoleic acid normalizes impaired glucose tolerance in the Zucker diabetic fatty *fa/fa* rat, *Biochem. Biophys. Res. Commun.* 244 (1998) 678–682.
- [19] N. Sehat, M.P. Yurawecz, J.A.G. Roach, M.M. Mossoba, J.K.G. Kramer, Y. Ku, Silver-ion high-performance liquid chromatographic separation and identification of conjugated linoleic acid isomers, *Lipids* 33 (1998) 217–221.
- [20] W.W. Christie, G. Dobson, F.D. Gunstone, Isomers in commercial samples of conjugated linoleic acid, *J. Am. Oil Chem. Soc.* 74 (1997) 1231.
- [21] M. Sugano, A. Tsujita, M. Yamasaki, M. Noguchi, K. Yamada, Conjugated linoleic acid modulates tissue levels of chemical mediators and immunoglobulins in rat, *Lipids* 33 (1998) 521–527.
- [22] M.A. Belury, A. Kempa-Sterczko, Conjugated linoleic acid modulates hepatic lipid composition in mice, *Lipids* 32 (1997) 199–204.
- [23] K.N. Lee, M.W. Pariza, J.M. Ntambi, Conjugated linoleic acid decreases hepatic stearoyl-CoA desaturase mRNA expression, *Biochem. Biophys. Res. Commun.* 248 (1998) 817–821.
- [24] Y. Park, K.J. Albright, J.M. Storkson, W. Liu, M.W. Pariza, Evidence that the *trans*-10, *cis*-12 isomer of conjugated linoleic acid induces body composition changes in mice, *Lipids* 34 (1999) 235–241.
- [25] C.W. Miller, J.M. Ntambi, Peroxisome proliferators induce mouse liver stearoyl-CoA desaturase I gene expression, *Proc. Natl. Acad. Sci. USA* 93 (1996) 9443–9448.
- [26] H.E. May, P.B. McCay, Reduced triphosphopyridine nucleotide oxidase-catalyzed alterations of membrane phospholipids, *J. Biol. Chem.* 243 (1968) 2288–2295.
- [27] P. Legrand, A. Bensadoun, Stearoyl-CoA desaturase activity in cultured rat hepatocytes, *Biochim. Biophys. Acta* 1086 (1991) 89–94.
- [28] T.K.L. Dinh, J.M. Bourre, G. Durand, Effect of age and

- α -linolenic acid deficiency on Δ^6 desaturase activity and liver lipids in rats, *Lipids* 28 (1993) 517–523.
- [29] R.C. Littell, G.A. Milliken, W.W. Stroup, R.D. Wolfinger, SAS[®] System for Mixed Models, SAS Institute, Cary, NC, 1996.
- [30] L. Bretilon, J.M. Chardigny, S. Gregoire, O. Berdeaux, J.L. Sebedio, Effects of conjugated linoleic acid isomers on the hepatic microsomal desaturation activities in vitro, *Lipids* 34 (1999) 965–969.
- [31] S.D. Clarke, D.B. Jump, Polyunsaturated fatty acid regulation of hepatic gene transcription, *Lipids* 31 (1996) S7–S11.
- [32] J.M. Ntambi, Regulation of Stearoyl-CoA desaturase by polyunsaturated fatty acids and cholesterol, *J. Lipid Res.* 40 (1999) 1546–1558.
- [33] D.B. Jump, S.D. Clarke, A. Thelen, M. Lämatta, Coordinate regulation of glycolytic and lipogenic gene expression by polyunsaturated fatty acids, *J. Lipids Res.* 35 (1994) 1076–1084.
- [34] K.E. Hovik, O.S. Spydevold, J. Bremer, Thia fatty acids as substrates and inhibitors of stearoyl-CoA desaturase, *Biochim. Biophys. Acta* 1349 (1997) 251–256.
- [35] B.H. Jones, M.A. Maher, W.J. Banz, M.B. Zemel, J. Whelan, P.J. Smith, N. Moustaid, Adipose tissue stearoyl-CoA desaturase mRNA is increase by obesity and decreased by polyunsaturated fatty acids, *Am. J. Physiol.* 271 (1996) E44–E49.
- [36] D.A. Pan, A.J. Hulbert, L.H. Storlien, Dietary fats, membrane phospholipids and obesity, *J. Nutr.* 124 (1994) 1555–1565.
- [37] M. Enser, Desaturation of stearic acid by liver and adipose tissue from obese-hyperglycaemic mice (ob/ob), *Biochem. J.* 148 (1975) 551–555.
- [38] Y. Choi, Y.-C. Kim, Y.-B. Han, Y. Park, M.W. Pariza, J.M. Ntambi, The *trans*-10, *cis*-12 isomer of conjugated linoleic acid down regulate stearoyl-CoA desaturase gene expression in 3T3-L1 adipocytes, *J. Nutr.*, in press.
- [39] A.M. Sessler, N. Kaur, J.P. Palta, J.M. Ntambi, Regulation of stearoyl-CoA desaturase 1 mRNA stability by polyunsaturated fatty acids in 3T3-L1 adipocytes, *J. Biol. Chem.* 271 (1996) 29854–29858.
- [40] M. Cook, B.M. Drake, M.W. Pariza, Derivatives of conjugated linoleic acid (CLA) and fat reduction, *FASEB J.* 14 (2000) A525.

Editor-in-Chief

Howard R. Knapp
University of Iowa College of Medicine, Department of Internal Medicine/C31, GH, Iowa City, IA 52242-1081
Phone 1-319-353-7338 Fax 1-319-353-7340 E-mail howard-knapp@uiowa.edu

Associate Editors

Alan R. Brash
John G. Coniglio
William McLean Grogan
James A. Hamilton
Bruce J. Holub
Lloyd A. Horrocks
David Y. Hui
John M. Johnston
John K.G. Kramer

Arnis Kuksis
Robert A. Moreau
John J. Murray
Gary J. Nelson
Etsuo Niki
Ernst H. Oliw
Robert E. Pitas
Alfred Poulos

Colin Ratledge
Miriam D. Rosenthal
Norman Salem
Friedhelm Schroeder
Andrew J. Sinclair
Larry L. Swift
Maret G. Traber
Mike VanRollins

Editorial Advisory Board Members

Robert G. Ackman
A. Richard Baldwin
Wolfgang J. Baumann
Konrad E. Bloch
Bengt Borgström
Robert M. Burton
Kenneth K. Carroll
David J. Chitwood
Harold W. Gardner
Frank D. Gunstone
Mats Hamberg

Sven Hammarström
Raymond J. Hohl
Ralph T. Holman
Ching-hsien Huang
David Kritchevsky
Suzanne G. Laychock
Burton J. Litman
Helmut K. Mangold
David E. Ong
Rodolfo Paoletti
Forrest Quackenbush

Dwight R. Robinson
Angelo M. Scanu
Hannsjörg W. Seyberth
Leland L. Smith
Fred L. Snyder
Howard W. Sprecher
Paul K. Stumpf
Thomas E. Thompson
Moseley B. Waite
Randall D. Wood

Index Editor

Erika A. Kistler

AOCS Press Staff

Director, Mary C. Lane; Production Manager, Melissa Madsen Blankenship;
AOCS Press, P.O. Box 3489, Champaign, IL 61826-3489 USA;
Phone 1-217-359-2344; Fax 1-217-351-8091.

Printed on acid-free paper in the United States of America

Lipids (ISSN:0024-4201) is published monthly by AOCS Press at 1608 Broadmoor Dr., Champaign, IL 61821-5930. Annual subscription rate for AOCS U.S. members is \$60.00, for non-U.S. members \$75.00. Library/institution rates are \$215.00 for one year; individual nonmember subscriptions are \$95.00; student subscriptions are \$45.00. There is also an additional postage charge of \$15.00 per year for subscriptions from countries outside the United States. Optional air mail delivery is \$75.00 per year. Periodicals paid at Champaign, Illinois, and additional mailing offices. POSTMASTER: Send address changes to P.O. Box 3489, Champaign, IL 61826-3489 USA.

Editorial information: Current instructions to authors are printed in *Lipids*, Vol. 32, Nos. 1 and 7 (January and July 1997). Five copies of manuscripts intended for publication should be submitted to the Editor-in-Chief, Howard R. Knapp, at AOCS Press in Champaign, Illinois. Correspondence regarding manuscripts in the reviewing process should be directed to the Editor-in-Chief. Correspondence concerning page proofs should be addressed to the Production Manager, Melissa Madsen Blankenship, at AOCS Press. Page charges of \$55.00 per printed page are expected to be paid for papers

published in *Lipids*; payment of page charges will not affect review of papers. AOCS Press and the Editor-in-Chief of *Lipids* cannot assume responsibility for statements and opinions advanced by contributors.

Notice to copiers: Authorization to photocopy items for internal or personal use, or the internal or personal use of specific clients, is granted by AOCS Press for libraries and other users registered with the Copyright Clearance Center (CCC) Transactional Reporting Service, provided that the base fee of \$3.00 and a per page charge of \$0.50 per copy are paid directly to CCC, 27 Congress St., Salem, MA 01970.

Correspondence concerning new subscriptions, subscription renewals, changes of address, and claims for missing issues should be addressed to Circulation Department, *Lipids*, AOCS Press, P.O. Box 3489, Champaign, IL 61826-3489 USA. Requests for changes of address must include both old and new addresses, including zip codes, and a recent mailing label; notices must be received 30 days before the date of issue. Claims for missing issues must be received by the Circulation Department within 30 days (90 days for subscriptions from outside the United States).

Effect of Conjugated Linoleic Acid on Body Composition in Mice

Yeonhwa Park^a, Karen J. Albright^a, Wei Liu^a, Jayne M. Storkson^a,
Mark E. Cook^{a,b}, and Michael W. Pariza^{a,*}

Food Research Institute, ^aDepartment of Food Microbiology and Toxicology and ^bDepartment of Animal Sciences,
University of Wisconsin-Madison, Madison, Wisconsin 53706

ABSTRACT: The effects of conjugated linoleic acid (CLA) on body composition were investigated. ICR mice were fed a control diet containing 5.5% corn oil or a CLA-supplemented diet (5.0% corn oil plus 0.5% CLA). Mice fed CLA-supplemented diet exhibited 57% and 60% lower body fat and 5% and 14% increased lean body mass relative to controls ($P < 0.05$). Total carnitine palmitoyltransferase activity was increased by dietary CLA supplementation in both fat pad and skeletal muscle; the differences were significant for fat pad of fed mice and skeletal muscle of fasted mice. In cultured 3T3-L1 adipocytes CLA treatment (1×10^{-4} M) significantly reduced heparin-releasable lipoprotein lipase activity (−66%) and the intracellular concentrations of triacylglyceride (−8%) and glycerol (−15%), but significantly increased free glycerol in the culture medium (+22%) compared to control ($P < 0.05$). The effects of CLA on body composition appear to be due in part to reduced fat deposition and increased lipolysis in adipocytes, possibly coupled with enhanced fatty acid oxidation in both muscle cells and adipocytes. *Lipids* 32, 853–858 (1997).

CLA is the acronym for a group of positional and geometric isomers of conjugated dienoic derivatives of linoleic acid. They were identified as comprising an anticarcinogenic principal present in grilled ground beef and other food sources, especially dairy products (1–3). Synthetically prepared CLA inhibited carcinogen-induced neoplasia in several animal models (2,4,5) and reduced development of atherosclerosis in rabbits (6) and hamsters (7). CLA reduced tissue arachidonic acid levels and protected against the catabolic effects of endotoxin administration in mice, rats, and chickens without adversely affecting immune function (8,9). Recently it was established that CLA is a growth factor for young rats in that it enhanced body mass and feed efficiency (10). We now report the effects of CLA on body composition in mice.

*To whom correspondence should be addressed at Food Research Institute, 1925 Willow Drive, Madison, WI 53706.
E-mail: mwp@facstaff.wisc.edu.

Abbreviations: CLA, conjugated linoleic acid; CPT, carnitine palmitoyltransferase; DMEM, Dulbecco's modified Eagle's medium; LPL, lipoprotein lipase.

METHODS AND PROCEDURES

Materials. Linoleic acid was purchased from Nu-Chek-Prep Corporation (Elysian, MN); triolein, [9,10-³H(N)], (specific activity 12 Ci/mmol) from American Radiolabeled Chemicals Incorporated (St. Louis, MO); and [1-¹⁴C]linoleic acid (specific activity 55 mCi/mmol) from Amersham Life Science (Arlington Heights, IL). CLA was prepared as described (1).

Animals and body composition analyses. Weanling male ICR mice (experiment 1), 6-wk-old female (experiment 2) ICR mice, 6-wk-old male ICR mice [carnitine palmitoyltransferase (CPT) assay], feed ingredients (except for CLA), and semi-purified diet (TD94060, 99% basal mix) (Table 1) were purchased from Harlan Sprague-Dawley (Madison, WI). Diet was stored at −20°C until use. Mice were housed in a windowless room with a 12-h light-dark cycle in strict accordance to guidelines established by the Research Animal Resources Center of University of Wisconsin-Madison. Diet was fed *ad libitum*, freshly provided every day (experiment 1) or three times per week (experiment 2). After a 5-d adaptation period mice were randomly separated into groups and fed either control diet (5.5% corn oil) or CLA-containing diet

TABLE 1
Diet Composition^a

Ingredients	Treatment	
	Control (g/kg)	CLA (g/kg)
Sucrose	481	481
Casein, "Vitamin-Free" Test	210	210
Corn starch	150	150
DL-Methionine	3	3
Corn oil	55	50
CLA	0	5
Cellulose	50	50
Mineral mix, AIN-76	35	35
Vitamin mix, AIN-76A	10	10
Calcium carbonate	4	4
Choline bitartrate	2	2
Ethoxyquin	0.1	0.1

^aIn experiment 2, diet was prepared as described in Reference 37, which contained the same amount of ingredients except 495 g sucrose and 200 g casein per kg diet, without additional calcium carbonate. CLA, conjugated linoleic acid.

(5.0% corn oil plus 0.5% CLA). For body composition analyses, animals were sacrificed, gut contents were removed (to obtain empty carcass weight), and the carcasses frozen at -20°C . Frozen carcasses were chopped, ground, and freeze-dried to determine water content. Total nitrogen was analyzed by the Kjeldahl method (11). Carcass fat content was measured by extraction with diethyl ether overnight using a Soxhlet apparatus. Total ash content was determined by incineration (500 – 600°C , overnight).

Cell culture. 3T3-L1 preadipocytes were purchased from American Type Culture Collection (Rockville, MD) and cultured as described (12). Briefly, 3T3-L1 preadipocytes were grown to confluence at 37°C in Dulbecco's modified Eagle's medium (DMEM) containing 10% calf serum. At 2-d post-confluence (designated day 0) cell differentiation was induced with a mixture of methylisobutylxanthin (0.5 mM), dexamethasone ($0.25\text{ }\mu\text{M}$), and insulin ($1\text{ }\mu\text{g/mL}$) in DMEM containing 10% fetal bovine serum. On day 2 this medium was replaced with medium containing 10% fetal bovine serum and insulin only. On day 4 and thereafter the medium consisted of DMEM plus 10% fetal bovine serum only; this medium was subsequently replaced with fresh medium at 2-d intervals.

Fatty acid-albumin complexes were prepared as described (13) with slight modifications and added to culture media 48 h prior to cell harvest. CLA ($10\text{ }\mu\text{mol}$) or linoleic acid ($10\text{ }\mu\text{mol}$), dissolved in 0.1 M KOH, was added to $10\text{ }\mu\text{mol}$ bovine serum albumin solution in phosphate-buffered saline and incubated overnight at 4°C . The pH and volume were adjusted, respectively, to 7.2 and 5.0 mL (before filter sterilization) before use. Final concentrations of linoleic acid and CLA were $100\text{ }\mu\text{M}$ except as indicated for the dose-response experiment. All dishes including control had final concentration of $100\text{ }\mu\text{M}$ albumin.

Cell viability was tested using a number of parameters, as follows. The activities of citrate cleavage enzyme (14) and fatty acid synthetase (15) were determined. In addition 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (Sigma Chemical Co., St. Louis, MO) was used as described (16). Briefly, cells were cultured in a 96-well plate and treated with fatty acid-albumin complexes for 48 h as described above. After rinsing, cells were incubated with $50\text{ }\mu\text{L}$ of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (5 mg/mL) in DMEM at 37°C for 4 h. Medium was then removed and $200\text{ }\mu\text{L}$ of dimethyl sulfoxide added to each well to dissolve formazan crystals before quantifying at 570 nm with a Microplate Reader Model MR600 (Dynatech Industries, Inc., Alexandria, VA).

Free and esterified glycerol were determined using a Sigma Diagnostic Kit. Heparin-releasable lipoprotein lipase (LPL) activity (E.C. 3.1.1.34, 10 U heparin/mL media for 1 h at 37°C) was measured as described (17). Recovery of free fatty acid was estimated at 71% by using [^{14}C]linoleic acid. Protein was determined as described (18).

CPT (E.C. 2.3.1.23) assay. Mice were fed CLA-supplemented or control diet for 1 wk. Half the mice in each group were then fasted overnight prior to sacrifice by cervical dis-

cation. Epididymal fat pad, skeletal muscle (hind leg), and liver were removed. Tissues were then homogenized in 3 vol (wt/vol) of 0.25 M sucrose, 1 mM EDTA, 10 mM Tris HCl, pH 7.4 at 4°C . Homogenates were centrifuged ($700\times g$ for 10 min); supernatant fluid was again centrifuged ($12,000\times g$ for 15 min). Pellets (mitochondrial fraction) were washed and stored at -70°C after resuspension in 70 mM sucrose, 220 mM mannitol, 2 mM HEPES buffer, pH 7.4 and 1 mM EDTA. Total CPT activity was assayed by measuring the initial rates of CoASH formation (19).

Statistical analyses. Data in Figures 1 and 2 and Tables 2 and 3 were analyzed using the "2 independent sample *t*-test." Feed intake per cage was determined every day for experiment 1 (two mice per cage) or at 2–3-d intervals for experiment 2 (one mouse per cage). Feed intake data shown in Figures 1 and 2 are calculated under the assumption that in experiment 1 each mouse acted independently, and in experiment 2 mice consumed an equal amount of feed on each of the days constituting the feeding period. Two-way analysis of variance were performed on data (treatments and experiments) for Figure 3, Table 4, and the carcass protein analyses described in the Results section. Of major interest here are the comparisons among the treatments; these were computed using the Statistics Analysis System (SAS Users' Guide: Statistics, SAS Institute Inc., Cary, NC) with the General Linear Mean procedure and Least Square Means option. If the interaction between treatment and experiment was significant, this interaction was then used as the error term in the Least Square Means analysis.




FIG. 1. Body weight (circle) and feed intake (triangle) from male mice fed control or conjugated linoleic acid (CLA)-supplemented diet (experiment 1). Four-week-old male mice were fed either control (open) or CLA (filled)-supplemented diet (0.5% of diet) for 32 d. Body weights ($n = 8$) and feed intake were subjected to *t*-test ($*P < 0.05$). Numbers are mean \pm SE.




FIG. 2. Body weight (circle) and feed intake (triangle) from female mice fed control or CLA-supplemented diet (experiment 2). Seven-week-old female mice were fed either control (open) or CLA (filled) supplemented diet (0.5% of diet) for 28 d. Body weights and feed intake were subjected to *t*-test (* $P < 0.05$). Numbers are mean \pm SE ($n = 6$). See Figure 1 for abbreviation.

RESULTS

Figure 1 shows that male mice fed CLA-supplemented diet exhibited body weights that were indistinguishable from controls. By contrast, the data of Figure 2 indicate that female mice fed CLA appeared to exhibit slightly reduced body weight relative to controls, although the apparent reduction was not statistically significant. In both studies (Figs. 1 and 2) *ad libitum* feed intake for the CLA-fed animals appeared to be reduced at some experimental time points. We have previously shown that small differences in feed intake do not affect body composition (20).

Table 2 shows the effect of CLA supplementation on body composition in these animals. Relative to their respective controls, the percentage body fat in CLA-fed mice was reduced by 57% (males) and 60% (females). Hence, despite similar body weights (Figs. 1 and 2), dietary CLA supplementation resulted in significantly less body fat (Table 2).

By contrast, the percentages of whole body protein and

carcass water were significantly enhanced for CLA-fed mice in both experiments (Table 2). Because of the relatively small numbers of animals studied it is not possible to conclude from these data alone that CLA induced a significant increase in whole body protein accretion (whole body protein mass for controls and CLA-fed groups, respectively, were 5.7 and 6.0 g in experiment 1 and 4.4 and 4.6 g in experiment 2; in neither case was the difference significant). However, to date we have conducted 10 such mouse CLA feeding trials, only two of which are reported here in Figures 1 and 2 and Table 2. An analysis of the combined data from these 10 studies indicates that mice fed CLA-supplemented diets do in fact exhibit increased whole body protein relative to controls ($P = 0.0401$).

The mechanism of action of body composition changes is of considerable importance. Accordingly we studied the effect of CLA on total CPT, which is rate-limiting for fatty acid β -oxidation (Table 3). CPT activity was increased by dietary CLA supplementation in both fat pad and skeletal muscle; the differences were significant for fat pad of fed mice, and skeletal muscle of fasted mice. Liver CPT activity was not affected by CLA feeding.

Another key enzyme in lipid metabolism is adipocyte LPL, which hydrolyzes free fatty acids from circulating triacylglyceride; the fatty acids are then taken up by the adipocytes and re-esterified. We utilized 3T3-L1 adipocytes to measure LPL as well as fat mobilization. The amount of CLA added to the cell culture media was based on our determination that the mean CLA content of sera from rats fed diet supplemented with 0.5% CLA for 14 or 28 d is 72 μ M (range 23–120 μ M).

The data of Figure 3 show that supplementing the culture medium of fully differentiated 3T3-L1 adipocytes with CLA (but not linoleic acid) significantly reduced LPL activity. The inhibitory effect was linear between 20 and 200 μ M CLA (Fig. 4). The data indicate a threshold below 20 μ M where inhibition is not observed; the apparent enhancement at 5 μ M CLA is not understood and requires further study. In separate experiments to determine cell viability we found that incubating 3T3-L1 adipocytes for 2 d with 5 to 200 μ M CLA or linoleic acid did not change cell viability as measured using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (data not shown). We also found that incubating 3T3-L1 adipocytes for 2–3 d with CLA under similar conditions did not reduce citrate cleavage enzyme or fatty acid synthetase ac-

TABLE 2
CLA-Induced Changes in Body Composition^a

Experiment		ECW (g) ^b	Fat (%)	Protein (%)	Water (%)	Ash (%)
1	Control	32.4 \pm 1.1	10.13 \pm 1.17	17.76 \pm 0.30	66.3 \pm 0.8	3.08 \pm 0.14
	CLA	32.2 \pm 0.8	4.34 \pm 0.40 ^f	18.58 \pm 0.14 ^d	70.9 \pm 0.4 ^c	3.24 \pm 0.05 ^c
2	Control	25.0 \pm 0.9	18.68 \pm 3.08	17.67 \pm 0.61	58.3 \pm 2.2	3.67 \pm 0.15
	CLA	23.1 \pm 1.0	7.47 \pm 0.59 ^d	20.09 \pm 0.24 ^d	66.2 \pm 0.7 ^f	4.07 \pm 0.09

^aStudy duration was 32 d (experiment 1) and 28 d (experiment 2). Reported values are means \pm SE ($n = 6$ to 8).

^bEmpty carcass weight.

^{c–f}Indicates CLA-fed mice are different from control, where ^c $P < 0.05$, ^d $P < 0.01$, ^e $P < 0.001$, and ^f $P < 0.0001$. For other abbreviation, see Table 1.

TABLE 3
Total Carnitine Palmitoyltransferase Activity in Fat Pad, Muscle, and Liver^a

	Carnitine palmitoyltransferase (nmol ^b /min/mg protein)		
	Fat pad	Muscle	Liver
Fasted			
Control	15.9 ± 1.7	16.8 ± 2.5	16.7 ± 1.6
CLA	18.4 ± 1.9	25.9 ± 2.1 ^c	17.7 ± 1.3
Fed			
Control	14.7 ± 2.1	11.7 ± 1.2	5.3 ± 0.9
CLA	22.4 ± 1.4 ^c	14.3 ± 1.7	5.3 ± 0.6

^aMice were fed control or CLA-supplemented diet (0.5%) for 1 wk. Reported values are means ± SE (n = 6).

^bnmols L-carnitine exchange (Ref. 23).

^cP < 0.05 compared to control. For abbreviation, see Table 1.

tivities (data not shown). Finally, there were no CLA- or linoleic acid-induced differences in total protein or heparin-released protein from the cells. Hence, the CLA effect reported here appears to be specific and not due to toxicity.

Table 4 shows the amount of esterified and free glycerol within the cells, and the free glycerol in the culture medium. The findings are consistent with Figures 3 and 4 in that CLA treatment resulted in a significant reduction in intracellular triacylglyceride (determined as esterified glycerol). However, the significant increase in free glycerol in the media indicates that CLA may have stimulated lipolysis as well. Linoleic acid also appears to have stimulated lipolysis in these cells, which is consistent with the findings of Takada *et al.* (21) that body fat was modestly reduced in rats fed gamma-linolenic acid (a metabolite of linoleic acid).

DISCUSSION

The data of Table 2 establish that mice fed CLA-supplemented diets exhibited significant changes in body composition




FIG. 3. CLA inhibits lipoprotein lipase (LPL) activity in 3T3-L1 adipocytes. Reported values are means ± SE (n = 15–17, collected from four independent experiments). *P < 0.05. See Figure 1 for abbreviation.




FIG. 4. Effect of CLA dose on LPL activity in 3T3-L1 adipocytes. Reported values are means ± SE of 3–4 culture dishes normalized against control. The r^2 is 0.9901 for the data between 20 and 200 μM CLA. For abbreviations, see Figures 1 and 3.

tion relative to controls. This observation should be considered with regard to effects on the adipocyte (the principal site of fat storage) and the skeletal muscle cell (the principal site of fat oxidation).

The study described in Table 3 was conducted with tissues from control and CLA-fed mice. The findings are consistent with the interpretation that feeding mice a diet supplemented with CLA enhances fatty acid β-oxidation in skeletal muscle and fat pad, but not liver. These data (Table 3) may also explain at least in part our previous report (6) that rabbits fed CLA-supplemented diet exhibited reduced serum triglyceride levels.

The experiments described in Figures 3 and 4 and Table 4, which were conducted with 3T3-L1 adipocytes, indicate that CLA treatment reduced LPL activity while apparently enhancing lipolysis. If we assume that the *in vitro* experiments

TABLE 4
Esterified and Free Glycerol in Cell Sonicates, and Free Glycerol in Culture Medium from Control and CLA-Treated 3T3-L1 Adipocytes^a

	Percentage of control		
	Control	CLA	Linoleic acid
Esterified glycerol			
in cell	100 ^a	91.7 ± 6.4 ^b	117.1 ± 8.0 ^c
Free glycerol			
in cell	100 ^a	84.9 ± 7.2 ^b	107.4 ± 5.9 ^a
Free glycerol			
in culture medium	100 ^a	122.0 ± 12.9 ^b	106.6 ± 2.2 ^b

^aCells were treated with 100 μM of CLA or linoleic acid for 48 h before assay. The means are for 10 independent experiments, normalized against their respective controls. Data (2–6 dishes per treatment in each experiment) were analyzed as log value with a two-way analysis of variance using fatty acid (as treatments) and experiment. If the interaction between treatment and experiment was significant, this interaction was used as the error term. Means with different superscript letters are significantly different (P < 0.05). For abbreviation, see Table 1.

with 3T3-L1 adipocytes (Figs. 3, 4, Table 4) reflect *in vivo* physiological changes, then these data taken together with the data of Table 3 appear to provide a framework for partially understanding the reduction in body fat that is evident in the data of Table 2. It should be noted that the data of Figures 3 and 4 and Table 4 support our very recent findings (22) that adipocytes isolated from rats fed CLA exhibited increased hormone-sensitive lipase activity and enhanced norepinephrine-induced lipolysis.

A possible explanation for the apparent enhancement of lean body mass (Table 2 and accompanying discussion of whole body protein accretion in the Results section) is less clear. This may be related to our previous reports that CLA protects against the catabolic effects of immune stimulation (8,9), which is modulated by prostaglandin E_2 and mediated by interleukin-1 and tumor necrosis factor- α , both of which are linked to obesity (23–26). It is also possible that CLA affects pathways induced by anabolic hormones, perhaps in conjunction with an adaptive response by skeletal muscle necessitated by increased lipid substrate for β -oxidation. Further investigation is required.

There appear to be parallels between the effects of CLA reported and discussed herein and the reported effects of dietary fish oil. Like CLA, fish oil is reported to reduce fat pad size (27, but also see 28), prevent cachexia and body weight loss following immune stimulation (29,30), and reduce tissue arachidonic acid levels (31,32). The omega-3 fatty acids of fish oil modulate numerous biological and physiological effects that are linked to tumor necrosis factor- α , interleukin-1, and the eicosanoid pathways (30,33), indicating that the effects of CLA may also involve eicosanoid mechanism(s).

The isomers of CLA may be metabolized to one or more biologically active products which exert biological effect. For example CLA-derived eicosanoids could regulate the synthesis of arachidonic acid metabolites like prostaglandin E_2 . Alternatively, CLA-derived eicosanoids might exhibit biological activity in their own right. These possibilities are at this time speculation, but they are also the focus of considerable ongoing research (8,9,34–36). Understanding the biochemical mechanism(s) of action of CLA will be crucial to fully utilizing the potential of this class of fatty acids in animal husbandry and human application.

ACKNOWLEDGMENTS

We thank Drs. Richard Atkinson, Charles Elson, and Nancy Block for helpful critique in preparing this manuscript. We thank Professor E. Nordheim for assistance with statistical analysis. This research was supported in part by research grants from Kraft Foods Inc., the National Livestock and Meat Board, and gift funds administered through the University of Wisconsin–Madison Food Research Institute. Three of the authors (MWP, MEC, YP) are inventors of CLA-use patents assigned to the Wisconsin Alumni Research Foundation (WARF).

REFERENCES

- Chin, S.F., Liu, W., Storkson, J.M., Ha, Y.L., and Pariza, M.W. (1992) Dietary Sources of Conjugated Dienoic Isomers of Linoleic Acid, a Newly Recognized Class of Anticarcinogens, *J. Food Comp. Anal.* 5, 185–197.
- Ha, Y.L., Grimm, N.K., and Pariza, M.W. (1987) Anticarcinogens from Fried Ground Beef: Heat-Altered Derivatives of Linoleic Acid, *Carcinogenesis* 8, 1881–1887.
- Ha, Y.L., Grimm, N.K., and Pariza, M.W. (1989) Newly Recognized Anticarcinogenic Fatty Acids: Identification and Quantification in Natural and Processed Cheeses, *J. Agric. Food Chem.* 37, 75–81.
- Ha, Y.L., Storkson, J., and Pariza, M.W. (1990) Inhibition of Benzo(a)pyrene-Induced Mouse Forestomach Neoplasia by Conjugated Dienoic Derivatives of Linoleic Acid, *Cancer Res.* 50, 1097–1101.
- Ip, C., Chin, S.F., Scimeca, J.A., and Pariza, M.W. (1991) Mammary Cancer Prevention by Conjugated Dienoic Derivative of Linoleic Acid, *Cancer Res.* 51, 6118–6124.
- Lee, K.N., Kritchevsky, D., and Pariza, M.W. (1994) Conjugated Linoleic Acid and Atherosclerosis in Rabbits, *Atherosclerosis* 108, 19–25.
- Nicolosi, R.J., Courtemanche, K.V., Laitinen, L., Scimeca, J.A., and Huth, P.J. (1993) Effect of Feeding Diets Enriched in Conjugated Linoleic Acid on Lipoproteins and Aortic Atherogenesis in Hamsters, *Circulation* 88 (suppl.), 2458.
- Cook, M.E., Miller, C.C., Park, Y., and Pariza, M.W. (1993) Immune Modulation by Altered Nutrient Metabolism: Nutritional Control of Immune-Induced Growth Depression, *Poultry Sci.* 72, 1301–1305.
- Miller, C.C., Park, Y., Pariza, M.W., and Cook, M.E. (1994) Feeding Conjugated Linoleic Acid to Animals Partially Overcomes Catabolic Responses Due to Endotoxin Injection, *Biochem. Biophys. Res. Commun.* 198, 1107–1112.
- Chin, S.F., Storkson, J.M., Albright, K.J., Cook, M.E., and Pariza, M.W. (1994) Conjugated Linoleic Acid Is a Growth Factor for Rats as Shown by Enhanced Weight Gain and Improved Feed Efficiency, *J. Nutr.* 124, 2344–2349.
- Helrich, K. (1990) *Official Methods of Analysis*, 15th edn., pp. 935–937, Association of Official Analytical Chemists Inc., Arlington.
- Frost, S.C., and Lane, M.D. (1985) Evidence for the Involvement of Vicinal Sulfhydryl Groups in Insulin-Activated Hexose Transport by 3T3-L1 Adipocytes, *J. Biol. Chem.* 260, 2646–2652.
- Calder, P.C., Bond, J.A., Harvey, D.J., Gordon, S., and Newsholme, E.A. (1990) Uptake and Incorporation of Saturated and Unsaturated Fatty Acids into Macrophage Lipids and Their Effect upon Macrophage Adhesion and Phagocytosis, *Biochem. J.* 269, 807–814.
- Martyn, P., and Hansen, L.A. (1981) Initiation of Lipogenic Enzyme Activities in Rat Mammary Glands, *Biochem. J.* 198, 187–192.
- Lynen, F. (1962) Fatty Acid Synthesis from Malonyl CoA, *Methods Enzymol.* 1, 739–741.
- Plumb, J.A., Milroy, R., and Kaye, S.B. (1989) Effects of the pH Dependence of 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium Bromide-Formazan Absorption on Chemosensitivity Determined by a Novel Tetrazolium-based Assay, *Cancer Res.* 49, 4435–4440.
- Nilsson-Ehle, P., and Schotz, M.C. (1976) A Stable, Radioactive Substrate Emulsion for Assay of Lipoprotein Lipase, *J. Lipid Res.* 17, 536–541.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L., and Randall, R.J. (1951) Protein Measurement with the Folin Phenol Reagent, *J. Biol. Chem.* 193, 265–275.
- Bieber, L.L., Abraham, T., and Helmrath, T. (1972) A Rapid Spectrophotometric Assay for Carnitine Palmitoyltransferase, *Anal. Chem.* 50, 509–518.
- Boissonneault, G.A., Elson, C.E., and Pariza, M.W. (1986) Net

- Energy Effects of Dietary Fat on Chemically Induced Mammary Carcinogenesis in F344 Rats, *J. Natl. Cancer Inst.* 76, 335-338.
21. Takada, R., Saitoh, M., and Mori, T. (1994) Dietary γ -Linolenic Acid-Enriched Oil Reduces Body Fat Content and Induces Liver Enzyme Activities Relating to Fatty Acid β -Oxidation in Rats, *J. Nutr.* 124, 469-474.
 22. Pariza, M.W., Park, Y., Kim, S., Sugimoto, K., Albright, K., Liu, W., Storkson, J., and Cook, M. (1997) Mechanism of Body Fat Reduction by Conjugated Linoleic Acid, *FASEB J.* 11, A139.
 23. Hotamisligil, G.S., Arner, P., Caro, J.F., Atkinson, R.L., and Spiegelman, B.M. (1995) Increased Adipose Tissue Expression of TNF- α in Human Obesity and Insulin Resistance, *J. Clin. Invest.* 95, 2409-2415.
 24. Hotamisligil, G.S., Peraldi, P., Budavari, A., Ellis, R., White, M.F., and Spiegelman, B.M. (1996) IRS-1-Mediated Inhibition of Insulin Receptor Tyrosine Kinase Activity in TNF- α - and Obesity-Induced Insulin Resistance, *Science* 271, 665-668.
 25. Hotamisligil, G.S., Shargill, N.S., and Spiegelman, B.M. (1993) Adipose Expression of Tumor Necrosis Factor- α : Direct Role in Obesity-Linked Insulin Resistance, *Science* 259, 87-91.
 26. Bunout, D., Munoz, C., Lopez, M., de la Maza, M.P., Schlesinger, L., Hirsch, S., and Pettermann, M. (1996) Interleukin 1 and Tumor Necrosis Factor in Obese Alcoholics Compared with Normal-Weight Patients, *Am. J. Clin. Nutr.* 63, 373-376.
 27. Parrish, C.C., Pathy, D.A., and Angel, A. (1990) Dietary Fish Oils Limit Adipose Tissue Hypertrophy in Rats, *Metabolism* 39, 217-219.
 28. Awad, A.B., Bernardis, L.L., and Fink, C.S. (1990) Failure to Demonstrate an Effect of Dietary Fatty Acid Composition on Body Weight, Body Composition and Parameters of Lipid Metabolism in Mature Rats, *J. Nutr.* 120, 1277-1282.
 29. Beck, S., Smith, K.L., and Tisdale, M.J. (1991) Anticachectic and Antitumor Effect of Eicosapentaenoic Acid and Its Effect on Protein Turnover, *Cancer Res.* 51, 6089-6093.
 30. Hellerstein, M.K., Meydani, S.N., Meydani, M., Wu, K., and Dinarello, C.A. (1989) Interleukin-1-Induced Anorexia in the Rat, *J. Clin. Invest.* 84, 228-235.
 31. Leslie, C.A., Gonnerman, W.A., Ullman, M.D., Hayes, K.C., Franzblau, C., and Cathcart, E.S. (1985) Dietary Fish Oil Modulates Macrophage Fatty Acids and Decreases Arthritis Susceptibility in Mice, *J. Exp. Med.* 162, 1336-1349.
 32. Lokesh, B.R., Hsieh, H.L., and Kinsella, J.E. (1986) Peritoneal Macrophages from Mice Fed Dietary (n-3) Polyunsaturated Fatty Acids Secrete Low Levels of Prostaglandins, *J. Nutr.* 116, 2547-2552.
 33. Endres, S., Ghorbani, R., Kelley, V.E., Georgilis, K., Lonnemann, G., van der Meer, J.W.M., Cannon, J.G., Rogers, T.S., Klempner, M.S., Weber, P.C., Schaefer, E.J., Wolff, S.M., and Dinarello, C.A. (1989) The Effect of Dietary Supplementation with n-3 Polyunsaturated Fatty Acids on the Synthesis of Interleukin-1 and Tumor Necrosis Factor by Mononuclear Cells, *New Engl. J. Med.* 320, 265-271.
 34. Liew, C., Schut, H.A.J., Chin, S.F., Pariza, M.W., and Dashwood, R.H. (1995) Protection of Conjugated Linoleic Acids Against 2-Amino-3-methylimidazo[4,5-f]quinoline-Induced Colon Carcinogenesis in the F344 Rat: a Study of Inhibitory Mechanisms, *Carcinogenesis* 16, 3037-3043.
 35. Belury, M.A. (1995) Conjugated Dienoic Linoleate: A Polyunsaturated Fatty Acid with Unique Chemoprotective Properties, *Nutr. Rev.* 53, 83-89.
 36. Banni, S., Day, B.W., Evans, R.W., Corongin, F.P., and Lombardi, B. (1995) Detection of Conjugated Diene Isomers of Linoleic Acid in Liver Lipids of Rats Fed a Choline-Devoid Diet Indicates That the Diet Does Not Cause Lipid Peroxidation, *Nutr. Biochem.* 6, 281-289.
 37. Cohen, L.A., Choi, K., and Wang, C.-X. (1988) Influence of Dietary Fat, Caloric Restriction, and Voluntary Exercise on N-Nitrosomethylurea-Induced Mammary Tumorigenesis in Rats, *Cancer Res.* 48, 4276-4283.

[Received January 17, 1997, and in final revised form and accepted June 6, 1997]

Conjugated Linoleic Acid Is a Growth Factor for Rats as Shown by Enhanced Weight Gain and Improved Feed Efficiency^{1,2}

SOU F. CHIN, JAYNE M. STORKSON, KAREN J. ALBRIGHT, MARK E. COOK*
AND MICHAEL W. PARIZA³

Food Research Institute, Department of Food Microbiology and Toxicology and

*Department of Poultry Science, University of Wisconsin-Madison, Madison, WI 53706

ABSTRACT We studied the effect of conjugated linoleic acid (CLA) on rat development and growth. Primigravid female Fischer rats were fed control or CLA-supplemented (0.25% or 0.5% CLA) diets during gestation and/or lactation. Conjugated linoleic acid was incorporated into milk fat and tissue lipids proportional to the level of CLA fed and the duration of CLA feeding. Conjugated linoleic acid was incorporated into fetal and neonatal tissues; it did not affect litter size nor induce apparent abnormalities. To the contrary, feeding CLA to the dams during gestation and lactation improved the postnatal body weight gain of pups ($P < 0.05$), measured on d 10 of lactation. Pups that continued to receive the CLA-supplemented diet after weaning had significantly greater body weight gain and improved feed efficiency relative to control animals ($P < 0.05$). *J. Nutr.* 124: 2344-2349, 1994.

INDEXING KEY WORDS:

- conjugated linoleic acid • rats
- feed efficiency • growth enhancement

Conjugated linoleic acid (CLA) was first isolated from grilled ground beef and established as a potent cancer inhibitor with apparent antioxidant activity in several animal models (Ha et al. 1987 and 1990, Ip et al. 1991 and 1994). Conjugated linoleic acid was then shown to reduce the catabolic effects of immune stimulation in mice, rats and chickens without adversely affecting immune function (Cook et al. 1993, Miller et al. 1994). These findings indicate a complex mechanism of action for CLA that may involve effects on prostaglandin synthesis and signal transduction pathways. It should be noted that the catabolism of skeletal muscle, which is induced by immune stimulation, partitions energy away from other biological processes, including growth (Benson et al. 1993, Klasing and Austic 1984, Klasing et al. 1987).

Conjugated linoleic acid is present in virtually all foods (Chin et al. 1992), but the principal dietary sources are dairy products and other foods derived from ruminant animals (Bartlett and Chapman 1961, Chin et al. 1992). Interestingly, turkey lipid contains CLA at levels similar to those of ruminant animals (Chin et al. 1992).

Kepler et al. (1966) identified the *cis*-9, *trans*-11 CLA isomer as an intermediate in the biohydrogenation of linoleic acid by the rumen bacterium *Butyrivibrio fibrisolvens*. Seemingly some of the CLA produced in this way escapes conversion to stearic acid, is absorbed from the digestive system, and is subsequently incorporated into tissue lipid (including milk phospholipid). In this way CLA may be seen as an inherent component of cow's milk. Conjugated linoleic acid is also generated in the colon of conventional but not germ-free rats fed diets containing free linoleic acid (Chin et al. 1994). This finding indicates that microorganisms in the digestive tracts of monogastric animals are able to synthesize CLA.

It follows from these observations that CLA may be a previously unrecognized growth factor for at least some animal species. The purpose of this study was to investigate this possibility. We found that feeding CLA to rat dams during gestation and lactation enhanced pup growth while producing no observed evidence of harm. Consumption of the CLA-fortified diets by the pups after weaning further extended the

¹Supported in part by the College of Agricultural and Life Sciences, University of Wisconsin-Madison, and gift funds administered through the Food Research Institute, University of Wisconsin-Madison.

²The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

³To whom correspondence should be addressed.

body weight gain and improved feed efficiency as well.

MATERIALS AND METHODS

All protocols for animal use were approved by the Research Animal Resources Center of University of Wisconsin-Madison, and experimental procedures were conducted in accordance with the *Guide for the Care and Use of Laboratory Animals* (NRC 1985).

Animals. Two experiments were performed. In Experiment 1, 8-wk-old virgin female Fisher rats (Harlan Sprague Dawley, Indianapolis, IN) were used; in Experiment 2, 10-wk-old animals were used. Animals were maintained in individual cages at 20–22°C with 12-h light:dark photoperiods and fed a nonpurified diet (Ralston Purina, St. Louis, MO). They were transferred to separate cages for mating with males fed the same diet. Matings were confirmed by the presence of sperm in the vaginal smear. Day zero of gestation was defined as the first day of visible sperm in the vaginal smear.

Immediately after mating the females were removed and assigned to dietary groups. In Experiment 1, the animals were fed control diet or control diet fortified with 0.5 g CLA/100 g (Table 1). To ensure a homogenous mixture, CLA was mixed in corn oil before adding in basal diet. Diets were mixed weekly and stored under nitrogen at 4°C. Fresh diet was provided daily to each rat to minimize autoxidation. At d 20 of gestation, 10 rats each from the two treatment groups were killed by carbon dioxide suffocation; liver, mammary gland, skeletal muscle and abdominal adipose tissues were then removed. Fetuses were removed, weighed and examined visually for abnormalities. Maternal and fetal tissues were stored at –80°C until analyzed. The remaining rats (see footnotes to Tables 4 and 5) were allowed to deliver at term, and the day of parturition was designated as day zero of lactation.

The same diets were used in Experiment 2, with the addition of a third group fed control diet fortified with 0.25 g CLA/100 g. A fourth group received the control diet during gestation and the 0.5% CLA diet during lactation. Pups were weighed on d 10 and weaned on d 22 of lactation. Immediately after weaning, male and female pups were housed in individual cages and fed for 8 and 10 wk, respectively, the same diet their mothers had received. Body weights and food intakes were determined weekly.

Milk collection and analysis. Milk was collected on d 10 of lactation between 1200 and 1400 h, to minimize possible diurnal variations in milk composition [Picciano 1978]. Dams were separated from their litters 4 h before milking; longer periods of separation can affect milk composition [Keen et al. 1980]. Dams were anesthetized with sodium pentobarbital (Butler Company, Columbus, OH). Oxy-

TABLE 1

Diet composition¹

Component	Amount g/kg diet
Casein, vitamin-free test	216.5
DL-Methionine	3.1
Sucrose	495.83
Cornstarch	154.64
Corn oil	25.78
Cellulose	51.55
Mineral mix, AIN-76	36.09
Calcium carbonate	4.13
Vitamin mix, AIN-76A	10.31
Choline bitartrate	2.07

¹To facilitate the mixing of conjugated linoleic acid (CLA) into the diet, only 25.78 g corn oil/kg diet was included in the basal diet. Additional corn oil (25 g/kg diet) was used as a carrier for CLA. The final compositions of the diets were as follows (g/kg): 0.5% CLA, 50.78 g corn oil + 5 g CLA; 0.25% CLA, 53.28 g corn oil + 2.5 g CLA; control, 55.78 g corn oil.

tocin (1.2 USP units) (Fermentia Animal Health Company, Kansas City, MO) was injected intraperitoneally to stimulate milk flow. Milk was drawn by applying intermittent mild suction to each nipple through a Tygon plastic tube (Norton, Akron, OH) and collected in plastic tubes. Milk was not pooled; rather, milk from each dam was portioned into a number of plastic vials and stored frozen (–80°C).

Total protein in milk was determined by the BCA protein assay kit (Pierce Chemical, Rockford, IL). Total milk lipid was extracted as described by Supelco Inc. with slight modification due to limited sample size. To a 16-mL screw-capped tube the following was added: 0.5 mL of rat milk, 1 mL of ethanol, 0.3 mL of 16.44 mol/L ammonium hydroxide, 2.5 mL of petroleum ether, and 2.5 mL of diethyl ether. Samples were shaken for 5 min and allowed to stand for 20 min. The bottom phase was discarded; the ether layer was washed with water, filtered into a tared screw-capped test tube, and then dried under nitrogen. Net fat was weighed, capped under argon gas, and stored at –80°C.

Conjugated linoleic acid. Conjugated linoleic acid was synthesized as previously described [Chin et al. 1992]. Quantification of total CLA methyl ester in milk fat and animal tissues was performed by HPLC [Chin et al. 1992]. Gas chromatographic analyses of CLA isomers were conducted as described [Ha et al. 1989].

Nucleic acid analysis. Nucleic acids were extracted from the mammary gland [Ogur and Rosen 1950]. RNA was quantified [Ceriotti 1955] using baker's yeast RNA [Sigma Chemical, St. Louis, MO] as the standard. DNA was quantified [Hubbard et al. 1970]

using calf thymus DNA (Sigma Chemical) as the standard.

Statistical analyses. Data were analyzed by ANOVA using the general linear model procedure of the SAS statistical software (SAS Institute, Cary, NC). Significant differences among means were determined using the Tukey's comparisons (Steel and Torrie 1980). Litter size was tested for a potential covariate between pup weight and milk protein, but no significant effect was observed ($P > 0.12$). Therefore only one-way analysis was performed on pup body weight and milk protein. Differences were considered significant at $P < 0.05$.

RESULTS

Experiment 1. Consumption of CLA during gestation did not affect the food intake or body weights of the dams (Table 2). Mammary gland weights and DNA and RNA concentrations were comparable for CLA-fed and control animals, indicating that CLA did not directly affect mammary gland development. Liver weights for the two groups were also similar.

Litter sizes (0.5% CLA, 9.3; control, 9.2) and fetal body weights (0.5% CLA, 3.27 ± 0.05 g; control, 3.27 ± 0.28 g) at 20 d of gestation were unaffected by CLA consumption. Fetal liver (0.5% CLA, 0.22 ± 0.01 g; control 0.23 ± 0.02 g) and brain weights (0.5% CLA, 0.16 ± 0.01 g; control, 0.15 ± 0.01 g) were similar in the two groups. There was no gross visible evidence of fetal abnormality.

TABLE 2

Lack of effect of consumption of a diet containing 0.5 g conjugated linoleic acid/100 g diet (0.5% CLA) on mammary gland development in rats^{1,2}

	Dietary group	
	Control	0.5% CLA
Total food intake, g	251.1 \pm 3.4	247.7 \pm 6.33
Maternal body weight, g	226.3 \pm 4.14	221.9 \pm 2.84
Mammary gland weight, g	6.70 \pm 0.32	6.69 \pm 0.36
Net body weight of dams, g	196.1 \pm 4.42	192.0 \pm 2.19
Mammary gland weight/ 100 g net body wt, g	3.43 \pm 0.19	3.49 \pm 0.19
Liver weight, g	12.7 \pm 0.25	13.1 \pm 0.72
RNA ³	73.8 \pm 5.11	77.6 \pm 5.60
DNA ³	34.2 \pm 4.16	37.8 \pm 3.37
RNA:DNA ratio	2.45 \pm 0.34	2.16 \pm 0.18

¹Values are means \pm SE. There were 9 dams in the control group and 10 dams in the CLA-fed group.

²Body weight and mammary gland weight were measured on d 20 of gestation. Net body weight was calculated as maternal body weight minus fetal body weight.

³Milligrams per gram of dry fat-free mammary gland.

TABLE 3

Maternal and fetal tissue CLA concentrations in rats fed a diet containing 0.5 g conjugated linoleic acid/100 g diet (0.5% CLA)^{1,2}

	Dietary group	
	Control	0.5% CLA
	$\mu\text{mol CLA/g tissue lipid}$	
Maternal liver	0.53 \pm 0.18	12.0 \pm 1.03
Maternal muscle	0.78 \pm 0.11	15.9 \pm 1.36
Maternal mammary gland	1.21 \pm 0.11	29.7 \pm 1.43
Fetal liver	0.21 \pm 0.03	4.96 \pm 0.64

¹Values are means \pm SE. There were 9 dams in the control group and 10 dams in the CLA-fed group.

²Tissues were collected on d 20 of gestation. Livers from a litter were pooled together for fat extraction. Litter sizes were comparable (control group, 9.3 pups per dam; CLA-fed group, 9.2 pups per dam).

Tissue from control dams and fetuses contained low but detectable amounts of CLA (Table 3). In contrast, the CLA concentration in comparable tissue from animals fed the diet containing 0.5% CLA was substantially elevated. Conjugated linoleic acid was incorporated into milk fat of dams fed the CLA-supplemented diet (Table 4). Pups receiving CLA during gestation and lactation were significantly heavier than pups from control dams ($P < 0.036$).

Experiment 2. The CLA concentration of milk increased in direct proportion to dietary CLA level (Table 5). Pup development and survival were not affected by dietary treatment. However, for the group fed the diet containing 0.5% CLA, pup weight on d 10 of lactation was significantly higher than for controls or for the group fed the diet containing 0.25% CLA ($P < 0.03$). Pups from dams fed the 0.5% CLA-supplemented diet during lactation only were intermediate in weight.

TABLE 4

Effect of consumption of a diet containing 0.5 g conjugated linoleic acid/100 g diet (0.5% CLA) on rat pup weight, litter size and CLA concentration in milk^{1,2}

	Dietary group	
	Control	0.5% CLA
Mean pup weight, g	9.61 \pm 0.4	10.7 \pm 0.5*
Litter size	6.81	6.92
CLA in milk, $\mu\text{mol/g milk fat}$	1.68 \pm 0.21	46.7 \pm 3.0

¹Values are means \pm SE. There were 11 dams in the control group and 13 dams in the CLA-fed group. * $P < 0.036$ as determined by ANOVA.

²Pups were weighed on d 10 of lactation. Approximately 3 mL of milk was collected on d 10 of lactation.

TABLE 5

Effects of consumption of diets containing conjugated linoleic acid (CLA) during gestation and lactation, or during lactation only^{1,2}

	Dietary group ³			
	Control (n = 15)	0.25, 0.25 (n = 7)	0.5, 0.5 (n = 17)	0, 0.5 (n = 10)
CLA in milk, $\mu\text{mol/g fat}$	0.96 \pm 0.03	22.3 \pm 1.43	46.8 \pm 1.43	42.5 \pm 0.71
Litter size	10.4	9.9	10.3	9.7
Mean pup weight, g	12.8 \pm 0.4 ^a	12.8 \pm 0.6 ^a	14.0 \pm 0.3 ^b	13.5 \pm 0.3 ^{ab}

¹Values are means \pm SE. Numbers in parentheses indicate number of dams. Values for mean pup weight that do not share a common superscript are significantly different ($P < 0.05$, Tukey's test).

²Pups were weighed on d 10 of lactation for litters of nine to 11 pups. Approximately 3 mL of milk was collected on d 10 of lactation.

³The first and second numbers indicate the dietary percentages of CLA fed during gestation and lactation, respectively.

Figure 1 depicts body weights and food intakes, from weaning to 8 wk of age, for the male pups from the three treatment groups. Animals fed the 0.5% CLA-supplemented diet exhibited a small but consistent and statistically significant enhancement in body weight compared with control rats, whereas animals fed the 0.25% CLA-supplemented diet were intermediate in weight. Food intake over this period was not affected by treatment (Table 6). Feed efficiency (gram of body weight gain per gram of food

consumed over this time period) was of the order 0.5% CLA-fed group > 0.25% CLA-fed group > control group. The difference in feed efficiency values for the 0.5% CLA-fed group vs. the control group was significant ($P < 0.05$). Hence, supplementing the diet with 0.5% CLA resulted not only in greater body size but also in greater efficiency of feed utilization.

Female pups exhibited similar characteristics (Fig. 2, Table 7). Feed efficiencies for both the 0.5% and 0.25% CLA-fed groups were significantly higher than for the control group ($P < 0.05$).




FIGURE 1 Body weights of male rats fed control or treatment diets from 3 to 8 wk of age. Values are means \pm SE at each week for rats fed the control diet ($n = 43$ rats), the diet containing 0.25% conjugated linoleic acid (CLA) ($n = 34$ rats), and the diet containing 0.5% CLA ($n = 59$ rats). SE bars are within the size of symbol. Final body weights for both the 0.5% CLA-fed and the 0.25% CLA-fed groups are significantly higher ($P < 0.05$) than that of the control group (Tukey's test).




FIGURE 2 Body weights of female rats fed control or treatment diets from 3 to 10 wk of age. Values are means \pm SE at each week for rats fed the control diet ($n = 47$ rats), the diet containing 0.25% conjugated linoleic acid (CLA) ($n = 28$ rats), and the diet containing 0.5% CLA ($n = 60$ rats). SE bars are within the size of symbol. Final body weights for both the 0.5% CLA-fed and the 0.25% CLA-fed groups are significantly higher ($P < 0.05$) than that of the control group (Tukey's test).

TABLE 6

Effect of consumption of diets containing conjugated linoleic acid (CLA) on growth performance of male rats (5 wk after weaning)^{1,2}

	Dietary group		
	Control (n = 43)	0.25% CLA (n = 34)	0.5% CLA (n = 59)
Total body weight gain, g	126.8 ± 1.34 ^a	132.4 ± 1.79 ^b	134.9 ± 0.81 ^b
Total food intake, g	350.3 ± 2.86	359.5 ± 3.55	354.8 ± 2.44
Feed efficiency ³	0.362 ± 0.01 ^a	0.374 ± 0.01 ^{ab}	0.381 ± 0.01 ^b

¹Values are means ± SE, with the number of rats in parentheses. Within a row, values that do not share a common superscript are significantly different ($P < 0.05$, Tukey's test).

²Rats were weaned (on d 22 of lactation) from dams that had been fed control or CLA-supplemented diet during gestation and lactation. Immediately after weaning, rats were housed in individual cages and fed treatment diets until 8 wk old.

³Feed efficiency = gram of body weight gain per gram of food intake.

We also studied the effect of CLA consumption on the protein concentration of rat milk. In Experiment 1 there was more protein in milk from dams fed diet containing CLA (mice fed diet supplemented with 0.5% CLA, 114.1 ± 4.3 g protein/L; controls, 95.15 ± 3.0 g protein/L, $P < 0.006$). However, in Experiment 2 there were no differences among groups (mice fed diet supplemented with 0.5% CLA, 98.6 ± 1.7 g protein/L; mice fed diet supplemented with 0.25% CLA, 88.1 ± 4.5 g protein/L; controls, 96.5 ± 2.1 g protein/L). Hence, the effects reported here were not related to difference in milk protein concentration.

DISCUSSION

Conjugated linoleic acid is detectable in virtually every animal-derived food examined (Chin et al. 1992) and is consistently found in the tissues of rodents irrespective of diet (Table 4) (Chin et al. 1994, Ha et

al. 1990, Ip et al. 1991). Conjugated linoleic acid is elevated in meat derived from ruminant animals and seems to be an inherent component of cow's milk (Bartlett and Chapman 1961, Chin et al. 1992). In addition, elevated CLA levels occur in fresh and processed turkey meat (Chin et al. 1992) and in tissues of conventional but not germ-free rats fed diets containing free linoleic acid (Chin et al. 1994).

Conjugated linoleic acid displays a number of biological activities that may be regarded as beneficial. Conjugated linoleic acid inhibits chemically induced neoplasia at some sites in rats and mice (Ha et al. 1987 and 1990, Ip et al. 1991 and 1994) and seems to reduce blood LDL concentrations and inhibit the development of atherosclerosis in rabbits (Lee et al. 1994) and hamsters (Nicolosi et al. 1993) fed atherogenic diets. Conjugated linoleic acid reduces the catabolic response induced by immune stimulation in mice, rats and chickens without adversely affecting immune function (Cook et al. 1993, Miller et al.

TABLE 7

Effect of consumption of diets containing conjugated linoleic acid (CLA) on growth performance of female rats (7 wk after weaning)^{1,2}

	Dietary group		
	Control (n = 47)	0.25% CLA (n = 28)	0.5% CLA (n = 60)
Total body weight gain, g	113.6 ± 1.07 ^a	120.7 ± 0.78 ^b	122.8 ± 1.43 ^b
Total food intake, g	475.7 ± 4.83	473.4 ± 3.69	483.1 ± 5.91
Feed efficiency ³	0.239 ± 0.01 ^a	0.255 ± 0.01 ^b	0.254 ± 0.01 ^b

¹Values are means ± SE, with the number of rats in parentheses. Within a row, values that do not share a common superscript are significantly different ($P < 0.05$, Tukey's test).

²Rats were weaned (d 22 of lactation) from dams that had been fed control or CLA diet during gestation and lactation. Immediately after weaning, rats were housed in individual cages and fed treatment diets until 10 wk old.

³Feed efficiency = gram of body weight gain per gram of food intake.

1994). This response is cytokine-mediated and regulated by prostaglandin E_2 synthesis. It is possible that these seemingly disparate effects of CLA are mechanistically related through effects on prostaglandin E_2 metabolism and possibly also signal transduction pathways; the observation that CLA exhibits antioxidant activity in vitro and in vivo, especially in tissues (e.g., mammary gland) in which it is also an anticarcinogen [Ha et al. 1990, Ip et al. 1991] may also be mechanistically important.

Fish oil reduces the catabolic response induced by immune stimulation [Meydani 1992], but in mice and chickens CLA seems much more effective in this regard [Cook et al. 1993, Miller et al. 1994]. Bacteria, dust and related materials, which occur naturally in virtually all environments, serve as immune stimulants. The catabolism of skeletal muscle that follows immune stimulation partitions energy away from other biological processes. Reducing this response enhances growth and improves feed efficiency [Benson et al. 1993, Klasing and Austic 1984, Klasing et al. 1987].

Conjugated linoleic acid may be a previously unrecognized nutrient that functions to modulate the potentially devastating negative effects of immune stimulation. Growth promotion would be an expected reflection of such a physiological activity.

LITERATURE CITED

- Bartlett, J. C. & Chapman, D. G. (1961) Detection of hydrogenated fats in butter fat by measurement of cis-trans conjugated unsaturation. *J. Agric. Food Chem.* 9: 50-53.
- Benson, B. N., Calvert, C. C., Roura, E. & Klasing, K. C. (1993) Dietary energy source and density modulate the expression of immunologic stress in chicks. *J. Nutr.* 123: 1714-1723.
- Cerotti, G. (1955) The determination of nucleic acid in animal tissues. *J. Biol. Chem.* 214: 59-70.
- Chin, S. F., Liu, W., Storkson, J. M., Ha, Y. & Pariza, M. W. (1992) Dietary sources of conjugated dienoic isomers of linoleic acid, a newly recognized class of anticarcinogens. *J. Food Compos. Anal.* 5: 185-197.
- Chin, S. F., Storkson, J. M., Liu, W., Albright, K. J. & Pariza, M. W. (1994) Conjugated linoleic acid (9,11- and 10,12-octadecadienoic acid) is produced in conventional but not germ-free rats fed linoleic acid. *J. Nutr.* 124: 694-701.
- Cook, M. E., Miller, C. C., Park, Y. & Pariza, M. W. (1993) Immune modulation by altered nutrient metabolism: nutritional control of immune-induced growth depression. *Poult. Sci.* 72: 1301-1305.
- Ha, Y. L., Grimm, N. K. & Pariza, M. W. (1987) Anticarcinogens from fried ground beef: heat-altered derivatives of linoleic acid. *Carcinogenesis* 8: 1881-1887.
- Ha, Y. L., Grimm, N. K. & Pariza, M. W. (1989) Newly recognized anticarcinogenic fatty acids: identification and quantification in natural and processed cheese. *J. Agric. Food Chem.* 37: 75-81.
- Ha, Y. L., Storkson, J. M. & Pariza, M. W. (1990) Inhibition of benzo[a]pyrene-induced mouse forestomach neoplasia by conjugated dienoic derivatives of linoleic acid. *Cancer Res.* 50: 1097-1101.
- Hubbard, R. N., Mattaw, W. Y. & Dubonick, D. A. (1970) Factors influencing the determination of DNA with indole. *Anal. Biochem.* 38: 190-201.
- Ip, C., Chin, S. F., Scimeca, J. A. & Pariza, M. W. (1991) Mammary cancer prevention by conjugated dienoic derivative of linoleic acid. *Cancer Res.* 51: 6118-6124.
- Ip, C., Singh, M., Thompson, H. J. & Scimeca, J. A. (1994) Conjugated linoleic acid suppresses mammary carcinogenesis and proliferative activity of the mammary gland in the rat. *Cancer Res.* 54: 1212-1215.
- Keen, C. L., Lonnerdal, B., Sloan, M. V. & Hurley, L. S. (1980) Effects of milking procedure on rat milk composition. *Physiol. & Behav.* 24: 613-615.
- Kepler, C. R., Hiron, K. P., McNeill, J. J. & Tove, S. B. (1966) Intermediates and products of the biohydrogenation of linoleic acid by *Butyrivibrio fibrisolvens*. *J. Biol. Chem.* 241: 1350-1354.
- Klasing, K. C. & Austic, R. E. (1984) Changes in protein degradation in chickens due to an inflammatory challenge. *Proc. Soc. Exp. Biol. Med.* 176: 292-296.
- Klasing, K. C., Laurin, D. E., Peng, R. K. & Fry, D. M. (1987) Immunologically mediated growth depression in chicks: influence of feed intake, corticosterone and interleukin-1. *J. Nutr.* 117: 1629-1637.
- Lec, K., Kritchevsky, D. & Pariza, M. W. (1994) Conjugated linoleic acid and atherosclerosis in rabbits. *Atherosclerosis* 108: 19-25.
- Meydani, S. N. (1992) Modulation of cytokine production by dietary polyunsaturated fatty acids. *Proc. Soc. Exp. Biol. Med.* 200: 189-193.
- Miller, C. C., Park, Y., Pariza, M. W. & Cook, M. E. (1994) Feeding conjugated linoleic acid to animals partially overcomes catabolic response due to endotoxin injection. *Biochem. Biophys. Res. Commun.* 198: 1107-1112.
- Nicolosi, R. J., Courtemanch, K. V., Laitinen, L., Scimeca, J. A. & Huth, P. J. (1993) Effect of feeding diets enriched in conjugated linoleic acid on lipoproteins and aortic atherosclerosis in hamsters. *Circulation [Suppl.]* 88, no. 4, part 2, abs. 2458.
- National Research Council (1985) Guide for the Care and Use of Laboratory Animals. Publication no. 85-23 (rev.), National Institutes of Health, Bethesda, MD.
- Ogur, M. & Rosen, G. (1950) The nucleic acids of plant tissues. I. The extraction and estimation of deoxypentose nucleic acid and pentose nucleic acid. *Arch. Biochem.* 25: 262-276.
- Picciano, M. F. (1978) Mineral content of human milk during a single nursing. *Nutr. Rep. Int.* 18: 5-10.
- Steel, R.G.D. & Torrie, J. H. (1980) Principles and Procedures of Statistics. McGraw-Hill, New York, NY.

FEEDING CONJUGATED LINOLEIC ACID TO ANIMALS PARTIALLY
OVERCOMES CATABOLIC RESPONSES DUE TO ENDOTOXIN INJECTION

C.C. Miller, Y. Park¹, M.W. Pariza¹, and M. E. Cook²

Poultry Science Dept., Food Research Institute,
U.W. Madison, Madison, WI 53706

Received December 25, 1993

Summary: The ability of conjugated linoleic acid to prevent endotoxin-induced growth suppression was examined. Mice fed a basal diet or diet with 0.5% fish oil lost twice as much body weight after endotoxin injection than mice fed conjugated linoleic acid. By 72 hours post injection, mice fed conjugated linoleic acid had body weights similar to vehicle injected controls; however, body weights of basal and fish oil fed mice injected with endotoxin were reduced. Conjugated linoleic acid prevented anorexia from endotoxin injection. Splenocyte blastogenesis was increased by conjugated linoleic acid.

© 1994 Academic Press, Inc.

Recently, nonessential fatty acids have been shown to modulate acute inflammatory and immune responses possibly by changing cellular fatty acid profiles (1). The ability of two major fatty acid components of fish oil, eicosapentaenoate (EPA) and docosahexaenoate (DHA), to modulate the synthesis of arachidonate and its metabolites was recognized as one mechanism of immune regulation by fatty acids. EPA has been shown to depress arachidonic acid derived prostaglandin E₂ (PGE₂) synthesis in peritoneal leukocytes from rats (2). Unlike corn oil-fed rats, fish oil-fed rats did not exhibit the characteristic depression in food intake when injected with interleukin-1 (IL-1) (3). This phenomena has been partially attributed to the ability of fish oil to block or alter the cyclooxygenase pathway thereby decreasing PGE₂ synthesis and altering the ability of PGE₂ to negatively feed back on IL-1 synthesis.

CLA (conjugated dienoic derivative of linoleic acid) was shown to be a naturally occurring substance in food which has an anticarcinogenic effect on dimethylbenz(a)anthracene induced

²To whom correspondence should be addressed.

mammary tumors (4). Recent studies (5) have shown that CLA decreases arachidonic acid content of select tissues. It was postulated that like fish oil, CLA may exert an effect on the immune system by altering product formation of either the cyclooxygenase or lipoxygenase pathway. The purpose of the following investigation was to determine the effect of CLA on endotoxin induced growth inhibition and food intake depression. Fatty acid levels in tissue and *in vitro* lymphocyte proliferation was examined.

Materials and methods

Preparation of CLA and purity determination: CLA was prepared from linoleic acid by alkali isomerization as previously described (6, 7). Purity was determined to exceed 97%. CLA was stored in an argon atmosphere at -20°C.

Tissue fatty acid determination: Tissue fatty acid levels were determined by the following procedure. Total fat was extracted by chloroform:methanol (2:1, v/v) as described by Polch et al (8). Fatty acid methyl esters were prepared by reaction with 4% HCL in methanol for 20 minutes at 60 C (9) and extracted with hexane. Tridecanoic acid methyl ester was used as an internal standard and fatty acid methyl esters were identified by comparison with standards using gas chromatography (Hewlett Packard 5890 series II) (7).

Mouse experimental protocol: Three-week old mixed-sex mice (U.W. Madison Dairy Science Dept. outbred colony) (4 per group and 3 groups per dietary treatment) were fed a semi-purified basal diet (containing 2.5% linoleic acid) (Harlan Teklad, Madison, WI) or the basal diet containing either 0.5% added Menhaden fish oil with 25-30% omega-3 fatty acids as triglycerides (Sigma Chemical Co., St. Louis, MO) or 0.5% added CLA. All fatty acids were mixed into the basal diet following the protocol of Frische and Johnston (10) to minimize autooxidation. At the end of 15 consecutive feeding days, mice were weighed and i.p. injected with lipopolysaccharide (E. Coli 055:B5, Sigma Chemical Co., St. Louis, MO) at 1 mg/kg body weight in sterile HEPES buffer (Sigma Chemical Co., St. Louis, MO) or HEPES buffer alone. All mice were weighed 3, 8, 24, 48 and 72 hours post injection, and feed intake was determined at 3, 8 and 24 hours post injection.

At sacrifice, spleens were excised and single cell suspensions were made by teasing the spleen apart in calcium and magnesium free buffer (pH 7.0) and passing through an 18-gauge needle. Cells were then centrifuged through Histopaque 1077 (density 1.077) (Sigma Chemical Co., St. Louis, MO) and the buffy layer containing mononuclear cells was collected. Mononuclear cells were counted on a hemocytometer and tested for viability using trypan blue exclusion. Blastogenesis was measured following the procedure of Hughes et al (11). Results are expressed as a DPM index (disintegrations per minute in stimulated sample cells/ DPM in unstimulated sample cells).

Rat and chick experimental protocol: CLA (0.5% mixed into a semi-purified rat diet (Harlan Teklad, Madison, WI) or a standard chick mash) was fed to 20 rats (Harlan Sprague Dawley, Madison, WI) and 24 chicks (Leghorn X New Hampshire, Poultry Research Laboratory, Madison, WI) for 14 and 7 consecutive days, respectively. Control animals were fed their respective basal diets. After the feeding period, animals were injected i.p. with

endotoxin (see mouse protocol) and body weight losses due to the injection were monitored.

Statistics: Data were subjected to analysis of variance using the general linear models procedure of the Statistical Analysis System (12). Mean differences were determined using the least significant difference method.

Results

Feeding CLA to mice prevented the severe body weight loss caused by endotoxin injection as observed in mice fed the basal diet (figure 1). Fish oil (.5% of diet) was ineffective in preventing endotoxin-induced weight loss. Within the first 12 hours after endotoxin injection, body weight loss was equal among treatment groups. However, by 24 hours after injection, body weight losses in basal and fish oil fed mice were twice that of CLA fed mice. Return to pre-injection body weight also appeared to be faster in the groups fed CLA. In a similar manner, twenty-four hours post endotoxin injection, the reduction in body weight gain in CLA fed chicks (+1 g/24 hrs vs. +3 g/24 hrs for noninjected controls) and rats (-8 g/24 hrs vs +9 g/24 hrs for noninjected controls) was significantly less than endotoxin injected chicks (-6 g/24 hrs) and rats (-15 g/24 hrs) fed their respective basal diets.

Mice injected with buffer consumed significantly more feed at all time periods measured than mice fed either the basal diet or basal diet plus fish oil and injected with endotoxin (figure 2). The latter two groups consumed no feed during the 24 hour period after injection. However, mice fed CLA also consumed significantly more feed than mice fed fish oil or the basal diet respectively. Mice from all treatments which were injected with




Figure 1. Change in body weight due to endotoxin injection in mice consuming diets containing CLA or fish oil.

*Not different from CLA or fish oil fed and control inject.

abc Means with different letters are significantly different ($p < 0.05$).




Figure 2. Effect of endotoxin injection on cumulative feed intake in mice.

^{ab}Means with different letters are significantly different ($p < .05$).

buffer had feed consumptions and body weight gains which did not differ.

Mice fed CLA had a 1.5 fold increase in PHA-P induced spleen lymphocyte blastogenesis over basal fed mice (figure 3). CLA fed mice also displayed greater responses than fish oil fed mice, although this difference was not significant.

The muscle content of CLA was increased in rats (6.25 vs. .94 mg/g fat; $p < .05$) and chicks (4.07 vs. not detectable; $p < .05$) fed CLA relative to basal fed controls. Muscle arachidonic acid levels were decreased in CLA fed animals relative to the basal fed control animals (60.8 vs. 39.9 mg/g fat in rats and 43.6 vs. 38.8 mg/g fat in chicks).




Figure 3. Effect of feeding fish oil or CLA on PHA-P* induced spleen lymphocyte blastogenesis in mice.

*PHA-P level = .78 μ g/500,000 cells.

**Significantly greater than control ($p < .08$).

***DPM sample/DPM control.

Discussion

Klasing et al (13) showed that endotoxin induced weight loss was the result of not only reduced food intake, but also reduced fractional protein synthesis rate in the gastrocnemius and increased skeletal muscle degradation. Likewise, the ability of animals fed CLA to maintain body weight gain during endotoxin challenge may be due to factors other than maintenance of feed intake. Endotoxin injection has been shown to enhance IL-1 production, and semi-purified IL-1 was capable of decreasing rate of gain and feed intake to the same extent as endotoxin injection (13). Recently it has been demonstrated that feed intake depression associated with immune stimulation via IL-1 injection can be prevented by feeding 8% fish oil containing omega-3 fatty acids (3). Fish oils added in human diets increased cell membrane EPA (an n-3 fatty acid) and decreased cell membrane arachidonic acid (an n-6 fatty acid). The change in cell membrane fatty acids was thought to alter prostaglandin production and thus prevented body weight loss due to IL-1 injection (3). It was also thought that a depression of PGE₂ at the skeletal muscle level depressed protein degradation during IL-1 stimulation (14) since IL-1 was shown to enhance muscle catabolism by inducing PGE₂ production (15). It is possible that CLA acted through a similar mechanism and thereby altered cell membrane fatty acids and depressed PGE₂ production. Indeed CLA decreased skeletal muscle arachidonic acid levels. Thus animals maintained growth during immune stimulation.

Michel et al (16) recently reported that CLA stimulated mitogen-induced lymphocyte blastogenesis, cytotoxic activity and macrophage killing ability. In agreement with Michel and co-workers, we also have shown that CLA enhances PHA-P induced lymphocyte blastogenesis relative to basal fed controls.

Acknowledgment

Research was partially supported by the College of Agriculture and Life Sciences at the University of Wisconsin.

References

1. Lefkowitz, J.B., Morrison, A., Lee, V., and Rogers, M. (1990) *J. Immunology*. 145, 1523-1529.
2. Endres, S., Ghorbani, R., Kelley, V., Georgilis, K., Lonnemann, G., VanderMeer, J.W.M., Cannon, J.G., Rogers, T.S., Klempner, M.S., Weber, P.C., Schaefer, E.J., Wolff, S.M., Dinarello, C.A. (1989) *N. Engl. J. Med.* 320, 265-271.


3. Hellerstein, M.K., Meydani, S.N., Meydani, M., Wu, K., and Dinarello, C.A. (1989) *J. Clin. Invest.* 84, 228-235.
4. Ip, C., Chin, S.F., Scimeca, J.A., and Pariza, M. (1991) *Cancer Res.* 51, 6118-6124.
5. Cook, M.E., Miller, C.C., Park, Y., and Pariza, M. (1993) *Poultry Sci.* 72, 1301-1305.
6. Chin, S.F., Storkson, J., Ha, Y.L., and Pariza, M.W. (1992) *Food Comp. Anal.* 5, 185-197.
7. Ha, Y.L., Storkson, J. and Pariza, M. (1990) *Cancer Res.* 50, 1097-1101.
8. Folch, J., Lees, M., and Sloane-Stanley, G.H. (1957) *J. Biol. Chem.* 226, 497-509.
9. Carreau, J.P., and Dubaco, J.P. (1978) *J. Chromatograph.* 151, 384-390.
10. Frische, K.L., Johnston, P.V. (1988) *J. Nutr.* 118, 425-426.
11. Hughes, B.J., Hsich, G.C., Jarvis, B.B., and Sharma, R.P. (1989) *Arch. Environ. Contam. Toxicol.* 18, 388-395.
12. SAS Institute Inc. (1985) *SAS Users Guide: Statistics*, pp. 139-199. SAS Institute Inc., Cary, NC.
13. Klasing, K.C., Laurin, D.E., Peng, R.K., and Fry, D.M. (1987) *J. Nutr.* 117, 1629-1637.
14. Rodemann, H.P., and Goldberg, A.L. (1982) *J. Biol. Chem.* 257, 1632-1638.
15. Dinarello, C.A. (1984) *Infect. Dis.* 6, 51-95.
16. Michel, J.J., Chew, B.P., Schultz, T.D., Wong, T.S., and Magnuson, N.S. (1992) *Fed. Am. Soc. Exp. Biol. J.* 6, A1102.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: Jan Wadstein *et al.*
Serial No.: 09/410,484
Filed: 09/30/99
Entitled: **Method Of Treating Hypertension And Reducing Serum Lipase Activity**
Group No.: 1614
Examiner: Webman

Declaration of Asgeir Sæbo

Assistant Commissioner for Patents
Washington, D.C. 20231

CERTIFICATE OF MAILING UNDER 37 C.F.R. § 1.8(a)(1)(i)(A)	
<small>I hereby certify that this correspondence (along with any referred to as being attached or enclosed) is, on the date shown below, being deposited with the U.S. Postal Service with sufficient postage as first class mail in an envelope addressed to: Assistant Commissioner for Patents, Washington, D.C. 20231</small>	
Dated: <u>4-25-07</u>	By: 

I, Asgeir Sæbo, state as follows:

1. My present position is Research Director, Natural ASA.
2. It is my understanding that the Examiner has stated the following in the Office Action dated January 25, 2007:

First, Kawamura et al. provide a nexus teaching between weight loss in hypertensive patients and lowering of blood pressure. It remains obvious to one of ordinary skill in the art that the method of Cook et al. can lower blood pressure via weight loss. The Examiner did not assert that CLA would have a positive or negative on hypertension like ephedrine as submitted by Applicant. The Examiner asserts that weight loss is directly related to lowering of blood pressure, as supported by Kawamura et al., and CLA can be used to effect weight loss as taught by Cook et al. Secondly, one of ordinary skill in the art would have a reasonable expectation of success because of the nexus teaching of Kawamura et al.

3. When a biologically active agent, such as CLA, is administered to a subject there can be a variety of effects. Just because CLA causes weight loss does not also mean that it would reduce hypertension. A person of skill in the art would not reasonable expect CLA to reduce hypertension for two reasons.

4. First, CLA has been shown to elevate the level of F2-isoprostane. Taylor et al., *Conjugated Linoleic Acid Impairs Endothelial Function, Arteriosclerosis, Thrombosis, and Vascular Biology* 26(2), 307-312 (2006)(attached at Tab 1). F2-isoprostanes have a vasoconstrictive effect. Cracowski et al., *Cardiovascular pharmacology and physiology of the isoprostanes, Fundamental & Clinical Pharmacology* 20(5): 417-427 (2006)(attached at Tab 2). Taken together, it should be expected that administration of CLA would result in an increase in blood pressure.

5. Second, administration of other agents known to be effective for weight loss can result in increased hypertension. Ephedrine, a commonly used, biologically active weight loss supplement is one such example. As established in Haller and Benowitz, *Adverse Cardiovascular and Central Nervous System Events Associated with Dietary Supplements Containing Ephedra Alkaloids*, *New England J. Med.* 343(25):1833-1838 (2000)(attached at Tab 3), ephedrine can cause an increase in hypertension.

6. The Examiner's argument that it would be obvious to use CLA to decrease hypertension because CLA administration also causes weight loss lacks scientific merit. How an agent such as CLA acts in the body is complex. Whether CLA causes an increase or decrease in hypertension, or has no effect at all, is determined by a variety of factors that have no relation to weight loss. It is not scientifically valid to draw a conclusion that because an agent causes weight loss, it can also be expected to decrease hypertension. The references cited by the Examiner contain no data that can be interpreted in this manner.

7. I further declare that all statement made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of title 18 of the United States

PATENT

U.S. Appln. Ser. No.: 09/410,484
Attorney Docket No. NATNUT-03972

Code, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.



Asgeir Sæbo

Date: March 26th. 2007

MAIN Ser CIST1/ICIST NR CNRC
RM1 MAIN Ser
F981 0767-3981
v. 20 Received on: 06-10-25
no. 5 Fundamental & clinical
2006 Oct pharmacology.

ISSN 0767-3981 (print)
ISSN 1472-8206 (online)

Fundamental & Clinical **Pharmacology**

REVIEW
ARTICLECardiovascular pharmacology and
physiology of the isoprostanesJean-Luc Cracowski^{a*}, Thierry Durand^b^aLaboratoire de Pharmacologie, Inserm ESPRI, HP2 EA 3745, Faculté de Médecine de Grenoble, France^bUMR CNRS, 5074, Faculté de Pharmacie, Université Montpellier 1, Montpellier, France

Keywords

atherosclerosis,
hypertension,
ischemia,
isoprostane,
lipid peroxidation,
vasoconstriction

ABSTRACT

F_2 -isoprostanes are a complex family of compounds produced from arachidonic acid via a free radical-catalyzed mechanism. Their quantification as a pathophysiological biomarker provides a unique opportunity to investigate lipid peroxidation in vascular diseases. Their measurement also provides an interesting biomarker for the rational dose selection of antioxidants in vascular diseases where oxidative stress might be involved. In addition to their use as biomarkers, some isoprostanes possess a biological activity. The 15-series F_2 - and E_2 -isoprostanes mediate vasoconstriction in different vascular beds and species. In addition, 15- F_{2t} -IsoP induces smooth muscle cells mitogenesis and monocyte adhesion to endothelial cells. The data available supports but does not prove the hypothesis that isoprostanes are involved in vascular physiology and pathogenesis.

Received 2 March 2006;
revised 27 April 2006;
accepted 13 June 2006*Correspondence and reprints:
jean-luc.cracowski@ujf-
grenoble.fr

INTRODUCTION

Isoprostanes are a complex family of compounds produced from arachidonic acid via a free radical-catalyzed mechanism. In vitro generation of auto-oxidation products derived from polyunsaturated fatty acids was described more than 30 years ago [1,2]. However, the first demonstration that these compounds were produced in humans was shown in 1990 by Morrow et al. [3], who reported the discovery of prostaglandin- F_2 -like compounds, termed F_2 -isoprostanes, generated by free radical-induced peroxidation of arachidonic acid. Since that time, F_2 -isoprostanes have been used extensively as clinical markers of lipid peroxidation in cardiovascular disorders. These compounds are not only biomarkers. Indeed, the 15-series F_2 - and E_2 -isoprostanes possess a pharmacological activity on the blood vessels.

ISOPROSTANE STRUCTURE AND
SYNTHESIS

Depending on which of the labile hydrogen atoms of arachidonic acid is first abstracted by free radicals, three

initial arachidonoyl radicals can be formed following free radical attack. These radicals form four prostaglandin- H_2 -like compounds that can then be fully reduced to form four prostaglandin F_{2x} regioisomers (Figure 1), or rearranged to form prostaglandin E_2 and D_2 regioisomers. Two mechanisms, based on the formation of a 'dioxetane' intermediate, via a 4-*exo*-cyclization or a β -fragmentation followed by successive 5-*exo*-cyclizations have been proposed recently for the formation of these compounds [4]. Because each F_2 -isoprostane regioisomer comprises eight diastereoisomers, 64 different F_2 -isoprostanes can be generated.

Isoprostanes were formerly named according to the prostaglandin F_{2x} chemical structure. They differ from prostaglandins by the *cis*-stereochemistry of the five-membered ring junction instead of the *trans*-stereochemistry of the prostaglandin F_{2x} . Because the first isoprostanes described were the 15-series, they were formerly named according to this major difference, the first isoprostane being named 8-*iso*-prostaglandin F_{2x} or 8-*epi*-prostaglandin F_{2x} . However, such a nomenclature does not allow the differentiation of the numerous isomeric structures. Two nomenclatures were proposed




Figure 1 The isoprostane (IsoP) pathway. Free radical attack of arachidonic acid results in the formation of arachidonoyl radicals, which, following peroxidation, form four prostaglandin-H₂-like compounds that can then be fully reduced to form four prostaglandin F_{2s} regioisomers [those of the 15-series (type III), 8-series (type IV), 12-series (type V) and 5-series (type VI)], or rearranged to form prostaglandin E₂ and D₂ regioisomers. Each regioisomer comprises eight diastereoisomers and so 64 different F₂-isoprostanes can be generated.

recently, both of which enable an easy differentiation of the isoprostane isomers. Taber et al. [5] nomenclature was filed with the Eicosanoid Nomenclature Committee, and approved by the International Union of Pure and Applied Chemistry. Rokach et al. [6] also proposed a nomenclature that enables the differentiation of the regioisomers. In 1997, this nomenclature was modified to be applicable to isoprostane-like compounds derived from eicosapentaenoic and docosahexaenoic acid. The concomitant use of these three different nomenclatures is confusing for the nonspecialist, and we propose that the old nomenclature (e.g. 8-iso-prostaglandin F_{2α}) be definitively abandoned, and that Taber's nomenclature be encouraged. The different nomenclatures used to name the main isoprostanes are given in Figure 2.

Several *in vitro* studies have suggested a cyclooxygenase (COX)-dependent formation of 15-F_{2t}-IsoP [7–9]. An efficient *in vivo* production of 15-F_{2t}-IsoP through the COX pathway would reduce its accuracy as

a valid marker of lipid peroxidation. In contrast to the *in vitro* data, clinical studies clearly showed that COX inhibition was unable to decrease the formation of F₂-isoprostanes in healthy subjects as well as patients, suggesting that F₂-isoprostanes are formed via a non-COX-dependent mechanism *in vivo* [10–14]. Furthermore, in conditions of increased COX-2 expression following intravenous lipopolysaccharide challenge, the formation of 15-F_{2t}-IsoP and of 5-series isomers was not altered by COX inhibitors in healthy volunteers, whereas prostanoid production was decreased, further suggesting a COX-independent pathway of F₂-isoprostane synthesis [15]. Finally, an *in vivo* COX-dependent formation of IPF₂-III has been shown in the rat but not in humans [16]. Altogether, these data suggest that although a COX-dependent formation can be demonstrated *in vitro*, this does not occur *in vivo* in humans, meaning that 15-F_{2t}-IsoP as well as the 5-series isomers can be used as biomarkers of lipid peroxidation *in vivo*.

Chemical structure	Former nomenclature	Taber's nomenclature	Rokach's nomenclature
	8-iso-PGF _{2α}	15-F _{2t} -IsoP	IPF _{2α} -III
	8-iso-PGE ₂	15-E _{2t} -IsoP	1PE _{2t} -III
	2,3-dinor-5,6-dihydro-8-iso-PGF _{2α}	2,3-dinor-5,6-dihydro-15-F _{2t} -IsoP	2,3-dinor-5,6-dihydro-IPF _{2α} -III
	Not available	5-F _{2t} -IsoP	IPF _{2α} -VI
	Not available	5-F _{2t} -IsoP	8,12-iso-IPF _{2α} -VI
	8-iso-PGF _{1α}	15-F _{2t} -IsoP	IPF _{2α} -III

Figure 2 Nomenclature of the isoprostanes.

ISOPROSTANE QUANTIFICATION

Quantification of F₂-isoprostanes is used as a reliable marker of lipid peroxidation in vivo [17], and several methods are currently used [18] including gas chromatography (GC)-mass spectrometry (MS), which might be associated with an immunoaffinity extraction, GC-tandem MS, and liquid chromatography-tandem MS. These methods are reviewed in detail elsewhere [19]. They are specific but their cost and technology limit their routine use. Measurement of urinary 15-F_{2t}-IsoP by radioimmunoassay has been validated and constitutes a valid and easier alternative to GC-MS [20]. Enzyme immunoassays have also been developed to measure levels of F₂-isoprostanes but the antibodies used have not been tested for cross-reactivity with the numerous F₂-isoprostane isomers and their metabolites. The results obtained using enzyme immunoassays sometimes differ from those obtained using GC-MS assays and therefore immunoassays should be considered as semi-quantitative indices of F₂-isoprostanes.

VASCULAR PHARMACOLOGY OF THE ISOPROSTANES

Isoprostanes are formed in situ on phospholipids, at sites of free radical generation. Once released from cell

membranes by phospholipases, isoprostanes circulate in the plasma in free forms and are therefore susceptible to activate membrane receptors. Most studies have focused on the biological activity of 15-F_{2t}-IsoP, the first isoprostane commercially available. 15-F_{2t}-IsoP is a vasoconstrictor in most species and vascular beds, both in vitro and in vivo, following intravenous administration (see [21,22] for full reviews). These constrictor properties are not specific to the blood vessels and have been demonstrated in the lymphatic vessels, the bronchi, the gastrointestinal tract and the uterus. In addition, it stimulates mitogenesis in uterine vascular smooth muscle cells [23]. The available data strongly suggest that the effects of 15-F_{2t}-IsoP on blood vessels are mediated by the activation of the TP receptors (thromboxane A₂/prostaglandin H₂ receptors), acting as a full or partial agonist [24,25], although some responses including mitogenesis appear to be at least in part TP receptor independent. The existence of a specific isoprostane receptor has been suggested but remains to be elucidated [26]. In addition, preliminary data suggest that isoprostanes, as well as their precursor arachidonic acid, are other lipid ligands for the peroxisome proliferator activated receptors (PPAR) [27].

The effects of 15-F_{2t}-IsoP on platelets are complex. When incubated with subthreshold concentrations of ADP, thrombin, collagen and arachidonic acid, 15-F_{2t}-IsoP causes irreversible platelet aggregation, dependent




Figure 3 Schematic representation of the pharmacological activities of the isoprostane 15-F_{2t}-IsoP on the interface blood vessels. COX, cyclo-oxygenase; TXA₂, thromboxane A₂; PGI₂, prostacyclin; NO, nitric oxide; TP, prostaglandin H₂/thromboxane receptor; IPR, specific isoprostane receptor; PLC, phospholipase C; ET-1, endothelin-1; ECE, endothelin conversion enzyme. For a better comprehension, the thickness of the arrows are correlated to the scientific evidence of these mechanisms. Such mechanisms are likely to differ within species, as well as within the vessel types.

on thromboxane generation, while 15-F_{2t}-IsoP alone induces weak, reversible aggregation, only at high concentrations [28]. As 15-F_{2t}-IsoP is a partial agonist at the prostanoid TP receptor on platelets, it might inhibit the pro-aggregatory effects of TP receptor stimulation. Indeed, in human whole blood, 15-F_{2t}-IsoP is anti-aggregatory [29]. Several authors suggested that increased isoprostane formation is one of the factors involved in aspirin resistance, but a full demonstration of this hypothesis is not available to date [30–33].

15-F_{2t}-IsoP-induced contraction is modulated by the endothelium through the release of NO, i.e. endothelium removal increases 15-F_{2t}-IsoP contraction [21]. In addition, 15-F_{2t}-IsoP induces both thromboxane A₂ and endothelin-1 release from endothelial cells (Figure 3). In comparison with the huge data available for 15-F_{2t}-IsoP, few are available for other isomers. Nevertheless, other isoprostanes belonging to the 15 series of the F-family isoprostanes, such as 9-epi-15-F_{2t}-IsoP and 15-epi-15-F_{2t}-IsoP are biologically active, although less potent than 15-F_{2t}-IsoP [21]. The 5-series and 15-series F₂-isoprostanes are produced in approximately equal amounts in vivo whereas the 8-series and 12-series F₂-isoprostanes are produced in lower amounts [34]. In human urine and plasma, the 5-series (e.g. 5-F_{2t}-IsoP and 5-F_{2c}-IsoP) was found to be the most abundant F₂-isoprostanes [35]. Both the 15-series and the 5-series are easily detectable in human urine and plasma. However, in contrast to the

15-series F₂-isoprostanes, the 5-series F₂-isoprostanes have no vasomotor effect in different species and vascular beds [36] and as such are unlikely to be involved in the pathogenesis of vascular diseases. In addition to the vasoconstrictor and mitogenic effects, 15-F_{2t}-IsoP and 15-E_{2t}-IsoP induce monocyte adhesion to endothelial cells [37,38], whereas 15-F_{2t}-IsoP is a specific activator of rapid neutrophil adhesion [39]. Furthermore, 15-F_{2t}-IsoP induces cerebral endothelial cell death [40]. Endothelial cells are one of the first targets of oxidative stress in atherogenesis and ischemia–reperfusion injury. Whether isoprostanes may be one of pathogenic mediators remains to be tested.

Compared with the F₂-isoprostanes, E₂-isoprostanes are more potent in vitro. 15-E_{2t}-IsoP is more potent than 15-F_{2t}-IsoP in systemic and pulmonary vessels, its contraction being mediated through TP receptor, and BP₃ receptor activation in the pulmonary vasculature [41,42]. In addition, 15-E_{2t}-IsoP may induce a relaxation through EP receptors [43]. However, because no data are available concerning the production of E₂-isoprostanes in cardiovascular disorders, it is premature to conclude concerning the potential role of such compounds in cardiovascular pathogenesis.

One should keep in mind that Morrow et al. showed [44] that A₂-isothromboxanes are formed in vivo. Due to the inherent instability of the thromboxane A₂ ring, no data concerning the vascular effects of A₂-isothromb-

oxanes are available. However, regarding the potency of thromboxane A_2 -induced contractions, studies using stable A_2 -isothromboxanes analogues are awaited with interest.

To date, the metabolism of 15- F_{2t} -IsoP leads to two major metabolites in humans: 2,3-dinor-15- F_{2t} -IsoP and 2,3-dinor-5,6-dihydro-15- F_{2t} -IsoP [45,46]. The metabolite 2,3,4,5-tetranor-13,14-dihydro-15-keto-15- F_{2t} -IsoP [47] was identified as the major metabolite in rabbits but there is no evidence that such a metabolite is produced in humans. Although most prostanoid metabolites are biologically inactive, surprisingly, a recent report showed that 2,3-dinor-5,6-dihydro-15- F_{2t} -IsoP exhibits contraction that is comparable with that of 15- F_{2t} -IsoP in porcine brain microvessels [48]. By contrast, unlike 15- F_{2t} -IsoP, neither 2,3-dinor-5,6-dihydro-15- F_{2t} -IsoP nor 2,3-dinor-15- F_{2t} -IsoP had any constrictor or dilator effects on the rat thoracic aorta [49]. Such disparate observations need to be further investigated in different species and vascular beds.

Recently, isoprostane-like compounds derived from eicosapentaenoic and docosahexaenoic acids have been discovered *in vivo* [50–52]. Among the F_3 -isoprostanes formed, 15- F_3 -IsoP possesses either no biological effect or might induce a weak relaxation in human airways. Preliminary data suggest that F_4 -neuroprostanes possess no vascular effects.

ISOPROSTANES AS A BIOMARKER OF LIPID PEROXIDATION IN VASCULAR DISEASES

Isoprostanes have been measured in biological fluids such as urine, plasma, exhaled breath condensate, bronchoalveolar lavage fluid, bile, cerebrospinal, seminal and pericardial fluids. They are also detectable in normal tissues, including umbilical cords [53]. The main advantage of urinary measurements is that both 15- F_{2t} -IsoP and 5- F_{2t} -IsoP are not formed *ex vivo* by auto-oxidation in urine, unlike in plasma samples.

Cigarette smoking was one of the first conditions in which an increase in F_2 -isoprostane levels was demonstrated [54]. This increase is reduced after 2 weeks of abstinence from smoking [14,54] and almost reaches the values of nonsmokers 4 weeks after quitting smoking [55]. Short-term cigarette smoking increases exhaled breath condensate F_2 -isoprostane concentrations [56], but not plasma levels of F_2 -isoprostane [54]. Restarting smoking after quitting and passive smoking are associated with an increase in plasma levels of F_2 -isoprostane

[57,58]. Interestingly, 15- F_{2t} -IsoP concentrations were approximately twice as high in umbilical cords from newborn babies of smoking mothers compared with those of nonsmoking mothers [53]. Together, these data provide evidence that cigarette smoking is associated with a chronic increased lipid peroxidation *in vivo*.

The measurement of isoprostanes in biological fluids has prompted clinical investigations on the pathophysiological role of lipid peroxidation in cardiovascular diseases (Table 1). Among the biological fluids available, most studies were performed on urine because of the non-invasiveness of the procedure and the lack of artifactual generation. A strong link between lipid peroxidation and vascular diseases associated with ischemia-reperfusion, atherosclerosis and inflammation has been suggested by the elevated levels of lipid peroxidation observed in such diseases.

In addition to being a pathophysiological marker, the quantification of F_2 -isoprostanes might represent a prognostic marker. Indeed, Schwedhelm *et al.* [59] showed in a case-control study that urinary 15- F_{2t} -IsoP level was a strong independent concentration-dependent risk marker of coronary heart disease. In addition, there is a relationship between plasma F_2 -isoprostanes and early development of coronary artery calcifications [60]. There are currently no published clinical studies aimed at testing isoprostanes as a long-term prognostic marker, with strong endpoints such as mortality or morbidity, but cohort studies are on-going.

ISOPROSTANES: EMERGING ROLE IN VASCULAR PHYSIOLOGY AND DISEASE?

An important issue to resolve is whether the same effects observed *in vitro* are observed consistently *in vivo* at physiological concentrations and whether these effects contribute to pathological states *in vivo*. Basal plasma concentrations of 15- F_{2t} -IsoP have been found to range from approximately 10^{-10} to 5×10^{-10} mol L⁻¹ in plasma samples. These concentrations are unlikely to induce a systemic vasoactive effect considering the EC_{50} values of 15- F_{2t} -IsoP observed *in vitro* [21,61]. However, F_2 -isoprostanes are released at the site of free radical injury and then diluted in the circulation and therefore local concentrations might be sufficiently high to induce regional vasoconstriction. The concentrations of 15- F_{2t} -IsoP and 5- F_{2t} -IsoP are increased markedly in the coronary sinus following coronary angioplasty [62]. However, 15- F_{2t} -IsoP concentrations are in the

Table 1 P₂-isoprostane quantification in human cardiovascular diseases.

Disease	Isoprostanes quantified	Tissue or biological fluid tested	Method	Results in comparison with a control group	References
Essential hypertension	15-F _{2t} -isoP	Urine	EIA and GC-MS	NSD	[86-88]
Hypertensive patients with renovascular disease	15-F _{2t} -isoP	Urine	EIA	Increased	[86]
Heart failure	15-F _{2t} -isoP	Urine, pericardial fluid	LC-MS-MS and EIA	Increased	[89-92]
Atherosclerosis	15-F _{2t} -isoP and 5-F _{2t} -isoP	Atherosclerotic lesions from carotid endarterectomy	GC-MS and RIA	Increased	[93-95]
Ruptured abdominal aortic aneurysm	15-F _{2t} -isoP	Plasma	EIA	Increased	[96]
Stable coronary heart disease	15-F _{2t} -isoP	Urine	GC-MS and RIA	NSD	[97,98]
Unstable angina	15-F _{2t} -isoP	Urine	RIA	Increased	[98]
Reperfusion following myocardial infarction and cardiopulmonary bypass	15-F _{2t} -isoP and 5-F _{2t} -isoP	Urine	GC-MS	Increased	[97,99]
Coronary angioplasty	15-F _{2t} -isoP and 5-F _{2t} -isoP	Urine and coronary sinus	GC-MS	Increased	[62,99]
Systemic sclerosis (scleroderma)	15-F _{2t} -isoP and 5-F _{2t} -isoprostane metabolites	Urine	GC-MS AND EIA	Increased	[100-104]
Antiphospholipid antibodies syndrome	15-F _{2t} -isoP and 5-F _{2t} -isoP	Urine	GC-MS and EIA	Increased	[105,106]
Raynaud's phenomenon	15-F _{2t} -isoP	Urine	GC-MS	NSD	[101,104]
Pulmonary hypertension	15-F _{2t} -isoP	Urine	GC-MS	Increased	[107]
Acute ischemic stroke	15-F _{2t} -isoP	Urine	RIA	No variation over 72 h	[108]
Migraine	15-F _{2t} -isoP	Urine	RIA	NSD	[109]
Preeclampsia	15-F _{2t} -isoP and 5-F _{2t} -isoP	Plasma, urine and saliva and placental tissue	GC-MS and EIA	Conflicting results among studies	[110-116]

EIA, enzyme immunoassay; GC-MS, gas chromatography-mass spectrometry; LC-MS-MS, liquid chromatography-tandem mass spectrometry; NSD, not significantly different; RIA, radioimmunoassay.

nanomolar range, and thus unlikely to contribute to epicardial coronary artery vasoconstriction [61,63].

No specific inhibition of 15-F_{2t}-IsoP or other isoprostanes vascular effects can currently be achieved. However, TP receptor antagonists but not aspirin are effective in atherosclerosis inhibition in apo E knock-out mice, showing that TP receptors blockade by S18886 is effective by a mechanism independent of platelet-derived thromboxane A₂ [64], whereas isoprostanes suppression with vitamin E retards atherogenesis in the same animal model [65]. Similarly, TP receptor antagonism by L670596, but not COX-2 inhibition prevented pulmonary hypertension and endothelin-1 upregulation in 60% O₂-mediated pulmonary hypertension in newborns rats [66]. In addition to these animal data, a recent study showed that in patients suffering from coronary artery disease, S18886, a TP receptor antagonist improved acetylcholine-induced and flow-mediated vasodilation in patients treated with aspirin [67]. An hypothesis is that endogenous TP receptor activation induced by 15-F_{2t}-IsoP or other isoprostanes may be involved in the COX-independent effects of TP receptors antagonists [68]. However, because TP receptors share other endogenous ligands such as prostaglandin H₂ or hydroxyeicosatetraenoic acids (HETEs), such data give strength to the hypothesis that isoprostanes are involved in the vascular physiology and pathogenesis, but does not enable a definitive conclusion. Given that many isomeric isoprostanes exist, one should not focus only on 15-F_{2t}-IsoP. Other 15-series F₂-isoprostanes are biologically active, as well as E₂-isoprostanes [22] and large field of investigations are still unexplored.

ISOPROSTANES AS A PHARMACOLOGICAL TOOL FOR DRUG EVALUATION

In the past decade, most attention has focused on the effect of an antioxidant therapeutic strategy, including the use of vitamin E, in cardiovascular and nephrological diseases, with mixed results. Measurement of F₂-isoprostanes currently represents a valuable pharmacological tool for the evaluation of antioxidant therapy, and should be used in the rational selection of antioxidant dosages. Vitamin E supplementation induced a reduction of urinary 15-F_{2t}-IsoP levels in patients with cystic fibrosis [69], type 2 diabetes [11] and homozygous homocystinuria [70], but not in patients with systemic sclerosis [71]. In addition, such a reduction was dose dependent in hypercholesterolemic patients [10]. The

vitamin E-dependent reduction in the concentrations of F₂-isoprostane was observed in clinical conditions of enhanced oxidative stress. By contrast, supplementation with vitamin E had no effects on F₂-isoprostane levels in either moderate cigarette smokers [72], or healthy adults [73]. In the latter studies, the supplementation was associated with a significant dose-dependent increase in circulating concentrations of vitamin E. In addition, vitamin E supplementation in cigarette smokers on a high polyunsaturated fatty acid diet caused an increase in the plasma levels of F₂-isoprostane [74]. Together, these studies suggest that vitamin E supplementation has antioxidant effects in patient populations that are characterized by high rates of lipid peroxidation. Patrignani et al. [72] hypothesized that the basal rate of lipid peroxidation may be an important determinant of the response to vitamin E supplementation, and could explain the variable effects of vitamin E supplementation in large clinical trials. Several studies favor such a hypothesis. Oral treatment with raxofelast, a vitamin-E-like antioxidant, induced a significant reduction of plasma concentrations of 15-F_{2t}-IsoP in type 2 diabetes but had no effect in healthy subjects [75]. In addition, vitamin C supplementation reduced urinary 15-F_{2t}-IsoP levels in patients with stroke [76], and chronic alcoholic liver disease but not in patients with hepatitis C cirrhosis, in whom endogenous vitamin C and E concentrations did not significantly differ compared with controls [77]. Furthermore, vitamin C did not decrease urinary F₂-isoprostane and metabolite levels in healthy young women [78], and in young subjects with a limited history of cigarette smoking [79]. Together, these data strengthen the need to incorporate the measurement of surrogate end-points such as F₂-isoprostanes in large-scale antioxidant clinical trials.

In addition to drug evaluation, F₂-isoprostane measurement could be used to test the antioxidant properties of the diet. F₂-isoprostane levels were decreased following supplementation with eicosapentaenoic acid or docosahexaenoic acid [80], fish meals in diabetic patients [81], olive oil [82], soy-containing isoflavone [83], gazpacho [84], and flavanol-rich cocoa [85].

CONCLUSION

Isoprostanes are a complex family of compounds produced from arachidonic acid via a free radical-catalyzed mechanism. Some isoprostanes induce vasoconstriction, mitogenesis and monocyte adhesion. The quantification of F₂-isoprostanes as a pathophysiological biomarker

provides a unique opportunity to investigate lipid peroxidation in vascular diseases. Their measurement provides an interesting biomarker for the rational dose selection of antioxidants in vascular diseases where oxidative stress might be involved. The data available supports but does not prove the hypothesis that isoprostanes are involved in vascular physiology and pathogenesis.

REFERENCES

- Nugteren D.H., Voncken H., Van Drop D.A. Non-enzymatic conversion of all-cis 8,11,14-eicosatrienoic acid into prostaglandin H₂. *Rec. Trav. Chim.* (1967) 86 1237–1245.
- Pryor W.A., Stanley J.P., Blair R. Autooxidation of polyunsaturated fatty acids: II. A suggested mechanism for the formation of TBA-reactive materials from prostaglandin-like endoperoxides. *Lipids* (1976) 11 370–379.
- Morrow J.D., Hill K.E., Burk R.F., Nammour T.M., Badr K.F., Roberts L.J. A series of prostaglandin F₂-like compounds are produced in vivo in humans by a non-cyclooxygenase, free radical-catalyzed mechanism. *Proc. Natl. Acad. Sci. U. S. A.* (1990) 87 9383–9387.
- Yin H., Havrilla C.M., Gao L., Morrow J.D., Porter N.A. Mechanisms for the formation of isoprostane endoperoxides from arachidonic acid. 'Dioxetane' intermediate versus beta-fragmentation of peroxy radicals. *J. Biol. Chem.* (2003) 278 16720–16725.
- Taber D.F., Morrow J.D., Robert L.J. II A nomenclature system for the isoprostanes. *Prostaglandins* (1997) 53 63–67.
- Rokach J., Khanapure S.P., Hwang S.W., Adiyaman M., Lawson J.A., Fitzgerald G.A. Nomenclature of isoprostanes: a proposal. *Prostaglandins* (1997) 54 853–873.
- Patriguani P., Santini G., Panara M.R. et al. Induction of prostaglandin endoperoxide synthase-2 in human monocytes associated with cyclo-oxygenase-dependent F₂-isoprostane formation. *Br. J. Pharmacol.* (1996) 118 1285–1293.
- Pratico D., Lawson J.A., Fitzgerald G.A. Cyclooxygenase-dependent formation of the isoprostane, 8-epi-prostaglandin F_{2α}. *J. Biol. Chem.* (1995) 270 9800–9808.
- Jourdan K., Mitchell J., Evans T. Release of isoprostanes by human pulmonary artery in organ culture: a cyclo-oxygenase and nitric oxide dependent pathway. *Biochem. Biophys. Res. Commun.* (1997) 233 668–672.
- Davi G., Alessandrini P., Mezzetti A. et al. In vivo formation of 8-epi-prostaglandin F_{2α} is increased in hypercholesterolemia. *Arterioscler. Thromb. Vasc. Biol.* (1997) 17 3230–3235.
- Davi G., Ciabattini G., Consoli A. et al. In vivo formation of 8-iso-prostaglandin F_{2α} and platelet activation in diabetes mellitus: effects of improved metabolic control and vitamin E supplementation. *Circulation* (1999) 99 224–229.
- Dworzki R., Murray J., Roberts L.J. et al. Allergen-induced synthesis of F₂-isoprostanes in atopic asthmatics. *Am. J. Respir. Crit. Care Med.* (1999) 160 1947–1951.
- Gopaul N.K., Anggard E.E., Mallet A.I., Betteridge D.J., Wolff S.P., Nourroz-Zadeh J. Plasma 8-epi-PGF_{2α} levels are elevated in individuals with non-insulin dependent diabetes mellitus. *FEBS Lett.* (1995) 368 225–229.
- Reilly M., Delanty N., Lawson J.A., Fitzgerald G.A. Modulation of oxidant stress in vivo in chronic cigarette smokers. *Circulation* (1996) 94 19–25.
- McAdam B.F., Mardini I.A., Habib A. et al. Effect of regulated expression of human cyclooxygenase isoforms on eicosanoid and isoeicosanoid production in inflammation. *J. Clin. Invest.* (2000) 105 1473–1482.
- Bachi A., Brambilla R., Fanelli R., Bianchi R., Zuccato E., Chiabrando C. Reduction of urinary 8-epi-prostaglandin F_{2α} during cyclo-oxygenase inhibition in rats but not in man. *Br. J. Pharmacol.* (1997) 121 1770–1774.
- Roberts L.J., Morrow J.D. Measurement of F₂-isoprostanes as an index of oxidative stress in vivo. *Free Radic. Biol. Med.* (2000) 28 505–513.
- Tsikas D. Application of gas chromatography-mass spectrometry and gas chromatography-tandem mass spectrometry to assess in vivo synthesis of prostaglandins, thromboxane, leukotrienes, isoprostanes and related compounds in humans. *J. Chromatogr. B Biomed. Sci. Appl.* (1998) 717 201–245.
- Berdeaux O., Scruel O., Cracowski J.L., Durand T. F₂-isoprostanes: review of analytical methods. *Curr. Pharm. Anal.* (2006) 2 69–78.
- Wang Z., Ciabattini G., Crémignon C. et al. Immunological characterization of urinary 8-epi-prostaglandin F_{2α} excretion in man. *J. Pharmacol. Exp. Ther.* (1995) 275 94–100.
- Cracowski J.L., Devillier P., Durand T., Stanke-Labesque F., Bessard C. Vascular biology of the isoprostanes. *J. Vasc. Res.* (2001) 38 93–103.
- Janssen L.J. Isoprostanes: an overview and putative roles in pulmonary pathophysiology. *Am. J. Physiol. Lung Cell. Mol. Physiol.* (2001) 280 L1067–L1082.
- Miggin S.M., Kinsella B.T. Thromboxane A₂ receptor mediated activation of the mitogen activated protein kinase cascades in human uterine smooth muscle cells. *Biochim. Biophys. Acta* (2001) 1539 147–162.
- Kinsella B.T. Thromboxane A₂ signalling in humans: a 'Tail' of two receptors. *Biochem. Soc. Trans.* (2001) 29 641–654.
- Daray F.M., Minvielle A.L., Pappo S., Rothlin R.P. Vasoconstrictor effects of 8-iso-prostaglandin E₂ and 8-iso-prostaglandin F_{2α} on human umbilical vein. *Eur. J. Pharmacol.* (2004) 499 189–195.
- Fukunaga M., Yura T., Grygorczyk R., Badr K.F. Evidence for the distinct nature of F₂-isoprostane receptors from those of thromboxane A₂. *Am. J. Physiol.* (1997) 272 F477–F483.
- McNamara P., Lawson J.A., Rokach J., Fitzgerald G.A. Isoprostane activation of the nuclear hormone receptor PPAR. *Adv. Exp. Med. Biol.* (2002) 507 351–355.
- Pratico D., Smyth B.M., Violi F., Fitzgerald G.A. Local amplification of platelet function by 8-epi prostaglandin F_{2α} is not mediated by thromboxane receptor isoforms. *J. Biol. Chem.* (1996) 271 14916–14924.
- Cranshaw J.B., Evans T.W., Mitchell J.A. Characterization of the effects of isoprostanes on platelet aggregation in human whole blood. *Br. J. Pharmacol.* (2001) 132 1699–1706.

- 30 Cambria-Kiely J.A., Gandhi P.J. Possible mechanisms of aspirin resistance. *J. Thromb. Thrombolysis* (2002) 13 49–56.
- 31 Maree A.O., Fitzgerald D.J. Aspirin and coronary artery disease. *Thromb. Haemost.* (2004) 92 1175–1181.
- 32 Martin C.P., Talbert R.L. Aspirin resistance: an evaluation of current evidence and measurement methods. *Pharmacotherapy* (2005) 25 942–953.
- 33 Patrignani P. Aspirin insensitive eicosanoid biosynthesis in cardiovascular disease. *Thromb. Res.* (2003) 110 281–286.
- 34 Waugh R.J., Morrow J.D., Roberts L.J., II, Murphy R.C. Identification and relative quantitation of F₂-isoprostane regioisomers formed in vivo in the rat. *Free Radic. Biol. Med.* (1997) 23 943–954.
- 35 Lawson J.A., Li H., Kokach J. et al. Identification of two major F₂ isoprostanes, 8,12-iso- and 5-epi-8, 12-iso-isoprostane F_{2α}-VI, in human urine. *J. Biol. Chem.* (1998) 273 29295–29301.
- 36 Marliere S., Cracowski J.L., Durand T. et al. The 5-series F₂-isoprostanes possess no vasomotor effects in the rat thoracic aorta, the human internal mammary artery and the human saphenous vein. *Br. J. Pharmacol.* (2002) 135 1276–1280.
- 37 Huber J., Bochkov V.N., Binder B.R., Leitinger N. The isoprostane 8-iso-PGF_{2α} stimulates endothelial cells to bind monocytes via cyclic AMP- and p38 MAP kinase-dependent signaling pathways. *Antioxid. Redox Signal.* (2003) 5 163–169.
- 38 Leitinger N., Huber J., Rizzo C. et al. The isoprostane 8-iso-PGF_{2α} stimulates endothelial cells to bind monocytes: differences from thromboxane-mediated endothelial activation. *FASEB J.* (2001) 15 1254–1256.
- 39 Fontana L., Giagulli C., Minn P., Lechi A., Laudanna C. 8-iso-PGF_{2α} induces beta 2-integrin-mediated rapid adhesion of human polymorphonuclear neutrophils: a link between oxidative stress and ischemia/reperfusion injury. *Arterioscler. Thromb. Vasc. Biol.* (2001) 21 55–60.
- 40 Brault S., Martinez-Bermudez A.K., Marrache A.M. et al. Selective neuromicrovascular endothelial cell death by 8-iso-prostaglandin F_{2α}: possible role in ischemic brain injury. *Stroke* (2003) 34 776–782.
- 41 Janssen L.J., Tazao T. Involvement of TP and EP3 receptors in vasoconstrictor responses to isoprostanes in pulmonary vasculature. *J. Pharmacol. Exp. Ther.* (2002) 301 1060–1066.
- 42 Cracowski J.L., Devillier P., Chavanon O., Sitchaping-Naspa F.A., Stanke-Labesque F., Bessard G. Isoprostaglandin E₂ type III (8-iso-prostaglandin E₂) evoked contractions in human internal mammary artery. *Life Sci.* (2001) 68 2405–2413.
- 43 Catali A., Zhang D., Janssen L.J. Receptors and signaling pathway underlying relaxations to isoprostanes in canine and porcine airway smooth muscle. *Am. J. Physiol. Lung Cell. Mol. Physiol.* (2002) 283 L1151–L1159.
- 44 Morrow J.D., Awad J.A., Wu A., Zachert W.E., Daniel V.C., Roberts L.J., II. Nonenzymatic free radical-catalyzed generation of thromboxane-like compounds (isothromboxanes) in vivo. *J. Biol. Chem.* (1996) 271 23185–23190.
- 45 Chiabrando C., Valagussa A., Rivalta C. et al. Identification and measurement of endogenous beta-oxidation metabolites of 8-epi-prostaglandin F_{2α}. *J. Biol. Chem.* (1999) 274 1313–1319.
- 46 Roberts L.J., II, Moore K.P., Zachert W.E., Oates J.A., Morrow J.D. Identification of the major urinary metabolite of the F₂-isoprostane 8-iso-prostaglandin F_{2α} in humans. *J. Biol. Chem.* (1996) 271 20617–20620.
- 47 Basu S. Radioimmunoassay of 8-iso-prostaglandin F_{2α}: an index for oxidative injury via free radical catalysed peroxidation. *Prostaglandins Leukot. Essent. Fatty Acids* (1998) 58 319–325.
- 48 Hou X., Roberts L.J., II, Taber D.F. et al. 2,3-Dinor-5,6-dihydro-15-F_{2α}-isoprostane: a bioactive prostanoid metabolite. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* (2001) 281 R391–R400.
- 49 Cracowski J.L., Camus L., Durand T. et al. Response of rat thoracic aorta to F₂-isoprostane metabolites. *J. Cardiovasc. Pharmacol.* (2002) 39 396–403.
- 50 Nourooz-Zadeh J., Halliwell B., Anggard E.E. Evidence for the formation of F₂-isoprostanes during peroxidation of eicosapentaenoic acid. *Biochem. Biophys. Res. Commun.* (1997) 236 467–472.
- 51 Nourooz-Zadeh J., Liu E.H., Anggard E., Halliwell B. F4-isoprostanes: a novel class of prostanoids formed during peroxidation of docosahexaenoic acid (DHA). *Biochem. Biophys. Res. Commun.* (1998) 242 338–344.
- 52 Roberts L.J., II, Montine T.J., Markesbery W.R. et al. Formation of isoprostane-like compounds (neuroprostanes) in vivo from docosahexaenoic acid. *J. Biol. Chem.* (1998) 273 13605–13612.
- 53 Obwegeser R., Ogungho A., Ulm M., Berghammer P., Sinzinger H. Maternal cigarette smoking increases F₂-isoprostanes and reduces prostacyclin and nitric oxide in umbilical vessels. *Prostaglandins Other Lipid Mediat.* (1999) 57 269–279.
- 54 Morrow J.D., Frei B., Longmire A.W. et al. Increase in circulating products of lipid peroxidation (F₂-isoprostanes) in smokers. *N. Engl. J. Med.* (1995) 332 1198–1203.
- 55 Ogungho A., Lupattelli G., Palumbo B., Sinzinger H. Isoprostanes quickly normalize after quitting cigarette smoking in healthy adults. *Vasa* (2000) 29 103–105.
- 56 Montuschi P., Collins J.V., Ciabattini G. et al. Exhaled 8-iso-prostaglandin as an in vivo biomarker of lung oxidative stress in patients with COPD and healthy smokers. *Am. J. Respir. Crit. Care Med.* (2000) 162 1175–1177.
- 57 Chehne F., Ogungho A., Lupattelli G., Budinsky A.C., Palumbo B., Sinzinger H. Increase of isoprostane 8-epi-PGF_{2α} after restarting smoking. *Prostaglandins Leukot. Essent. Fatty Acids* (2001) 64 307–310.
- 58 Ahmadzadehfar H., Ogungho A., Effimios Y., Kritz H., Sinzinger H. Passive cigarette smoking increases isoprostane formation. *Life Sci.* (2006) 78 894–897.
- 59 Schwedhelm E., Bartling A., Lenzen H. et al. Urinary 8-iso-prostaglandin F_{2α} as a risk marker in patients with coronary heart disease: a matched case-control study. *Circulation* (2004) 109 843–848.
- 60 Gross M., Steffes M., Jacobs D.R., Jr et al. Plasma F₂-isoprostanes and coronary artery calcification: the CARDIA Study. *Clin. Chem.* (2005) 51 125–131.

- 61 Greaves K., Dixon S.R., Coker I.O. et al. Influence of isoprostane F2alpha-HI on reflow after myocardial infarction. *Eur. Heart J.* (2004) 25 847–853.
- 62 Iuliano L., Pratico D., Green C. et al. Angioplasty increases coronary sinus F₂-isoprostane formation: evidence for in vivo oxidative stress during PTCA. *J. Am. Coll. Cardiol.* (2001) 37 76–80.
- 63 Kromer B., Tipples J.R. Coronary artery constriction by the isoprostane 8-epi-prostaglandin F_{2α}. *Br. J. Pharmacol.* (1996) 119 1276–1280.
- 64 Cayatte A.J., Du Y., Oliver-Krasinski J., Lavielle G., Verbeuren T.J., Cohen R.A. The thromboxane receptor antagonist S18886 but not aspirin inhibits atherogenesis in apo E-deficient mice: evidence that eicosanoids other than thromboxane contribute to atherosclerosis. *Arterioscler. Thromb. Vasc. Biol.* (2000) 20 1724–1728.
- 65 Pratico D., Tangirala R.K., Rader D.J., Rokach J., Fitzgerald G.A. Vitamin E suppresses isoprostane generation in vivo and reduces atherosclerosis in ApoE-deficient mice. *Nat. Med.* (1998) 4 1189–1192.
- 66 Janick K.P., Belcastro R., Ovcina E. et al. Thromboxane A₂ receptors mediate pulmonary hypertension in 60% oxygen-exposed newborn rats by a cyclooxygenase-independent mechanism. *Am. J. Respir. Crit. Care Med.* (2002) 166 208–214.
- 67 Belhassen L., Pelle G., Dubois-Rande J.L., Adnot S. Improved endothelial function by the thromboxane A₂ receptor antagonist S 18886 in patients with coronary artery disease treated with aspirin. *J. Am. Coll. Cardiol.* (2003) 41 1198–1204.
- 68 Pratico D., Cheng Y., Fitzgerald G.A. TP or not TP: primary mediators in a close runoff? *Arterioscler. Thromb. Vasc. Biol.* (2000) 20 1695–1698.
- 69 Ciabattini G., Davi G., Collura M. et al. In vivo lipid peroxidation and platelet activation in cystic fibrosis. *Am. J. Respir. Crit. Care Med.* (2000) 162 1195–1201.
- 70 Davi G., Di Minno G., Coppola A. et al. Oxidative stress and platelet activation in homozygous homocystinuria. *Circulation* (2001) 104 1124–1128.
- 71 Cracowski J.L., Girollet S., Imbert B. et al. Effects of short-term treatment with vitamin E in systemic sclerosis: a double blind, randomized, controlled clinical trial of efficacy based on urinary isoprostane measurement. *Free Radic. Biol. Med.* (2005) 38 98–103.
- 72 Patrignani P., Panara M.R., Tacconelli S. et al. Effects of vitamin E supplementation on F₂-isoprostane and thromboxane biosynthesis in healthy cigarette smokers. *Circulation* (2000) 102 539–545.
- 73 Meagher E.A., Barry O.P., Lawson J.A., Rokach J., Fitzgerald G.A. Effects of vitamin E on lipid peroxidation in healthy persons. *JAMA* (2001) 285 1178–1182.
- 74 Weinberg R.B., Vander Werken E.S., Anderson R.A., Siegner J.K., Thomas M.J. Pro-oxidant effect of vitamin E in cigarette smokers consuming a high polyunsaturated fat diet. *Arterioscler. Thromb. Vasc. Biol.* (2001) 21 1029–1033.
- 75 Chowienzyk P.J., Brett S.B., Gopaul N.K. et al. Oral treatment with an antioxidant (trolox) reduces oxidative stress and improves endothelial function in men with type II diabetes. *Diabetologia* (2000) 43 974–977.
- 76 Polidori M.C., Pratico D., Ingegni T. et al. Effects of vitamin C and aspirin in ischemic stroke-related lipid peroxidation: results of the AVASAS (Aspirin Versus Ascorbic acid plus Aspirin in Stroke) Study. *Biofactors* (2005) 24 265–274.
- 77 Meagher E.A., Barry O.P., Burke A. et al. Alcohol-induced generation of lipid peroxidation products in humans. *J. Clin. Invest.* (1999) 104 805–813.
- 78 Levine M., Wang Y., Padayatty S.J., Morrow J. A new recommended dietary allowance of vitamin C for healthy young women. *Proc. Natl. Acad. Sci. U. S. A.* (2001) 98 9842–9846.
- 79 Bruno R.S., Leonard S.W., Atkinson J. et al. Faster plasma vitamin E disappearance in smokers is normalized by vitamin C supplementation. *Free Radic. Biol. Med.* (2006) 40 689–697.
- 80 Higdon J.V., Liu J., Du S.H., Morrow J.D., Ames B.N., Wander R.C. Supplementation of postmenopausal women with fish oil rich in eicosapentaenoic acid and docosahexaenoic acid is not associated with greater in vivo lipid peroxidation compared with oils rich in oleate and linoleate as assessed by plasma malondialdehyde and F₂-isoprostanes. *Am. J. Clin. Nutr.* (2000) 72 714–722.
- 81 Mori T.A., Dunstan D.W., Burke V. et al. Effect of dietary fish and exercise training on urinary F₂-isoprostane excretion in non-insulin-dependent diabetic patients. *Metabolism* (1999) 48 1402–1408.
- 82 Visioli F., Caruso D., Galli C., Viappiani S., Galli G., Sala A. Olive oils rich in natural catecholic phenols decrease isoprostane excretion in humans. *Biochem. Biophys. Res. Commun.* (2000) 278 797–799.
- 83 Wiseman H., O'Reilly J.D., Adlercreutz H. et al. Isoflavone phytoestrogens consumed in soy decrease F₂-isoprostane concentrations and increase resistance of low-density lipoprotein to oxidation in humans. *Am. J. Clin. Nutr.* (2000) 72 395–400.
- 84 Sanchez-Moreno C., Cano M.P., de Ancos B. et al. Mediterranean vegetable soup consumption increases plasma vitamin C and decreases F(2)-isoprostanes, prostaglandin E(2) and monocyte chemotactic protein-1 in healthy humans. *J. Nutr. Biochem.* (2006) 17 183–189.
- 85 Wiswedel I., Hirsch D., Kropf S. et al. Flavanol-rich cocoa drink lowers plasma F(2)-isoprostane concentrations in humans. *Free Radic. Biol. Med.* (2004) 37 411–421.
- 86 Minuz P., Patrignani P., Gaiuso S. et al. Increased oxidative stress and platelet activation in patients with hypertension and renovascular disease. *Circulation* (2002) 106 2800–2805.
- 87 Cracowski J.L., Baguet J.P., Ormezzano O. et al. Lipid peroxidation is not increased in patients with untreated mild-to-moderate hypertension. *Hypertension* (2003) 41 286–288.
- 88 Ward N.C., Hodgson J.M., Puddey L.B., Mori T.A., Beilin L.J., Croft K.D. Oxidative stress in human hypertension: association with antihypertensive treatment, gender, nutrition, and lifestyle. *Free Radic. Biol. Med.* (2004) 36 226–232.

- 89 Mallat Z., Philip I., Lebret M., Chatel D., Macioun J., Tedgui A. Elevated levels of 8-iso-prostaglandin F_{2a} in pericardial fluid of patients with heart failure: a potential role for in vivo oxidant stress in ventricular dilatation and progression to heart failure. *Circulation* (1998) 97 1536–1539.
- 90 Li H., Lawson J.A., Reilly M. et al. Quantitative high performance liquid chromatography/tandem mass spectrometric analysis of the four classes of $F(2)$ -isoprostanes in human urine. *Proc. Natl. Acad. Sci. U. S. A.* (1999) 96 13381–13386.
- 91 Cracowski J.L., Tremel F., Marpeau C. et al. Increased formation of F_2 -isoprostanes in patients with severe heart failure. *Heart* (2000) 84 439–440.
- 92 Nonaka-Sarukawa M., Yamamoto K., Aoki H. et al. Increased urinary isoprostane excretion in patients with non-ischemic congestive heart failure: a marker of oxidative stress. *Heart* (2003) 89 871–874.
- 93 Gniwotta C., Morrow J.D., Roberts L.J., Kühn H. Prostaglandin F_2 -like compounds, F_2 -isoprostanes, are present in increased amounts in atherosclerotic lesions. *Arterioscler. Thromb. Vasc. Biol.* (1997) 17 3236–3241.
- 94 Pratico D., Iuliano L., Mauricillo A. et al. Localization of distinct F_2 -isoprostanes in human atherosclerotic lesions. *J. Clin. Invest.* (1997) 100 2028–2034.
- 95 Mehrabi M.R., Ekmekcioglu C., Tatzber F. et al. The isoprostane, 8-epi- PGF_{2a} , is accumulated in coronary arteries isolated from patients with coronary heart disease. *Cardiovasc. Res.* (1999) 43 492–499.
- 96 Lindsay T.F., Luo X.F., Lehotay D.C. et al. Ruptured abdominal aortic aneurysm, a 'two-hit' ischemia/reperfusion injury: evidence from an analysis of oxidative products. *J. Vasc. Surg.* (1999) 30 219–228.
- 97 Delanty N., Reilly M.P., Pratico D. et al. 8-epi PGF_{2a} generation during coronary reperfusion. *Circulation* (1997) 95 2492–2499.
- 98 Cipellone F., Ciabattini G., Patrignani P. et al. Oxidant stress and aspirin-insensitive thromboxane biosynthesis in severe unstable angina. *Circulation* (2000) 102 1007–1013.
- 99 Reilly M.P., Delanty N., Roy L. et al. Increased formation of the isoprostanes IPF_{2a} -I and 8-epi-prostaglandin F_{2a} in acute coronary angioplasty: evidence for oxidant stress during coronary reperfusion in humans. *Circulation* (1997) 96 3314–3320.
- 100 Stein C.M., Tanner S.B., Awad J.A., Roberts L.J., Morrow J.D. Evidence of free radical-mediated injury (isoprostane overproduction) in scleroderma. *Arthritis Rheum.* (1996) 39 1146–1150.
- 101 Cracowski J.L., Carpentier P.H., Imbert B. et al. Increased urinary F_2 -isoprostanes in systemic sclerosis, but not in primary Raynaud's phenomenon: effect of cold exposure. *Arthritis Rheum.* (2002) 46 1319–1323.
- 102 Cracowski J.L., Marpeau C., Carpentier P.H. et al. Enhanced in vivo lipid peroxidation in scleroderma spectrum disorders. *Arthritis Rheum.* (2001) 44 1143–1148.
- 103 Volpe A., Biasi D., Caramaschi P. et al. Levels of F_2 -isoprostanes in systemic sclerosis: correlation with clinical features. *Rheumatology (Oxford)* (2005) 45 314–320.
- 104 Cracowski J.L., Kom G.D., Salvat-Melis M. et al. Post occlusive reactive hyperemia inversely correlates to urinary 15- F_2t -isoprostane levels in systemic sclerosis. *Free Radic. Biol. Med.* (2006) 40 1732–1737.
- 105 Pratico D., Ferro D., Iuliano L. et al. Ongoing prothrombotic state in patients with antiphospholipid antibodies: a role for increased lipid peroxidation. *Blood* (1999) 93 3401–3407.
- 106 Martinuzzi M.E., Forastiero R.R., Kordich L., Carreras L.O. Increased lipid peroxidation correlates with platelet activation but not with markers of endothelial cell and blood coagulation activation in patients with antiphospholipid antibodies. *Br. J. Haematol.* (2001) 114 845–851.
- 107 Cracowski J.L., Cracowski C., Bessard G. et al. Increased lipid peroxidation in patients with pulmonary hypertension. *Am. J. Respir. Crit. Care Med.* (2001) 164 1038–1042.
- 108 Van Kooten F., Ciabattini G., Patrono C., Dippel D.W., Koudstaal P.J. Platelet activation and lipid peroxidation in patients with acute ischemic stroke. *Stroke* (1997) 28 1557–1563.
- 109 Helmersson J., Mattsson P., Basu S. Prostaglandin F_2 metabolite and F_2 -isoprostane excretion in migraine. *Clin. Sci.* (2002) 102 39–43.
- 110 Regan C.L., Levine R.J., Baird D.D. et al. No evidence for lipid peroxidation in severe preeclampsia. *Am. J. Obstet. Gynecol.* (2001) 185 572–578.
- 111 Walsh S.W., Vaughan J.E., Wang Y., Roberts L.J., II Placental isoprostane is significantly increased in preeclampsia. *FASEB J.* (2000) 14 1289–1296.
- 112 Morris J.M., Gopaul N.K., Endresen M.J. et al. Circulating markers of oxidative stress are raised in normal pregnancy and pre-eclampsia. *Br. J. Obstet. Gynaecol.* (1998) 105 1195–1199.
- 113 Barden A., Beilin L.J., Ritchie J., Croft K.D., Walters B.N., Michael C.A. Plasma and urinary 8-iso-prostane as an indicator of lipid peroxidation in pre-eclampsia and normal pregnancy. *Clin. Sci. (Colch)* (1996) 91 711–718.
- 114 McKinney E.T., Shouri R., Hunt R.S., Ahokas R.A., Sibai B.M. Plasma, urinary, and salivary 8-epi-prostaglandin F_{2a} levels in normotensive and preeclamptic pregnancies. *Am. J. Obstet. Gynecol.* (2000) 183 874–877.
- 115 Parra M., Rodrigo R., Barja P. et al. Screening test for preeclampsia through assessment of uteroplacental blood flow and biochemical markers of oxidative stress and endothelial dysfunction. *Am. J. Obstet. Gynecol.* (2005) 193 1486–1491.
- 116 Harseni N.K., Bracke K., Staff A.C. Augmented oxidative stress as well as antioxidant capacity in maternal circulation in preeclampsia. *Eur. J. Obstet. Gynecol. Reprod. Biol.* (2005) [Epub ahead of print].

Arteriosclerosis, Thrombosis, and Vascular Biology

Volume 26 Number 2 February 2006



This article contains online supplementary information that is available at <http://www.atvbaha.org>.

Brief Reviews

Diet and Murine Atherosclerosis

Godfrey S. Getz, Catherine A. Reardon

242

Murine models of atherosclerosis have been fed a variety of diets that vary in the level of cholesterol, the level and type of fatty acid, and the absence and presence of cholate. This review summarizes what is known about the effect of these dietary components on lipoprotein levels and/or atherosclerosis.

Smoking, Metalloproteinases, and Vascular Disease

Todd S. Perlestein, Richard T. Lee

250

The matrix metalloproteinases are emerging as strong candidate mediators of smoking-associated vascular disease. Smoking-induced inflammation and oxidative stress may increase metalloproteinase transcription, increase pro-enzyme activation, and limit endogenous inhibition of metalloproteinase activity. The relationship between smoking, metalloproteinases, and vascular disease is discussed in this brief review.

Influence of Cardiovascular Risk Factors on Endothelial Progenitor Cells: Limitations for Therapy?—ATVB In Focus

Nikos Werner, Georg Nickenig

257

Circulating endothelial progenitor cells play an important role in restoration of the endothelium after endothelial cell damage. The current review focuses on the role of cardiovascular risk factors on endothelial cell apoptosis and progenitor cell-mediated vasculoprotection.

Endothelial NO Synthase: Host Defense Enzyme of the Endothelium?

Ton J. Rabelink, Thomas F. Luscher

267

This article explores the physiology of superoxide generation by endothelial nitric oxide synthase, the so-called "uncoupled" state of the enzyme. The fact that this alternative chemistry of the eNOS enzyme is evolutionary strongly conserved, suggests that it may play a physiological role. It is proposed that this uncoupled state may contribute to defense against infections, and the central role of uncoupled eNOS in redox signaling in the endothelium may open up new avenues for therapy to prevent atherosclerosis.

Vascular Biology

Importance of Junctional Adhesion Molecule-A for Neointimal Lesion Formation and Infiltration in Atherosclerosis-Prone Mice

Alma Zernecke, Elisa A. Liehn ... Christian Weber

Web Site Feature

e10-e13

Abstract

272

Junctional adhesion molecule-A (JAM-A) has recently been implicated in leukocyte recruitment on early atherosclerotic endothelium. Our data provide the first evidence to our knowledge for a crucial role of JAM-A in accelerated lesion formation and monocyte infiltration in atherosclerosis-prone mice.

Aspirin Has A Gender-Dependent Impact on Antiinflammatory 15-Epi-Lipoxin A₄ Formation: A Randomized Human Trial

Nan Chiang, Shelley Hurwitz ... Charles N. Serhan

Web Site Feature

e14-e17

Abstract

273

In a double-blinded clinical trial, low-dose aspirin significantly increased the antiinflammatory mediator, aspirin-triggered-15-epi-lipoxinA₄. A post-trial analysis showed that the age trends in aspirin-triggered-15-epi-lipoxinA₄ formation are significantly different with genders: a positive correlation in women and a negative correlation in men. Therefore, aspirin-triggered-15-epi-lipoxinA₄ may be related to the gender-specific clinical impact of aspirin.

ARTERIOSCLEROSIS, THROMBOSIS, AND VASCULAR BIOLOGY (ISSN 1079-5642) is published monthly by Lippincott Williams & Wilkins, at 16522 Hunters Green Parkway, Hagerstown, MD 21740. Business offices are located at 530 Walnut Street, Philadelphia, PA 19106-3621. Production offices are located at 351 West Camden Street, Baltimore, MD 21201-2436. For institutional rates in Japan, contact Igaku-Shoin MYW, Ltd., 3-23-14 Hongo, Bunkyo-ku, Tokyo 113, Japan. Individuals may subscribe for their personal use at the following annual rates: \$217 for members of an American Heart Association scientific council and \$355 for nonmembers; international: \$266 for members of an American Heart Association scientific council and \$419 for nonmembers. Periodicals postage paid at Hagerstown, MD and additional mailing offices. POSTMASTER: Send address changes to ARTERIOSCLEROSIS, THROMBOSIS, AND VASCULAR BIOLOGY, American Heart Association, Lippincott Williams & Wilkins, 12107 Insurance Way, Hagerstown, MD 21740.

Conjugated Linoleic Acid Impairs Endothelial Function

Justin S.W. Taylor, Simon R.P. Williams, Rhian Rhys, Phillip James, Michael P. Frenneaux

Objectives—To determine the effect of dietary supplementation with conjugated linoleic acid (CLA) on body mass index (BMI), body fat distribution, endothelial function, and markers of cardiovascular risk.

Methods and Results—Forty healthy volunteers with BMI >27 kg/m² were randomized to receive a CLA isomeric mixture or olive oil in a 12-week double-blind study. Subcutaneous body fat and abdominal/hepatic fat content were assessed using skin-fold thicknesses and computed tomography scanning, respectively. Endothelial function was assessed by brachial artery flow-mediated dilatation (FMD). Plasma isoprostanes were measured as an index of oxidative stress. CLA supplementation did not result in a significant change in BMI index or total body fat. There was a significant decrease in limb (−7.8 mm, $P<0.001$), but not torso skin-fold thicknesses or abdominal or liver fat content. Brachial artery FMD declined (−1.3%, $P=0.013$), and plasma F2-isoprostanes increased (+91pg/mL, $P=0.042$).

Conclusions—A CLA isomeric mixture had at most modest effects on adiposity and worsened endothelial function. On the basis of these results, the use of the isomeric mixture of CLA as an aid to weight loss cannot be recommended. (*Arterioscler Thromb Vasc Biol.* 2006;26:307–312.)

Key Words: body composition ■ conjugated linoleic acid ■ endothelial function ■ obesity ■ oxidative stress

Abdominal obesity¹ and the associated dysmetabolic syndrome² confer increased cardiovascular risk. Dietary modification with n-3 polyunsaturated fatty acids appears to reduce the risk of coronary artery disease and improve mortality.^{3–5} Conjugated linoleic acid (CLA) is a naturally occurring fatty acid and is found in dairy products and meat from ruminants. It differs from the better known linoleic acid by having an extra carbon-carbon double bond. It has been widely promoted in the lay press,⁶ with claims that it can prevent and treat cancer,⁷ prevent heart disease,⁸ improve immune function,⁹ and treat obesity.¹⁰ Whereas some of these effects are supported by studies in animals,^{8,11} there is conflicting published human research on CLA, and in particular there have been no conclusive studies measuring its effect on markers of cardiovascular risk. Furthermore, it is now realized that the different isomers of CLA may have very different biological properties and may have different mechanisms of action.¹² The 2 main CLA isomers that have been studied are 9,11 and 10,12 CLA. Other studies have suggested that CLA supplementation may increase oxidative stress, although this was not proven.^{13,17} Furthermore, the small number of studies thus far published have suggested that in humans, weight loss produced by CLA supplementation is at most modest.^{10,13,16,18} To assess the efficacy of CLA as an aid to weight loss and its effect on cardiovascular risk factors, we undertook a double blind study examining the effects of a commercially available isomeric mixture of CLA on body weight, body fat mass and distribution, endothelial

function, insulin sensitivity, and markers of oxidative stress in overweight middle-aged men.

Methods

Subjects

Forty nonsmoking white men, aged 35 to 60, without diabetes, hypertension, or cardiovascular disease, with a body mass index (BMI) >27 kg/m² were recruited from the local community through media advertisements. All subjects gave informed written consent and the protocol was approved by the local research and ethics committee.

Protocol

Subjects were randomly assigned to receive 4.5 g/d of CLA (isomeric mixture 60 calories/d) or olive oil (54 calories/d). The randomization was performed by an independent observer who also matched subjects by age and BMI. The isomeric mixture contained 35% 9c,11t CLA, 36% t10,c12 CLA, 1% to 2% 9c,11c and 10c,12c CLA, 1.5% 9t,11t and 10t,11t CLA, and <1% t8,c10 and c11,t13 CLA. The CLA and olive oil capsules were supplied by Natural Lipids (Hovdebygd, Norway). All vascular measurements were made in the morning after an overnight fast, at the beginning of the study, and after 12 weeks of supplementation.

Body Composition

Body weight and height were measured and BMI calculated. Skin-fold thicknesses were measured according to standard guidelines¹⁹ with Harpenden skinfold calipers (Holtain Ltd, Crymych, UK) at the following sites: biceps, triceps, front mid-thigh, medial calf, subscapular, mid-axillary, and abdominal. Abdominal, waist, and hip girths were measured. All measurements were made in triplicate and averaged. Bioelectrical impedance analysis was performed using the

Original received January 24, 2005; final version accepted November 29, 2005.

From Morriston Hospital, Swansea, UK; University of Glamorgan, Pontypridd, UK; Royal Glamorgan Hospital, Llantrisant, UK; Wales Heart Research Institute, Cardiff, UK; Dept Cardiovascular Medicine, The University of Birmingham, Birmingham, UK.

Source of funding: British Heart Foundation Project Grant.

Correspondence to Justin S. W. Taylor, Department of Cardiology, Morriston Hospital, Morriston, Swansea SA6 6NL, UK. E-mail jswt@totalise.co.uk
© 2006 American Heart Association, Inc.

Arterioscler Thromb Vasc Biol. is available at <http://www.atvbaha.org>

DOI: 10.1161/01.ATV.0000199679.46501.ac

tetrapolar method and a Bodystat 1500 analyser (Bodystat Ltd, Isle of Man). Abdominal adipose tissue and liver fat were measured using two computed tomography (CT) images as described previously.²⁰ Abdominal adipose tissue is presented as a surface area at the level of the fourth lumbar vertebrae, and hepatic and splenic fat is presented as radiographic density in Hounsfield units (HU). Images were acquired using a Somatom Plus 4 scanner. CT slices were 10 mm in thickness and were obtained at 120 kV and 200 mA, with a 42-cm field of view and a 512×512 matrix. Image analysis was performed using dedicated software (SliceOmatic Version 4.2; Tomovision, Montreal, Canada).

Endothelial Function

Changes in brachial artery diameter in response to reactive hyperemia (FMD) were measured noninvasively using a high-resolution ultrasonic wall-tracking system (Vadirec Wall-track System™) as previously validated.^{21,49,50} Studies were performed at a controlled temperature of 21°C, with subjects supine and their arm held outstretched on a cushion. Baseline measurements of internal brachial artery diameter were taken after 15 minutes of rest. Reactive hyperemia was produced by releasing a pediatric sphygmomanometer wrist cuff inflated to systolic pressure plus 50 mm Hg for 5 minutes. Internal brachial artery diameter was measured every minute after cuff release, and the maximum change from baseline was used to calculate FMD. Data are presented as the percentage diameter change from baseline in the brachial artery.

Laboratory Measurements

Venous blood samples were freshly analyzed for glucose, insulin, cholesterol (including low-density lipoprotein and high-density lipoprotein), and C-reactive protein. Further samples were centrifuged and the supernatant frozen at -80°C. These samples were analyzed later using enzyme immuno-linked assays for leptin (Alexis), adiponectin (Biogenesis UK), F2-isoprostanes (Alexis), and tumor necrosis factor- α (paired antibody enzyme-linked immunosorbent assay). Insulin sensitivity was measured indirectly using HOMA-IR [fasting serum insulin (μ U/mL)×fasting plasma glucose (mmol/L)/22.5].

Statistical Analysis

Analysis was performed using SPSS v11.5. Baseline values are presented as mean±SD. Variables with a skewed distribution were logarithmically transformed to achieve a normal distribution prior to analysis. Baseline comparison between the CLA and olive oil groups was assessed by Student *t* test. The effects of treatment with CLA versus olive oil were assessed by using ANCOVA using baseline values as covariates and are presented with 95% confidence intervals. To avoid type I errors, post-hoc Bonferroni corrections were applied to the groups of primary objective measurements. Therefore, 2-tailed $P<0.025$, $P<0.005$, $P<0.001$, $P<0.025$, and $P<0.008$ were regarded as significant for blood pressure, skin-folds, girths, FMD, and abdominal CT measurements, respectively. A 2-tailed $P<0.05$ was regarded as significant for primary objective measurements weight, and bioimpedance, and the exploratory blood analyses.

Results

Baseline measurements did not differ significantly in the olive oil (19 subjects) and CLA (21 subjects) groups (Table 1). All subjects completed the study. The effect of 12 weeks of supplementation is shown in Table 2. The important changes are also displayed in the Figure. There were no significant changes in parameters in the olive oil group except for a decline in tumor necrosis factor- α (-61 pg/mL [95% CI, -3 to -120]; $P=0.04$). In the CLA group there was no significant change in body mass (-1.1 kg [95% CI, -2.3 to 0.04]; $P=0.06$), BMI (-0.4 kg/m² [95% CI, -0.8 to 0.03]; $P=0.07$), or total body fat (-1% [95% CI, -2.5 to 0.5];

TABLE 1. Characteristics of Olive Oil and CLA Groups at the Start of the Study

Baseline Characteristic	Olive Oil (n=19)	CLA (n=21)	P
Age, y	47±8	45±6	NS
Mass, kg	97±13	101±9	NS
BMI	33±3	33±3	NS
Fat, % (Bioimpedance)	29±3	28±4	NS
Systolic BP, mm Hg	128±13	122±10	NS
Diastolic BP, mm Hg	85±7	80±8	NS
Anthropometric measurements			
Skin-folds: Biceps, mm	12±4	11±4	NS
Skin-folds: Triceps, mm	20±5	19±5	NS
Skin-folds: Mid-thigh, mm	23±7	25±8	NS
Skin-folds: Medial calf, mm	16±4	16±5	NS
Skin-folds: Subscapular, mm	32±6	32±7	NS
Skin-folds: Mid-axillary, mm	23±4	24±6	NS
Skin-folds: Abdominal, mm	33±7	35±6	NS
Skin-folds: Limb, mm	72±17	72±18	NS
Skin-folds: Torso, mm	123±13	128±16	NS
Skin-folds: Torso:Limb	1.8±0.4	1.9±0.3	NS
Girth: Abdomen, cm	107±7	107±6	NS
Girth: Waist, cm	112±7	112±7	NS
Girth: Hip, cm	110±6	110±5	NS
Girth: Waist:Hip	1.02±0.03	1.01±0.04	NS
Girth: Abdominal:Hip	0.98±0.05	0.97±0.05	NS
Endothelial function			
Brachial artery, mm	3.8±0.5	3.7±0.4	NS
Brachial artery FMD, %	2.1±2.1	2.2±2.7	NS
Abdominal CT measurements			
Liver, HU	80±13	66±10	NS
Spleen, HU	64±2	61±2	NS
Liver:Spleen	0.93±0.18	1.08±0.18	NS
Visceral area, cm ²	242±82	204±70	NS
Subcutaneous fat area, cm ²	341±80	357±93	NS
Visceral:Subcutaneous fat	0.75±0.38	0.63±0.37	NS
Blood analyses			
LDL cholesterol, mmol/L	3.6±0.9	3.4±0.8	NS
HDL cholesterol, mmol/L	1.0±0.3	1.1±0.3	NS
Total cholesterol, mmol/L	5.6±0.9	5.4±0.8	NS
Triglycerides, mmol/L	2.3±1.3	1.8±1.0	NS
CRP, mg/L	5.2±2.9	4.8±2.7	NS
Glucose, mmol/L	5.2±0.8	5.1±0.7	NS
Insulin, mU/L	18±22	13±7	NS
HOMA-IR	4.8±6.4	3.1±1.8	NS
Leptin, ng/mL	21±12	22±16	NS
Adiponectin, pg/mL	8±6	9±6	NS
F2-isoprostanes, pg/mL	135±67	99±103	NS
Tumor necrosis factor- α , pg/mL	122±97	108±145	0.04

NS indicates not significant.


$P=0.18$). There was a significant decrease in limb (-7.8 mm [95% CI, -11.1 to -4.5]; $P<0.001$), but not torso (-4.0 mm [95% CI, -11.5 to 3.5]; $P=0.29$) skin folds, and a significant increase in the total torso-to-limb skin-fold ratio(+0.13 [95%

TABLE 2. Change in Measurements in Each Treatment Group After 12 Weeks of Supplementation Plus ANCOVA Analyses Showing the Overall Effect of Supplementation With CLA

	Olive Oil	CLA	Change in Measurement (ANCOVA) [95%CI]
Mass, kg	0.9±1.4	-0.2±1.9	-1.1 [-2.3 to 0.04] (<i>P</i> =0.06)
BMI	0.3±0.5	-0.1±0.7	-0.4 [-0.8 to 0.03] (<i>P</i> =0.07)
Fat, % (Bioimpedance)	1.1±2.9	0.1±1.7	-1 [-2.5 to 0.5] (<i>P</i> =0.18)
Systolic BP, mm Hg	0.2±6.7	-0.4±9.1	-2.2 [-7.1 to 2.7] (<i>P</i> =0.37)
Diastolic BP, mm Hg	-0.8±7.5	0.1±5.7	-1.3 [-5.4 to 2.8] (<i>P</i> =0.52)
Anthropometric measurements			
Skin-folds: Biceps, mm	0.1±2.3	-1.4±1.5	-1.6 [-2.7 to -0.5] (<i>P</i> =0.005)
Skin-folds: Triceps, mm	0.3±1.9	-1.1±2.4	-1.5 [-2.9 to 0.03] (<i>P</i> =0.046)
Skin-folds: Mid-thigh, mm	0.4±2.1	-2.6±2.6	-2.8 [-4.3 to -1.3] (<i>P</i> <0.001)
Skin-folds: Medial calf, mm	0.2±1.4	-1.5±1.5	-1.6 [-2.7 to -0.6] (<i>P</i> =0.001)
Skin-folds: Sub-scapular, mm	0.9±3.3	-0.3±4.8	-1.3 [-3.6 to 1.2] (<i>P</i> =0.29)
Skin-folds: Mid-axillary, mm	-0.4±2.8	-3.0±3.0	-2.3 [-4.0 to -0.6] (<i>P</i> =0.011)
Skin-folds: Abdominal, mm	2.5±6.8	2.2±3.4	+1.4 [-1.8 to 4.3] (<i>P</i> =0.36)
Skin-folds: Limb, mm	1.1±3.5	-6.6±6.0	-7.8 [-11.1 to -4.5] (<i>P</i> <0.001)
Skin-folds: Torso, mm	2.9±13.5	-2.1±9.9	-4.0 [-11.5 to 3.5] (<i>P</i> =0.29)
Skin-folds: Torso:Limb	0.02±0.18	0.15±0.15	+0.13 [0.03 to 0.24] (<i>P</i> =0.017)
Girth: Abdomen, cm	0.1±1.3	-0.1±1.9	-0.2 [-1.4 to 0.9] (<i>P</i> =0.69)
Girths: waist, cm	-0.3±2.1	0.5±2.5	+0.8 [-0.8 to 2.4] (<i>P</i> =0.30)
Girths: hip, cm	0.4±1.4	-0.01±1.5	-0.4 [-1.3 to 0.6] (<i>P</i> =0.44)
Girths: Waist:Hip	-0.003±0.03	0.01±0.22	+0.01 [-0.01 to 0.02] (<i>P</i> =0.44)
Girths: Abdominal:Hip	-0.01±0.03	-0.01±0.01	+0.01 [-0.01 to 0.02] (<i>P</i> =0.42)
Endothelial function			
Brachial artery, mm	-0.17±0.60	-0.03±0.62	+0.06 [-0.29 to 0.41] (<i>P</i> =0.72)
Brachial artery FMD, %	0.1±2.3	-1.3±2.5	-1.3 [-2.4 to -0.3] (<i>P</i> =0.013)
Abdominal CT measurements			
Liver, HU	-0.7±5.7	-4.1±10.2	-3.9 [-9.9 to 2.2] (<i>P</i> =0.20)
Spleen, HU	-2.2±4.8	0.7±2.8	1.1 [-1.7 to 3.8] (<i>P</i> =0.44)
Liver:Spleen	0.01±0.11	-0.08±0.17	-0.09 [-0.21 to 0.03] (<i>P</i> =0.12)
Visceral area, cm ²	11.1±34.5	3.7±55.0	-4.5 [-33.6 to 24.6] (<i>P</i> =0.76)
Subcutaneous fat area, cm ²	-9.8±32.6	3.43±40.0	-12.6 [-39.5 to 14.2] (<i>P</i> =0.34)
Visceral:Subcutaneous fat	-0.09±0.26	0.03±0.29	0.08 [-0.11 to 0.26] (<i>P</i> =0.41)
Blood analyses			
LDL cholesterol, mmol/L	-0.2±0.5	-0.01±0.4	+0.1 [-0.1 to 0.3] (<i>P</i> =0.41)
HDL cholesterol, mmol/L	-0.03±0.11	-0.09±0.13	-0.03 [-0.10 to 0.05] (<i>P</i> =0.46)
Total cholesterol, mmol/L	-0.09±0.50	-0.07±0.44	-0.07 [-0.34 to 0.20] (<i>P</i> =0.62)
Triglycerides, mmol/L	0.1±0.9	0.1±0.8	-0.1 [-0.6 to 0.4] (<i>P</i> =0.57)
CRP, mg/L	-0.8±2.9	0.4±4.8	+0.9 [-1.4 to 3.2] (<i>P</i> =0.41)
Glucose, mmol/L	0.3±0.8	0.2±0.8	-0.1 [-0.5 to 0.3] (<i>P</i> =0.51)
Insulin, mIU/L	0.7±22.2	2.3±12.6	-3 [-11 to 6] (<i>P</i> =0.50)
HOMA-IR	0.2±6.7	0.8±3.2	-0.8 [-3.2 to 1.6] (<i>P</i> =0.50)
Leptin, ng/mL	1.3±7.3	0.7±5.1	-0.7 [-4.8 to 3.4] (<i>P</i> =0.72)
Adiponectin, pg/mL	0.5±3.4	-0.2±4.0	-0.2 [-2.4 to 2.0] (<i>P</i> =0.84)
F2-isoprostanes, pg/mL	-36±95	94±200	+91 [3 to 178] (<i>P</i> =0.042)
Tumor necrosis factor-α, pg/mL	-51±70	15±21	+45 [-11 to 101] (<i>P</i> =0.11)

CI, 0.03 to 0.24]; *P*=0.017). There was no significant change in abdominal, waist, or hip girths, or in subcutaneous abdominal fat and liver fat measured by CT. However, there was a significant decrease in brachial artery FMD (-1.3% [95% CI,

-2.4 to -0.3]; *P*=0.013), and a significant increase in plasma F2-isoprostanes(+91pg/mL [95% CI, 3 to 178]; *P*=0.042). There was no change in estimated insulin sensitivity, total cholesterol, low-density lipoprotein cholesterol,



Main findings of changes caused by CLA using baseline values as covariates (ANCOVA).

high-density lipoprotein cholesterol, triglycerides, CRP, leptin, or adiponectin.

There was a significant negative correlation between change in F2-isoprostanes and change in total limb skin folds (ie, loss of limb skin-fold thickness was associated with an increase in F2-isoprostanes) for the entire group (CLA + olive oil) ($P=0.012$) but no significant correlation when each group was analyzed separately. There was no significant correlation between change in endothelial function and change in F2-isoprostanes or change in limb skin-fold thicknesses.

Discussion

Obesity, and in particular abdominal obesity, is associated with increased cardiovascular risk,³²⁻³⁴ and intentional weight reduction improves cardiovascular risk.^{35,36} Recently there has been a great deal of interest in the effect of the Mediterranean diet on cardiovascular risk. A 2-year study examining the effect of the Mediterranean diet in patients with the metabolic syndrome found a reduction in body weight, an improvement in endothelial function, a decrease in CRP and an improvement in insulin resistance.³⁷ A study that tried to identify which component of the Mediterranean diet was responsible for improving cardiovascular risk paradoxically found that olive oil impaired endothelial function, although this was inversely correlated with changes in triglycerides.²⁸ The conclusion of the study was that it was the antioxidant and omega-3-rich foods that conferred cardiovascular benefit. Nevertheless, these studies supported the view that dietary modification or supplementation may have a significant impact on obesity and, in particular, cardiovascular risk.

Experimental evidence in animal models suggests that CLA supplementation, in particular the 10,12 CLA isomer, induces fat mass loss.²⁹⁻³² On the basis of this initial evidence, there has been a great deal of interest in its use as an aid to lose fat and weight in humans. Blankson et al reported that 12-week supplementation with >3.4 g/d isomeric CLA significantly reduced body fat mass in overweight volunteers, although there was no change in weight or BMI,¹⁰

and Riserus et al found that only 4-week supplementation with 4.2 g/d isomeric CLA significantly improved sagittal abdominal diameter.³³ However, Zambell et al found no significant change in weight, BMI, or fat mass after 3-g/d supplementation with isomeric CLA (in women who were not overweight).¹⁸ The observation in two studies of an impairment of insulin sensitivity have raised concerns.¹³⁻¹⁵ This study was therefore designed to assess the effect of CLA supplementation on BMI, body fat distribution, and markers of cardiovascular risk, including endothelial function.

Our study found that an isomeric mixture of CLA did not cause significant weight loss (although there was a trend to weight loss of 1.1 kg). This is consistent with the 0.24 to 0.46 kg weight loss reported in previous studies.^{13,18} Although CLA reduced limb fat, it had no effect on abdominal fat or liver fat (although there was a nonsignificant trend to an increase in liver fat, measured as a decrease in liver density in Hounsfield units). This finding is in contrast to a previous report that found a decrease in sagittal abdominal diameter after 4 weeks of 4.2 g/d CLA.³³ However, the suggestion that CLA may have a lipodystrophic effect is not new. A study supplementing mice with isomeric CLA found a reduction in fat mass, liver hypertrophy, and an increase in insulin resistance,³⁴ whereas mice fed the 10,12 CLA isomer had hyperinsulinemia and an increase in liver fat develop.³⁵ The mechanism for this is not clear, although a rapid decrease in leptin and adiponectin has been observed in mice only 2 days after starting CLA supplementation.³⁶ A decrease in leptin has also been observed in rats.³⁷ This hypothesis is supported by the observation that hyperinsulinemia and liver steatosis are partially reversed when hypoleptinemia is normalized by leptin infusion in CLA lipodystrophic mice.³⁴ However, there is conflicting evidence regarding the effects of CLA supplementation on plasma leptin in humans. One study supplementing patients with type 2 diabetes with CLA found a decrease in leptin,³⁸ but another study supplementing obese men with CLA found no change in leptin.¹³ No clinical studies in humans have measured adiponectin after CLA supplementation. In contrast to these findings, a study supplementing Zucker diabetic fatty rats with CLA found that the previous impaired glucose tolerance improved.³⁹

Cell culture and animal studies have suggested several other potential mechanisms by which CLA may reduce body fat, including reducing apolipoprotein B secretion in HepG2 cells,⁴⁰ increasing carnitine palmitoyltransferase activity and decreasing lipoprotein lipase activity,³⁰ and increasing tumor necrosis factor and uncoupling protein levels.³⁴ The role of peroxisome proliferator-activated receptor gamma (PPAR γ) activity is not clear. PPAR γ activity was increased in Zucker diabetic fatty rats and genetically obese mice fed CLA,^{39,41,42} although several *in vitro* studies have found that 10,12 CLA downregulates PPAR γ activity in mice adipocytes.^{43,44} *In vivo* and *in vitro* studies in pigs have also found that CLA induced an increase in PPAR γ activity.⁴⁵ However, human studies have found an opposite effect on PPAR γ . Human adipocytes cultured *in vitro* with 10,12 CLA decreased the expression of PPAR γ ,^{41,43} and diabetic patients treated with PPAR γ agonists (glitazones) experience decreased central fat and increased peripheral fat.⁴⁶

Our observations are important because reducing nonabdominal fat is less likely to reduce cardiovascular risk, and an increase in hepatic fat will increase insulin requirements.⁴⁷ Thus any weight loss with this regime is at most modest and the pattern of weight loss is not metabolically favorable.

Furthermore, we found that CLA significantly impaired brachial artery endothelial function, consistent with an adverse impact on cardiovascular risk.⁵¹ The mechanism of this effect is not clear. We and others¹⁶ found an increase in F₂ isoprostanes, a lipid peroxidation product generally considered to be a marker of increased oxidative stress. Another previous study reported an increase in plasma CRP, a marker of inflammation.¹⁷ Taken together, these data suggest that this CLA regime impairs endothelial function and that this may, at least in part, be caused by increased oxidative stress. It is possible that the observed change in FMD in this study has been caused by the change in limb skin-fold thicknesses interfering with the FMD technique (for example, by altering wrist arterial occlusion pressure or changing depth from probe to brachial artery), thus giving a false measurement. However, we feel that this is unlikely because a high wrist arterial occlusion pressure was used and a change in depth to the brachial artery of 1 to 2 mm is well within the capability of the ultrasound probe. Brachial artery FMD measurements vary markedly from laboratory to laboratory, dependant on the exact technique used.⁴⁸ In this study, we used the wrist cuff technique, which results in lower values of brachial artery FMD than those obtained using the upper arm or mid-forearm techniques. We did not observe an adverse impact on insulin sensitivity. However, the study may have been under-powered to detect differences using the HOMA technique.⁵² We did not identify any change in plasma lipid profiles.

The effect of CLA on other cardiovascular risk factors has been examined. One study reported that CLA decreased platelet aggregability,⁵³ but another reported no change in platelet aggregability.⁵⁴ A small decrease in total, low-density lipoprotein, and high-density lipoprotein cholesterol was found in overweight men taking 1.7 g/d CLA and 3.4 g/d CLA, although this was not maintained at higher doses.¹⁰ A further study in obese men using 10,12 CLA and a CLA mixture lowered high-density lipoprotein cholesterol, although no change was observed in total or low-density lipoprotein cholesterol or triglycerides.¹³ No change in lipids was found in a study with healthy women supplemented with 3.9 g/d CLA.⁵⁵

Consistent with our observations, 4.2 g/d CLA for 1 month was found to increase urinary isoprostanes in men with abdominal obesity.¹⁶ Isoprostanes are produced from peroxidation of lipids, and it was suggested that the increase in isoprostanes might be simply a result of increased fat lipolysis, rather than indicating increased oxidative stress. However, more recent studies have found that 10,12 CLA increases insulin resistance¹³ and plasma CRP.¹⁷ Taken together with our observation of an impairment of endothelial function, it seems highly likely that the increase in isoprostanes does indeed imply an increase in oxidative stress.

Conclusions

CLA supplementation for 12 weeks using the regime used in this study had no significant effect on BMI. Even if this represents a type 2 error, the reduction is at most modest, consistent with previous reports in man. Furthermore, the pattern of fat loss is peripheral rather than central. Importantly, the observed impairment of endothelial function and increase in markers of oxidative stress raise concerns about the widespread use of this agent until further studies demonstrate its cardiovascular safety or otherwise.

References

1. Peiris AN, Sothmann MS, Hoffmann RG, Hennes ML, Wilson CR, Gustafson AB, Kissebah AH. Adiposity, fat distribution, and cardiovascular risk. *Ann Intern Med.* 1989;110:867-872.
2. Isomaa B, Almgren P, Tuomi T, Forsen B, Lahti K, Nissen M, Taskinen MR, Groop L. Cardiovascular morbidity and mortality associated with the metabolic syndrome. *Diabetes Care.* 2001;24:683-689.
3. Bigger JT, Jr, El Sherif T. Polyunsaturated fatty acids and cardiovascular events: a fish tale. *Circulation.* 2001;103:623-625.
4. Hu FB, Stamper MJ, Manson JE, Rimm EB, Wolk A, Colditz GA, Hennekens CH, Willett WC. Dietary intake of alpha-linolenic acid and risk of fatal ischemic heart disease among women. *Am J Clin Nutr.* 1999;69:890-897.
5. Dietary supplementation with n-3 polyunsaturated fatty acids and vitamin E after myocardial infarction: results of the GISSI-Prevenzione trial. Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto miocardico. *Lancet.* 1999;354:447-455.
6. Wahle KWI, Heys SD. Conjugated Linoleic Acids (CLA): Are they Ambrosia or Merely a Panacea. Rowett Research Institute - News & Reviews [1], 4-7. 1-12-2001. 1-12-2001.
7. Lee KW, Lee HJ, Cho HY, Kim YJ. Role of the conjugated linoleic acid in the prevention of cancer. *Crit Rev Food Sci Nutr.* 2005;45:135-144.
8. Kritchewsky D, Tepper SA, Wright S, Tso P, Czarnecki SK. Influence of conjugated linoleic acid (CLA) on establishment and progression of atherosclerosis in rabbits. *J Am Coll Nutr.* 2000;19:472S-477S.
9. Ntambi JM, Choi Y, Park Y, Peters JM, Pariza MW. Effects of conjugated linoleic acid (CLA) on immune responses, body composition and stearoyl-CoA desaturase. *Can J Appl Physiol.* 2002;27:617-628.
10. Blankson H, Stakkestad JA, Fagertun H, Thom E, Wadstein J, Gudmundsen O. Conjugated linoleic acid reduces body fat mass in overweight and obese humans. *J Nutr.* 2000;130:2943-2948.
11. Hargrave KM, Li C, Meyer BJ, Kachman SD, Hartzell DL, Della-Pera MA, Miner JL, Baile CA. Adipose depletion and apoptosis induced by trans-10, cis-12 conjugated linoleic acid in mice. *Obes Res.* 2002;10:1284-1290.
12. Pariza MW, Park Y, Cook ME. Mechanisms of action of conjugated linoleic acid: evidence and speculation. *Proc Soc Exp Biol Med.* 2000;223:8-13.
13. Riserus U, Arner P, Brismar K, Vessby B. Treatment with dietary trans-10,cis-12 conjugated linoleic acid causes isomer-specific insulin resistance in obese men with the metabolic syndrome. *Diabetes Care.* 2002;25:1516-1521.
14. Riserus U, Vessby B, Arnlov J, Basu S. Effects of cis-9,trans-11 conjugated linoleic acid supplementation on insulin sensitivity, lipid peroxidation, and proinflammatory markers in obese men. *Am J Clin Nutr.* 2004;80:279-283.
15. Muloney P, Yeow TP, Mullen A, Nolan H, Roche HM. Conjugated linoleic acid supplementation, insulin sensitivity, and lipoprotein metabolism in patients with type 2 diabetes mellitus. *Am J Clin Nutr.* 2004;80:887-895.
16. Basu S, Riserus U, Turpeinen A, Vessby B. Conjugated linoleic acid induces lipid peroxidation in men with abdominal obesity. *Clin Sci (Colch).* 2000;99:511-516.
17. Riserus U, Basu S, Jovinge S, Fredrikson GN, Arnlov J, Vessby B. Supplementation with conjugated linoleic acid causes isomer-dependent oxidative stress and elevated C-reactive protein: a potential link to fatty acid-induced insulin resistance. *Circulation.* 2002;106:1925-1929.
18. Zambell KL, Keim NL, Van Loan MD, Gale B, Benito P, Kelley DS, Nelson GJ. Conjugated linoleic acid supplementation in humans: effects on body composition and energy expenditure. *Lipids.* 2000;35:777-782.

19. Lohman TG, Roche AF, Martorell R. *Anthropometric Standardization Reference Manual*. Champaign, IL: Human Kinetics; 1991.
20. Nguyen-Duy TB, Nichaman MZ, Church TS, Blair SN, Ross R. Visceral fat and liver fat are independent predictors of metabolic risk factors in men. *Am J Physiol Endocrinol Metab*. 2003;284:E1065-E1071.
21. Goodfellow J, Bellamy MF, Ramsey MW, Jones CJ, Lewis MJ. Dietary supplementation with marine omega-3 fatty acids improves systemic large artery endothelial function in subjects with hypercholesterolemia. *J Am Coll Cardiol*. 2000;35:265-270.
22. Misra A, Vikram NK. Clinical and pathophysiological consequences of abdominal adiposity and abdominal adipose tissue depots. *Nutrition*. 2003;19:457-466.
23. Hwu CM, Fuh JL, Hsiao CF, Wang SJ, Lu SR, Wei MC, Kao WY, Hsiao LC, Ho LT. Waist circumference predicts metabolic cardiovascular risk in postmenopausal Chinese women. *Menopause*. 2003;10:73-80.
24. Lamarche B. Abdominal obesity and its metabolic complications: implications for the risk of ischaemic heart disease. *Coron Artery Dis*. 1998;9:473-481.
25. Tuomilehto J, Lindstrom J, Eriksson JG, Valle TT, Hämäläinen H, Ilanne-Parikka P, Keinänen-Kiukkaanniemi S, Laakso M, Louheranta A, Rastas M, Salminen V, Uusitupa M. Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med*. 2001;344:1343-1350.
26. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med*. 2002;346:393-403.
27. Esposito K, Marfella R, Ciotola M, Di Palo C, Giugliano F, Giugliano G, D'Armiento M, D'Andrea F, Giugliano D. Effect of a Mediterranean-style diet on endothelial dysfunction and markers of vascular inflammation in the metabolic syndrome: a randomized trial. *JAMA*. 2004;292:1440-1446.
28. Vogel RA, Correnti MC, Plotnick GD. The postprandial effect of components of the Mediterranean diet on endothelial function. *J Am Coll Cardiol*. 2000;36:1455-1460.
29. DeLany JP, Blohm F, Truett AA, Scimeca JA, West DB. Conjugated linoleic acid rapidly reduces body fat content in mice without affecting energy intake. *Am J Physiol*. 1999;276:R1172-R1179.
30. Park Y, Albright KJ, Liu W, Storkson JM, Cook ME, Pariza MW. Effect of conjugated linoleic acid on body composition in mice. *Lipids*. 1997;32:853-858.
31. Park Y, Storkson JM, Albright KJ, Liu W, Pariza MW. Evidence that the trans-10,cis-12 isomer of conjugated linoleic acid induces body composition changes in mice. *Lipids*. 1999;34:235-241.
32. Cook ME, Drake B, Jerome D, Pariza MW. The interaction of 9c, 11n9i, 11c and 10t, 12c conjugated linoleic acid on fat deposition in mice. *Fed Am Soc Exp Biol J*. 1999;13:A1023.
33. Riserus U, Berglund L, Vessby B. Conjugated linoleic acid (CLA) reduced abdominal adipose tissue in obese middle-aged men with signs of the metabolic syndrome: a randomised controlled trial. *Int J Obes Relat Metab Disord*. 2001;25:1129-1135.
34. Tsuboyama-Kasaoka N, Takahashi M, Tanemura K, Kim HJ, Tange T, Okuyama H, Kasai M, Ikemoto S, Ezaki O. Conjugated linoleic acid supplementation reduces adipose tissue by apoptosis and develops lipodystrophy in mice. *Diabetes*. 2000;49:1534-1542.
35. Clement L, Poirier H, Niot I, Bocher V, Guerre-Millo M, Krief S, Staels B, Besnard P. Dietary trans-10,cis-12 conjugated linoleic acid induces hyperinsulinemia and fatty liver in the mouse. *J Lipid Res*. 2002;43:1400-1409.
36. Poirier H, Rouault C, Clement L, Niot I, Monnot MC, Guerre-Millo M, Besnard P. Hyperinsulinaemia triggered by dietary conjugated linoleic acid is associated with a decrease in leptin and adiponectin plasma levels and pancreatic beta cell hyperplasia in the mouse. *Diabetologia*. 2005.
37. Yamasaki M, Mansho K, Ogino Y, Kasai M, Tachibana H, Yamada K. Acute reduction of serum leptin level by dietary conjugated linoleic acid in Sprague-Dawley rats. *J Nutr Biochem*. 2000;11:467-471.
38. Belury MA, Mahon A, Banni S. The conjugated linoleic acid (CLA) isomer, t10c12-CLA, is inversely associated with changes in body weight and serum leptin in subjects with type 2 diabetes mellitus. *J Nutr*. 2003;133:2578-2608.
39. Houseknecht KL, Vanden Heuvel JP, Moya-Camarena SY, Portocarrero CP, Peck LW, Nickel KP, Belury MA. Dietary conjugated linoleic acid normalizes impaired glucose tolerance in the Zucker diabetic fatty fa/fa rat. *Biochem Biophys Res Commun*. 1998;244:678-682.
40. Yotsumoto H, Hara E, Naka S, Adiof R, Emken E, Yanagita T. 10trans, 12cis-linoleic acid reduces apolipoprotein B secretion in HepG2 cells. *Food Res Int*. 1999;31:403-409.
41. Brown RM, Boyson MS, Jensen SS, Morrison RF, Sterkson J, Lea-Curie R, Pariza M, Mandrup S, McIntosh MK. Isomer-specific regulation of metabolism and PPARgamma signaling by CLA in human preadipocytes. *J Lipid Res*. 2003;44:1287-1300.
42. Wargent E, Sennitt MV, Stocker C, Mayes AE, Brown L, O'Dowd J, Wang S, Einerhand AW, Mohede I, Arch JR, Cawthorne MA. Prolonged treatment of genetically obese mice with conjugated linoleic acid improves glucose tolerance and lowers plasma insulin concentration: possible involvement of PPAR activation. *Lipids Health Dis*. 2005;4:3.
43. Granlund L, Juvel LK, Pedersen JJ, Nebb HJ. Trans-10, cis-12-conjugated linoleic acid prevents triacylglycerol accumulation in adipocytes by acting as a PPARgamma modulator. *J Lipid Res*. 2003;44:1441-1452.
44. Kang K, Liu W, Albright KJ, Park Y, Pariza MW. trans-10,cis-12 CLA inhibits differentiation of 3T3-L1 adipocytes and decreases PPARgamma expression. *Biochem Biophys Res Commun*. 2003;303:795-799.
45. Changhua L, Jindong Y, Defa L, Lidian Z, Shiyun Q, Jianjun X. Conjugated linoleic acid attenuates the production and gene expression of proinflammatory cytokines in weaned pigs challenged with lipopolysaccharide. *J Nutr*. 2005;135:239-244.
46. Smith SA. Central role of the adipocyte in the insulin-sensitising and cardiovascular risk modifying actions of the thiazolidinediones. *Biochimie*. 2003;85:1219-1230.
47. Ryysy L, Hakkinen AM, Goto T, Vehkavaara S, Westerbacka J, Halavaara J, Yki-Jarvinen H. Hepatic fat content and insulin action on free fatty acids and glucose metabolism rather than insulin absorption are associated with insulin requirements during insulin therapy in type 2 diabetic patients. *Diabetes*. 2000;49:749-758.
48. Bots ML, Westerink J, Rabelink TJ, de Koning HJ. Assessment of flow-mediated vasodilation (FMD) of the brachial artery: effects of technical aspects of the FMD measurement on the FMD response. *Eur Heart J*. 2005;26:363-368.
49. Smith JC, Evans LM, Wilkinson I, Goodfellow J, Cockcroft JR, Scanlon MF, Davies JS. Effects of GH replacement on endothelial function and large-artery stiffness in GH-deficient adults: a randomized, double-blind, placebo-controlled study. *Clin Endocrinol (Oxf)*. 2002;56:493-501.
50. Evans M, Anderson RA, Smith JC, Khan N, Graham JM, Thomas AW, Morris K, Deely D, Frencheaux MP, Davies JS, Raes A. Effects of insulin lispro and chronic vitamin C therapy on postprandial lipaemia, oxidative stress and endothelial function in patients with type 2 diabetes mellitus. *Eur J Clin Invest*. 2003;33:231-238.
51. Quyyumi AA. Prognostic value of endothelial function. *Am J Cardiol*. 2003;91:19H-24H.
52. Wallace TM, Levy JC, Matthews DR. Use and abuse of HOMA modeling. *Diabetes Care*. 2004;27:1487-1495.
53. Truitt A, McNeill G, Vanderhoek JY. Antiplatelet effects of conjugated linoleic acid isomers. *Biochim Biophys Acta*. 1999;1438:239-246.
54. Benito P, Nelson GJ, Kelley DS, Bartolini G, Schmidt PC, Simon V. The effect of conjugated linoleic acid on platelet function, platelet fatty acid composition, and blood coagulation in humans. *Lipids*. 2001;36:221-227.
55. Benito P, Nelson GJ, Kelley DS, Bartolini G, Schmidt PC, Simon V. The effect of conjugated linoleic acid on plasma lipoproteins and tissue fatty acid composition in humans. *Lipids*. 2001;36:229-236.

ADVERSE CARDIOVASCULAR AND CENTRAL NERVOUS SYSTEM EVENTS ASSOCIATED WITH DIETARY SUPPLEMENTS CONTAINING EPHEDRA ALKALOIDS

CHRISTINE A. HALLER, M.D., AND NEAL L. BENOWITZ, M.D.

ABSTRACT

Background Dietary supplements that contain ephedra alkaloids (sometimes called ma huang) are widely promoted and used in the United States as a means of losing weight and increasing energy. In the light of recently reported adverse events related to use of these products, the Food and Drug Administration (FDA) has proposed limits on the dose and duration of use of such supplements. The FDA requested an independent review of reports of adverse events related to the use of supplements that contained ephedra alkaloids to assess causation and to estimate the level of risk the use of these supplements poses to consumers.

Methods We reviewed 140 reports of adverse events related to the use of dietary supplements containing ephedra alkaloids that were submitted to the FDA between June 1, 1997, and March 31, 1999. A standardized rating system for assessing causation was applied to each adverse event.

Results Thirty-one percent of cases were considered to be definitely or probably related to the use of supplements containing ephedra alkaloids, and 31 percent were deemed to be possibly related. Among the adverse events that were deemed definitely, probably, or possibly related to the use of supplements containing ephedra alkaloids, 47 percent involved cardiovascular symptoms and 18 percent involved the central nervous system. Hypertension was the single most frequent adverse effect (17 reports), followed by palpitations, tachycardia, or both (13); stroke (10); and seizures (7). Ten events resulted in death, and 13 events produced permanent disability, representing 26 percent of the definite, probable, and possible cases.

Conclusions The use of dietary supplements that contain ephedra alkaloids may pose a health risk to some persons. These findings indicate the need for a better understanding of individual susceptibility to the adverse effects of such dietary supplements. (N Engl J Med 2000;343:1833-8.)

©2000, Massachusetts Medical Society.

DIETARY supplements that contain ephedra alkaloids (also known as ma huang) and guarana-derived caffeine are widely consumed in the United States for purposes of weight reduction and energy enhancement. A number of reports of adverse reactions to dietary supplements that contain ephedra alkaloids, some of which resulted in permanent injury or death, have appeared in the medical literature.¹⁻⁶ In response to

growing concern about the safety of ephedra alkaloids in dietary supplements, the Food and Drug Administration (FDA) requested an independent review of reports of adverse events related to the use of ephedra alkaloids to assess causation and determine the level of risk these products pose to consumers.

We conducted an in-depth review of 140 reports of adverse events involving dietary supplements containing ephedra alkaloids that were submitted to the FDA between June 1, 1997, and March 31, 1999, and applied a standardized rating system for assessing causation. We also evaluated factors that might increase the risk to consumers and the adequacy of warnings about potential risks included on product labels. The full report of our review of adverse events is available elsewhere.⁷ Here, we summarize our findings.

METHODS

The objective of the review was to determine the likelihood that ephedra alkaloids (which were usually combined with caffeine) caused the reported adverse events on the basis of the information provided in the FDA MedWatch report, along with supplemental medical records. We independently reviewed each of the 140 cases. Causation was assessed according to the criteria described by Blanc et al.⁸ and included an evaluation of the timing of the event in relation to the dose and duration of use of a product; an assessment of the pattern of response to determine whether it constituted a recognized reaction to the substance on the basis of previous reports of ephedrine or similar stimulants in the medical literature; and a determination of the contribution of any underlying diseases or medical conditions.

In general, we defined an adverse event as definitely related to the use of supplements containing ephedra alkaloids only if the symptoms recurred with the reintroduction of ephedra alkaloids or when the onset of symptoms coincided with the expected peak plasma concentration of the drug and resolved within an interval that was consistent with the expected duration of the effect of ephedrine. An adverse event was defined as probably related to the use of supplements containing ephedra alkaloids when the majority of the evidence supported the existence of a causal link but one or more aspects of the case, such as time since the last dose, were unknown or there was a minor inconsistency in the supporting evidence, such as a low reported dose. An adverse event was designated as possibly related to the use of supplements containing ephedra alkaloids when it was equally likely that the adverse event was not related to the use of ephedra alkaloids; for example, in the case of effects that have not been reported in the literature in association with ephedra alkaloids but that are pharmacologically plausible. Reports of ad-

From the Division of Clinical Pharmacology and Experimental Therapeutics, Departments of Medicine and Biopharmaceutical Sciences, University of California, San Francisco, and the California Poison Control System, San Francisco Division — both in San Francisco. Address reprint requests to Dr. Benowitz at the Division of Clinical Pharmacology and Experimental Therapeutics, University of California, San Francisco, Box 1220, San Francisco, CA 94143-1220, or at nbeno@itsa.ucsf.edu.

verse events that included scant medical history and incomplete information about the product involved were usually considered to have insufficient information to be assessed. This category was reserved for events in which the evidence was not substantial enough to consider them as possibly related to the use of supplements containing ephedra alkaloids. Adverse events were defined as probably unrelated to the use of supplements containing ephedra alkaloids if the evidence that ephedra alkaloids were the cause was weak or if the likelihood was strong that there was some other cause, either medical or toxicologic. When the scientific evidence or course of events was highly inconsistent with the known effects of ephedra alkaloids, the event was considered definitely unrelated; for example, in the case of symptoms that persisted long after the use of ephedra alkaloids had been discontinued or in the case of symptoms that had no association with the known pharmacodynamic effects of ephedra alkaloids. However, the event was considered related if the patient had a preexisting condition such as hypertension that could have been aggravated by the use of ephedra alkaloids and if the pattern of use met the criteria for causation.

In determining the likelihood of a causal link, we evaluated aspects of the medical history, dietary patterns, and social habits as possible contributing or causative factors. For example, we noted when events occurred while patients were fasting or in conjunction with high intakes of caffeine. We recognized that in the case of adverse events that were most likely not related to the use of supplements containing ephedra alkaloids one or more of the other ingredients in the supplement may have been causally related to the event.

RESULTS

Features of the Cases

The age and sex of the users of products containing ephedra alkaloids and the reported reasons for use are shown in Table 1. Although the labels of most such products state that they are not intended for use by persons less than 18 years of age, adverse events were recorded in at least 10 persons under this age. The youngest was 15 years old. Overall, 43 cases (31 percent) were considered to be definitely or probably related to the use of supplements containing ephedra alkaloids, 24 cases (17 percent) were considered to be unrelated to the use of such supplements, 44 cases (31 percent) were deemed possibly related to the use of such supplements, and in 29 cases (21 percent) the information provided was insufficient to assess causation. The types of adverse events that were definitely or probably related to the use of supplements containing ephedra alkaloids and those that were possibly related are summarized in Table 2.

Cardiovascular symptoms made up 47 percent of the adverse events that were definitely, probably, or possibly related to the use of supplements containing ephedra alkaloids. Hypertension was the single most frequent adverse effect, followed by palpitations, tachycardia, or both. Eighteen percent of related and possibly related adverse events involved the central nervous system. Strokes ($n=10$) and seizures ($n=7$) were the most frequent type of central nervous system event reported. The clinical outcomes of the definite, probable, and possible cases are listed in Table 3.

Ten events resulted in death (including 1 neonatal death and 1 fetal death), and 13 events resulted in permanent impairment, which represented 26 percent of

TABLE 1. AGE AND SEX OF USERS OF SUPPLEMENTS CONTAINING EPHEDRA ALKALOIDS AND REASONS FOR USE, ACCORDING TO ADVERSE-EVENT REPORTS.

VARIABLE	No. of Users (%)
Age	
<18 yr	10 (7)
18–29 yr	34 (24)
30–45 yr	45 (32)
>45 yr	32 (23)
Unknown	19 (14)
Sex	
Male	56 (40)
Female	84 (60)
Reason for use	
Weight loss	83 (59)
To increase athletic performance	23 (16)
To increase energy	9 (6)
Unknown	24 (17)
Intentional misuse	1 (1)

the definite, probable, and possible cases. Nine serious adverse events occurred in persons who were taking relatively low doses of ephedra alkaloids (range, 12 to 36 mg per day) and who had no important medical risk factors. Features of the definite or probable and possible cases that resulted in death or permanent impairment or that necessitated substantial medical intervention are given in Tables 4 and 5, respectively.

Of the sudden catastrophic cerebrovascular and cardiovascular events, 11 occurred in previously healthy persons. Some of these cases, which were definitely or probably related to the use of supplements containing ephedra alkaloids, are described in detail in the following sections.

Examples of Severe Cerebrovascular Adverse Events

Patient 1

Patient 1 was a healthy 35-year-old woman who had taken aerobic-exercise classes for several years without incident (Table 4). In July 1997, she began taking one capsule of Shape-Fast Plus (according to the label, each capsule contained 15 mg of ephedra alkaloids and 40 mg of caffeine) three times a day before meals for weight loss; she was taking no other medications. She had been taking the product for one week when she collapsed during an aerobics class. Bystanders observed that her arms and legs were flexing and tensing. In the emergency department, her blood pressure was 110/38 mm Hg and the heart rate was 104 beats per minute. A computed tomographic scan of the head showed a subarachnoid hemorrhage. Cerebral angiography showed no evidence of a vascular aneurysm. A urine toxicology screen was positive for amphetamine, a result presumed to reflect a cross-reaction with the ephedrine and therefore to be false positive.

TABLE 2. TYPES OF ADVERSE EVENTS THAT WERE DEFINITELY, PROBABLY, OR POSSIBLY RELATED TO THE USE OF SUPPLEMENTS CONTAINING EPHEDRA ALKALOIDS.

ADVERSE EVENT	DEFINITELY OR PROBABLY RELATED (N=43)	POSSIBLY RELATED (N=44)	TOTAL (N=87)
	no. of events (%)		
Cardiovascular			
Hypertension	10 (21)	7 (14)	17 (17)
Palpitations, tachycardia, or both	8 (17)	5 (10)	13 (13)
Arrhythmia	3 (6)	3 (6)	6 (6)
Myocardial infarction	2 (4)	0	2 (2)
Cardiac arrest or sudden death	5 (10)	3 (6)	8 (8)
Central nervous system			
Stroke	4 (8)	6 (12)	10 (10)
Transient ischemic attack	1 (2)	0	1 (1)
Seizure	1 (2)	6 (12)	7 (7)
Other	14 (29)	20 (40)	34 (35)
Total no. of events*	48	50	98

*The total number of events exceeds the total number of cases, since some cases involved more than one adverse event.

Neurogenic pulmonary edema rapidly developed, necessitating endotracheal intubation and mechanical ventilation. Electrocardiographic findings and cardiac-enzyme levels were consistent with the occurrence of a small myocardial infarction. The treating cardiologist and neurologist thought that ephedrine induced the subarachnoid hemorrhage. The finding of amphetamine on the urine toxicology test supports the presence of ephedrine at the time of the event. Laboratory analysis of the supplement determined that the ephedrine content was 12.0 mg per capsule. At that time, the FDA's recommendation was a maximal dose of 8 mg per serving.⁷

Patient 10

Patient 10 was an apparently healthy 39-year-old man who experienced numbness of the right arm and leg on March 17, 1998, 90 minutes after drinking Ultimate Orange, which according to the label contained 415 mg of ma huang (ephedra alkaloids) per serving as well as guarana (a source of caffeine), and 5 minutes after running 3 miles (4.8 km) (Table 4). He also regularly took multivitamins and amino acid supplements, but no other medications. On presentation at a nearby hospital, his blood pressure was 140/78 mm Hg and his pulse was 60 beats per minute. A computed tomographic scan of the head revealed a left-sided intrathalamic hemorrhage. Cerebral angiography showed no evidence of vascular anomalies. The patient had gradual clinical improvement, and his symptoms resolved except for persistent sensory loss on the right side of his face. Chemical analy-

TABLE 3. CLINICAL OUTCOMES OF ADVERSE EVENTS THAT WERE DEFINITELY, PROBABLY, OR POSSIBLY RELATED TO THE USE OF SUPPLEMENTS CONTAINING EPHEDRA ALKALOIDS.

OUTCOME	DEFINITELY OR PROBABLY RELATED (N=43)	POSSIBLY RELATED (N=44)	TOTAL (N=87)
	no. of events (%)		
Death	3 (7)	7 (16)	10 (11)
Permanent impairment	7 (16)	6 (14)	13 (15)
Ongoing medical treatment	4 (9)	4 (9)	8 (9)
Full recovery	29 (67)	13 (30)	42 (48)
Unknown	0	14 (32)	14 (16)

sis of the Ultimate Orange product confirmed the presence of ephedrine, as well as of pseudoephedrine, norephedrine, and norpseudoephedrine.

Examples of Severe Cardiovascular Adverse Events

Patient 2

Patient 2 was a 22-year-old man with a history of asthma who collapsed while lifting weights at a gym on March 31, 1998 (Table 4). His medications included theophylline (Theo-Dur; 300 mg twice daily), albuterol (Ventolin; administered as necessary through a metered-dose inhaler), and a combination of chlorpheniramine maleate, phenylephrine hydrochloride, and phenylpropanolamine hydrochloride (Atrohist Plus SR). According to friends, he had consumed one 18-oz bottle of Ripped Force (which is listed as containing 20 mg of ephedrine alkaloids, 100 mg of caffeine, 250 mg of L-carnitine, and 240 µg of chromium) before working out and was regularly drinking three bottles of Ripped Force per day. He also took creatine and protein supplements. Witnesses reported that he had a seizure. Paramedics initially found him apneic and in ventricular fibrillation. He was successfully resuscitated. Computed tomography of the head showed cerebral edema but no hemorrhage or masses. An initial electrocardiogram showed atrial flutter, which subsequently converted to sinus rhythm. An echocardiogram revealed mild left ventricular hypertrophy. The plasma theophylline level was 11 µg per milliliter (therapeutic range, 10 to 20), and urinalysis revealed 12 µg of ephedrine per milliliter, 0.38 µg of pseudoephedrine per milliliter, and 0.41 µg of phenylpropanolamine per milliliter. The treating cardiologist thought that the combination of ephedra alkaloids and caffeine in Ripped Force and the theophylline and albuterol medications caused a ventric-

TABLE 4. OUTCOME IN 11 PATIENTS WITH ADVERSE EVENTS THAT WERE DEFINITELY OR PROBABLY RELATED TO THE USE OF SUPPLEMENTS CONTAINING EPHEDRA ALKALOIDS.

PATIENT NO.	AGE (YR)/SEX	NAME OF SUPPLEMENT	ESTIMATED DAILY DOSE OF EPHEDRA ALKALOIDS mg	DURATION OF USE	ADVERSE EVENT	OUTCOME	PREEXISTING CONDITIONS OR CONCURRENT RISKS
1	35/F	Shape-Fast Plus	45	1 wk	Subarachnoid hemorrhage	Permanent disability	None
2	22/M	Ripped Force	20–60	Unknown	Arrhythmia, cardiac arrest	Permanent disability	Asthma
3	28/F	Herbalife's Thermo-jetics	21	1 day	Cardiac arrest	Permanent disability	None
4	43/M	Ripped Fuel	60	7 mo	Cardiac arrest	Death	Family history of coronary artery disease
5	37/F	Metabolife 350	36	1 wk	Severe hypertension, cardiac arrest, hypokalemia	Death	None
6	59/F	OmniTrim Extra Vitamin-Fortified tea	36	3 wk	Acute myocardial infarction	Coronary bypass surgery	Hypertension
7	38/M	Ripped Fuel	20	1 yr	Arrhythmia, cardiac arrest	Death	None
8	47/F	Total Control	44–66	9 mo	Hypertension, bilateral lacunar infarctions	Permanent disability	Concomitant ingestion of caffeine and ethanol
9	29/M	Ultimate Orange	30	2 wk	Stroke	Permanent disability	Concomitant use of dehydroepiandrosterone and androstenedione
10	39/M	Ultimate Orange	Unknown	Unknown	Hemorrhagic stroke	Permanent disability	None
11	47/M	Purple Blast	Unknown	3 wk	Hemorrhagic stroke	Permanent disability	Possible hypertension

ular arrhythmia that resulted in cardiac arrest. The patient suffered anoxic encephalopathy and remained in a vegetative state for several weeks. After one month in an acute care facility and six weeks at a rehabilitation facility, he was discharged with substantial residual neurologic impairment.

Patient 7

Patient 7 was an apparently healthy 38-year-old man who had been taking two capsules of Ripped Fuel (according to the label each capsule contains 10 mg of ephedrine and 100 mg of caffeine) each morning for one year as directed on the product label (Table 4). On June 6, 1996, he took his usual dose along with a cup of coffee and went jogging for 20 minutes. After returning home, he was talking with his family when he suddenly collapsed and appeared to have a tonic-clonic seizure. He had not reported any symptoms before collapsing. He was in full cardiac arrest when paramedics arrived and could not be resuscitated. Autopsy showed mild cardiomegaly with four-chamber dilatation and coronary artery disease, with narrowing of 50 to 75 percent in four vessels. The cause of death was acute arrhythmia resulting from atherosclerotic cardiovascular disease. Subsequent toxicology testing showed blood levels of 110 ng of ephedrine per milliliter (the therapeutic range used for bronchodilation is 20 to 80).⁹ An addendum

to the autopsy report included the comment, "ephedrine is a stimulant medication, and as such may have contributed to a fatal arrhythmia in the decedent."

DISCUSSION

Ephedrine and related alkaloids have been associated with adverse cardiovascular events, including acute myocardial infarction, severe hypertension, myocarditis, and lethal cardiac arrhythmias.^{10,11} Constriction of coronary arteries and, in some cases, vasospasm are believed to be the mechanisms of myocarditis and myocardial infarction. The adrenergic effects of ephedrine shorten cardiac refractory periods, permitting the development of reentrant cardiac arrhythmias. Ephedrine can predispose patients to both hemorrhagic and ischemic stroke.¹² Subarachnoid hemorrhage is thought to be a result of the hypertensive action of ephedrine, which can be short lived, or of cerebral vasculitis, which has been described in association with a variety of sympathomimetic drugs.^{13,14} Thrombotic stroke is presumably related to vasoconstriction of large cerebral arteries, which leads to local thrombosis as a result of stasis and sympathomimetic-induced platelet activation.

Caffeine is present in many products that contain ephedra alkaloids, and those who take these products might also be consuming considerable quantities of caffeine in coffee, tea, and soft drinks. Caffeine is like-

TABLE 5. OUTCOME IN 15 PATIENTS WITH ADVERSE EVENTS THAT WERE POSSIBLY RELATED TO THE USE OF SUPPLEMENTS CONTAINING EPHEDRA ALKALOIDS.

PATIENT No.	AGE (yr)/SEX	NAME OF SUPPLEMENT	ESTIMATED DAILY DOSE OF EPHEDRA ALKALOIDS mg	DURATION OF USE	ADVERSE EVENT	OUTCOME	PREEXISTING CONDITIONS OR CONCURRENT RISKS
1	46/M	Diet Fuel	Unknown	5-6 mo	Stroke	Death	None
2	22/M	Ripped Fuel	Unknown	Unknown	Hyperthermia, abnormal electrolyte levels, cardiac arrest	Death	None
3	64/F	Fit America Natural Weight Control Aid	Unknown	2 mo	Atrial fibrillation, stroke	Permanent disability	Hypertension, transient ischemic attack
4	47/F	Per-Form Dieter's Natural Tea	Unknown	6 mo	Rhabdomyolysis, hydronephrosis, hypokalemia	Prolonged hospital care	None
5	64/F	Shape-Fast	20	Unknown	Hemorrhagic stroke	Permanent disability	None
6	34/M	Herbalife's Thermojetics	Unknown	>3 wk	Atrial flutter, renal failure, hypokalemia, rhabdomyolysis	Death	None
7	32/F	Ripped Fuel	20	4 yr	Premature delivery (34 wk of gestation)	Death of neonate	Smoking
8	29/M	Ultimate Nutrition Product Ma Huang	Unknown	6-7 mo	Stroke	Permanent disability	None
9	15/F	Ripped Fuel	Unknown	2-3 wk	Arrhythmia, cardiac arrest	Death	None
10	41/F	Diet-Phen	12	1 mo	Hypertension, multiple brain-stem infarcts	Permanent disability	Possible hypertension
11	22/F	Magic Herb	72	3 mo	Spontaneous abortion at 9 wk	Death of fetus	None
12	43/F	Metabolife 356	12	6 mo	Severe hypertension, hemorrhagic stroke	Permanent disability	None
13	18/M	Ultimate Orange	Unknown	Unknown	Seizure, hemorrhagic stroke	Death	None
14	61/F	Metabolife 356	24; increased to 60	1 mo	Hypertension, unstable angina	Coronary bypass surgery	Asthma
15	26/M	Ripped Fuel	20-60	3 yr	Status epilepticus, hypokalemia	Permanent disability	None

ly to enhance the cardiovascular and central nervous system effects of ephedrine. Caffeine acts by competitively antagonizing the receptors for adenosine, a hormone released by endothelial cells that dilates blood vessels.¹⁵ By inhibiting adenosine-mediated dilatation of blood vessels, caffeine constricts blood vessels and may increase blood pressure in persons prone to hypertension. Caffeine also augments the release of catecholamines, an effect that, when combined with that of ephedrine, could lead to increased stimulation of the central nervous system and cardiovascular system.¹⁶

Phenylpropanolamine, another ephedrine alkaloid, was marketed with caffeine in various weight-reducing aids until 1983, when the combination was banned by the FDA after numerous reports of adverse effects. Several studies have shown that caffeine and phenylpropanolamine have an additive effect on blood pressure.¹⁷ These interactions between phenylpropanolamine and caffeine support the idea that the combination of ephedrine and caffeine in a dietary supplement could increase the risk of adverse effects.

The quantity of ephedrine in dietary supplements,

as reported on package labels, is typically about 20 mg per serving, and the usual dose frequency is two to three times per day. These products may contain larger or smaller amounts of ephedra alkaloids than are listed on the product label. For example, 11 of 20 supplements tested by Gurley et al.¹⁸ either failed to list the alkaloid content on the label or had more than a 20 percent difference between the amount listed on the label and the actual amount.

Often, the dose of ephedrine that was associated with an adverse event was less than a typical dose of ephedrine used for bronchodilation (25 to 50 mg). Experimental studies show that ephedrine has only moderate effects on heart rate and blood pressure at these doses.^{19,20} The discrepancy between such data and our findings of serious adverse events reported with the use of dietary supplements containing ephedra alkaloids may be due to individual susceptibility, the additive stimulant effects of caffeine, the variability in the contents of pharmacologically active chemicals in the products, or preexisting medical conditions.

Many of the cases we reviewed involved side effects

such as anxiety, tremulousness, insomnia, palpitations, and personality changes that are well known to occur with the use of stimulant drugs. When ephedrine is used for medical purposes, these types of reactions are considered side effects and must be included in the assessment of risks and benefits. In fact, ephedrine is rarely prescribed today for medical purposes, because newer drugs have more specific actions and fewer side effects. The risks of taking ephedra alkaloids as a dietary supplement, however, are difficult to justify because the alkaloids have no demonstrated benefit. Unlike vitamins and minerals, ephedra alkaloid supplements are not essential for proper nutrition. People who take these products to increase their exercise capacity or to lose weight place themselves at risk without a substantial likelihood of benefit.

A limitation to the use of reports of adverse events as an indicator of a product's safety is that the number of people at risk for the event is unknown. Manufacturers of dietary supplements that contain ephedra alkaloids reported that 3 billion servings were sold in 1999.²¹ The number of servings that were actually consumed is difficult to determine. Assuming that the products were consumed as directed — three doses per day for 12 weeks — then approximately 12 million people used these supplements in 1999.

Another limitation is that adverse events are known to be underreported. Several studies have shown that spontaneous reporting of adverse events to MedWatch is not routine, and the rate of reporting may be less than 15 percent.^{22,23} The frequency of reports of adverse reactions to herbal products is thought to be even lower.²⁴ Therefore, the frequency of serious adverse events associated with the use of supplements containing ephedra alkaloids cannot be precisely determined with the use of current reporting mechanisms.

Because of the severity of the adverse events that we reviewed and, in particular, the occurrence of events that caused permanent disability and death, we conclude that dietary supplements that contain ephedra alkaloids pose a serious health risk to some users. Although the incidence of serious adverse effects cannot be determined from our analysis, our findings arouse concern about the risks of these products, given that they have no scientifically established benefits. Our findings indicate the need for a better understanding of the determinants of individual susceptibility to the serious adverse effects of dietary supplements containing ephedra alkaloids so that appropriate dosing guidelines and warnings can be devised.

Supported by funds from the Food and Drug Administration and by grants (DA02277 and DA12393) from the National Institute on Drug

Abuse, National Institutes of Health. Dr. Haller was supported by a Veterans Affairs Medical Toxicology fellowship and by a National Research Service Award (T32GM07546) from the National Institutes of Health.

We are indebted to Dr. Raymond Woosley for his critical review of the manuscript and to Kaye Welch for editorial assistance.

REFERENCES

1. Adverse events associated with ephedrine-containing products — Texas, December 1993–September 1995. *JAMA* 1996;276:1711–2.
2. Theoharides TC. Sudden death of a healthy college student related to ephedrine toxicity from a ma huang-containing drink. *J Clin Psychopharmacol* 1997;17:437–9.
3. Josefson D. Herbal stimulant causes US deaths. *BMJ* 1996;312:1378–9.
4. Zahn KA, Li RL, Pursell RA. Cardiovascular toxicity after ingestion of "herbal ecstasy." *J Emerg Med* 1999;17:289–91.
5. Zaacks SM, Klein L, Tan CD, Rodriguez ER, Leikin JB. Hypersensitivity myocarditis associated with ephedra use. *J Toxicol Clin Toxicol* 1999;37:485–9.
6. Vahedi K, Domingo V, Amencio R, Boussier MG. Ischaemic stroke in a sportsman who consumed MaHuang extract and creatine monohydrate for body building. *J Neurol Neurosurg Psychiatry* 2000;68:112–3.
7. Dietary supplements containing ephedrine alkaloids. FDA docket no. 00N-1200. Rockville, Md.: Food and Drug Administration, 2000. (See www.accessdata.fda.gov/scripts/cdrh/cdrms/index.cfm.)
8. Blanc S, Leuenberger P, Berger JR, Brooke EM, Schelling JL. Judgments of trained observers on adverse drug reactions. *Clin Pharmacol Ther* 1979;25:493–8.
9. Costello JR, May CS, Paterson JW, Pickup ME. Pharmacokinetics of ephedrine in asthmatics receiving acute and chronic treatment. *Br J Clin Pharmacol* 1975;2:180–1, abstract.
10. Wiener I, Tilkian AG, Palazzolo M. Coronary artery spasm and myocardial infarction in a patient with normal coronary arteries: temporal relationship to pseudoephedrine ingestion. *Cathet Cardiovasc Diagn* 1990;20:51–3.
11. To LB, Sangster JF, Rampling D, Cammings I. Ephedrine-induced cardiomyopathy. *Med J Aust* 1980;2:35–6.
12. Bruno A, Nolte KB, Chapin J. Stroke associated with ephedrine use. *Neurology* 1993;43:1313–6.
13. Fallis RJ, Fisher M. Cerebral vasculitis and hemorrhage associated with phenylpropanolamine. *Neurology* 1985;35:405–7.
14. Wooten MR, Khangure MS, Murphy MJ. Intracerebral hemorrhage and vasculitis related to ephedrine abuse. *Ann Neurol* 1983;13:337–40.
15. Benowitz NL. Clinical pharmacology of caffeine. *Annu Rev Med* 1990;41:277–88.
16. Robertson D, Frölich JC, Carr RK, et al. Effects of caffeine on plasma renin activity, catecholamines and blood pressure. *N Engl J Med* 1978;298:181–6.
17. Brown NJ, Ryder D, Brauch RA. A pharmacodynamic interaction between caffeine and phenylpropanolamine. *Clin Pharmacol Ther* 1991;50:363–71.
18. Gurley BJ, Gardner SF, Hubbard MA. Content versus label claims in ephedra-containing dietary supplements. *Am J Health Syst Pharm* 2000;57:963–9.
19. Drew CD, Knight GT, Hughes DT, Bush M. Comparison of the effects of D-(–)-ephedrine and L-(+)-pseudoephedrine on the cardiovascular and respiratory systems in man. *Br J Clin Pharmacol* 1978;6:221–5.
20. White LM, Gardner SF, Gurley BJ, Marx MA, Wang PL, Estes M. Pharmacokinetics and cardiovascular effects of ma-huang (Ephedra sinica) in normotensive adults. *J Clin Pharmacol* 1997;37:116–22.
21. Dietary supplement market view. Chevy Chase, Md.: FDC Reports, August 2000.
22. Chyka PA, McCommon SW. Reporting of adverse drug reactions by poison control centres in the US. *Drug Saf* 2000;23:87–93.
23. Spontaneous adverse event reports. (See <http://www.fda.gov/ohrtm/dockets/ac/99/saides/3558slc/1slid003.htm>.)
24. Barnes J, Mills SY, Abbot NC, Willoughby M, Ernst E. Different standards for reporting ADRs to herbal remedies and conventional OTC medicines: face-to-face interviews with 515 users of herbal remedies. *Br J Clin Pharmacol* 1998;45:496–506.

X. RELATED PROCEEDINGS APPENDIX

No decisions have been rendered by a court or the Board in any proceeding identified pursuant to paragraph (c)(1)(ii) C.F.R. §41.37.

XI. CONCLUSION

For the foregoing reasons, Appellants respectfully submit that the Examiner's rejection of Claims 1-3, 7 and 9 is erroneous. Reversal of the rejections is respectfully requested. Appellants request that the Board render a decision as to the allowability of the Claims.

Respectfully submitted,

Dated: October 27, 2008 /John Mitchell Jones/

J. Mitchell Jones
Registration Number 44,174
Casimir Jones SC
440 Science Dr. Ste. 203
Madison WI 53711